

The Product of Two Eulerian Polynomials Author(s): L. Carlitz Source: *Mathematics Magazine*, Vol. 36, No. 1 (Jan., 1963), pp. 37-41 Published by: <u>Mathematical Association of America</u> Stable URL: <u>http://www.jstor.org/stable/2688134</u> Accessed: 23/01/2015 08:40

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at http://www.jstor.org/page/info/about/policies/terms.jsp

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



Mathematical Association of America is collaborating with JSTOR to digitize, preserve and extend access to *Mathematics Magazine*.

http://www.jstor.org

THE PRODUCT OF TWO EULERIAN POLYNOMIALS

L. CARLITZ, Duke University

The Bernoulli and Euler polynomials can be defined by means of

$$\frac{te^{xt}}{e^t-1} = \sum_{m=0}^{\infty} B_m(x) \frac{t^m}{m!}, \qquad \frac{2e^{xt}}{e^t+1} = \sum_{m=0}^{\infty} E_m(x) \frac{t^m}{m!}.$$

The formula

(1)
$$B_m(x)B_n(x) = \sum_r \left\{ \binom{m}{2r} n + \binom{n}{2r} m \right\} \frac{B_{2r}B_{m+n-2r}(x)}{m+n-2r}$$

is proved in Nielsen's book [3, p. 75]; a different proof occurs in [2]. Nielsen also obtains similar formulas for

 $E_m(x)E_n(x)$ and $E_m(x)B_n(x)$.

The Eulerian polynomial $H_m(x|\lambda)$ can be defined by means of

(2)
$$\frac{(1-\lambda)e^{xt}}{e^t-\lambda} = \sum_{m=0}^{\infty} H_m(x \mid \lambda) \frac{t^m}{m!};$$

for properties of $H_m(x|\lambda)$ see for example [1]. Since

$$H_m(x \mid -1) = E_m(x),$$

it may be of interest to get a formula for the product of two Eulerian polynomials.

We assume that $\alpha \neq 1$, $\beta \neq 1$, $\alpha \beta \neq 1$. It follows from (2) that

$$\begin{split} &\sum_{m,n=0}^{\infty} H_m(x \mid \alpha) H_n(x \mid \beta) \frac{u^m v^n}{m!n!} = \frac{(1-\alpha)e^{xu}}{e^u - \alpha} \frac{(1-\beta)e^{xv}}{e^v - \beta} \\ &= \frac{(1-\alpha)(1-\beta)}{1-\alpha\beta} \frac{(1-\alpha\beta)e^{x(u+v)}}{e^{u+v} - \alpha\beta} \frac{e^{u+v} - \alpha\beta}{(e^u - \alpha)(e^v - \beta)} \\ &= \frac{(1-\alpha)(1-\beta)}{1-\alpha\beta} \frac{(1-\alpha\beta)e^{x(u+v)}}{e^{u+v} - \alpha\beta} \left\{ 1 + \frac{\alpha}{e^u - \alpha} + \frac{\beta}{e^v - \beta} \right\} \\ &= \frac{1}{1-\alpha\beta} \sum_{m,n=0}^{\infty} H_{m+n}(x \mid \alpha\beta) \frac{u^m v^n}{m!n!} \\ &\cdot \left\{ (1-\alpha)(1-\beta) + \alpha(1-\beta) \sum_{r=0}^{\infty} H_r[\alpha] \frac{u^r}{r!} + \beta(1-\alpha) \sum_{s=0}^{\infty} H_s[\beta] \frac{v^s}{s!} , \right. \end{split}$$

where we have put

(3)
$$H_r[\alpha] = H_r(0 \mid \alpha)$$

the so-called Eulerian number. Comparison of coefficients evidently yields $H_m(x \mid \alpha) H_n(x \mid \beta) = H_{m+n}(x \mid \alpha\beta)$

Supported in part by National Science Foundation grant G16485.

37

[Jan.-Feb.

(4)
$$+ \frac{\alpha(1-\beta)}{1-\alpha\beta} \sum_{r=1}^{m} {m \choose r} H_r[\alpha] H_{m+n-r}(x \mid \alpha\beta) \\ + \frac{\beta(1-\alpha)}{1-\alpha\beta} \sum_{s=1}^{n} {n \choose s} H_s[\beta] H_{m+n-s}(x \mid \alpha\beta),$$

provided $\alpha \neq 1$, $\beta \neq 1$, $\alpha \beta \neq 1$.

In the next place we have

$$\sum_{m,n=0}^{\infty} B_m(x) H_n(x \mid \alpha) \frac{u^m v^n}{m! n!} = \frac{u e^{xu}}{e^u - 1} \frac{(1 - \alpha) e^{xv}}{e^v - \alpha}$$

$$= u \frac{(1 - \alpha) e^{x(u+v)}}{e^{u+v} - \alpha} \frac{e^{u+v} - \alpha}{(e^u - 1)(e^v - \alpha)}$$

$$= \frac{(1 - \alpha) e^{x(u+v)}}{e^{u+v} - \alpha} \left\{ u + \frac{u}{e^u - 1} + \frac{\alpha u}{e^v - \alpha} \right\}$$

$$= \sum_{m,n=0}^{\infty} H_{m+n}(x \mid \alpha) \frac{u^m v^n}{m! m!} \left\{ u + \sum_{r=0}^{\infty} B_r \frac{u^r}{r!} + \frac{u}{1 - \alpha} \sum_{s=0}^{\infty} H_s[\alpha] \frac{v^s}{s!} \right\}.$$

It follows that

(5)
$$B_{m}(x)H_{n}(x \mid \alpha) = mH_{m+n-1}(x \mid \alpha) + \sum_{r=0}^{m} \binom{m}{r} B_{r}H_{m+n-r}(x \mid \alpha) + \frac{m\alpha}{1-\alpha} \sum_{s=0}^{n} \binom{n}{s} H_{s}[\alpha]H_{m+n-s-1}(x \mid \alpha),$$

provided $\alpha \neq 1$.

If $\alpha \neq 1$ but $\alpha \beta = 1$ we take

$$(u+v) \sum_{m,n=0}^{\infty} H_m(x \mid \alpha) H_n(x \mid \alpha^{-1}) \frac{u^m v^n}{m! n!} = (u+v) \frac{(1-\alpha)e^{xu}}{e^u - \alpha} \frac{(1-\alpha^{-1})e^{xv}}{e^v - \alpha^{-1}} \\ = (1-\alpha)(1-\alpha^{-1}) \frac{(u+v)e^{x(u+v)}}{e^{u+v} - 1} \left\{ 1 + \frac{\alpha}{e^u - \alpha} + \frac{\alpha^{-1}}{e^v - \alpha^{-1}} \right\}.$$

This implies

$$\begin{split} mH_{m-1}(x \mid \alpha)H_n(x \mid \alpha^{-1}) &+ nH_m(x \mid \alpha)H_{n-1}(x \mid \alpha^{-1}) \\ &= (1-\alpha)(1-\alpha^{-1})B_{m+n}(x) - (1-\alpha)\sum_{r=0}^m \binom{m}{r}H_r[\alpha]B_{m+n-r}(x) \\ &- (1-\alpha^{-1})\sum_{s=0}^n \binom{n}{s}H_s[\alpha^{-1}]B_{m+n-s}(x) \\ &= - (1-\alpha)\sum_{r=1}^m \binom{m}{r}H_r[\alpha]B_{m+n-r}(x) \\ &- (1-\alpha^{-1})\sum_{s=1}^n \binom{n}{s}H_s[\alpha^{-1}]B_{m+n-s}(x). \end{split}$$

This content downloaded from 158.110.11.25 on Fri, 23 Jan 2015 08:40:38 AM All use subject to JSTOR Terms and Conditions Since

$$\frac{\partial}{\partial x} H_n(x \mid \alpha) = n H_{n-1}(x \mid \alpha),$$

it is clear from (6) that

(7)
$$H_{m}(x \mid \alpha) H_{n}(x \mid \alpha^{-1}) = -(1-\alpha) \sum_{r=0}^{m-1} {m \choose r+1} H_{r+1}[\alpha] \frac{B_{m+n-r}(x)}{m+n-r} -(1-\alpha^{-1}) \sum_{s=0}^{n-1} {n \choose s+1} H_{s+1}[\alpha^{-1}] \frac{B_{m+n-s}(x)}{m+n-s} + C_{m,n},$$

where $C_{m,n}$ is independent of x. To determine $C_{m,n}$ we notice first that (6) and (7) imply

$$mC_{m-1,n} + nC_{m,n-1} = 0,$$

so that

$$C_{m,n} = -\frac{n}{m+1} C_{m+1,n-1}.$$

Repeated application of this recursion leads to

(8)
$$C_{m,n} = (-1)^n \frac{m!n!}{(m+n)!} C_{m+n,0}.$$

Now if we put n=0, x=0 in (7) we get

$$H_m[\alpha] = -(1-\alpha) \sum_{r=0}^{m-1} {m \choose r+1} H_{r+1}[\alpha] \frac{B_{m-r}}{m-r} + C_{m,0}$$
$$= -\frac{1-\alpha}{m+1} \sum_{r=1}^{m} {m+1 \choose r} H_r[\alpha] B_{m-r+1} + C_{m,0}.$$

Similarly (5) implies

$$B_{m+1} = (m+1)H_m[\alpha] + \sum_{r=0}^{m+1} \binom{m+1}{r} H_r[\alpha]B_{m-r+1} + \frac{(m+1)\alpha}{1-\alpha} H_m[\alpha],$$

so that

$$(m+1)C_{m,0} = -(1-\alpha)H_{m+1}[\alpha].$$

Therefore by (8)

(9)
$$C_{m,n} = (-1)^{n+1} \frac{m!n!}{(m+n+1)!} (1-\alpha) H_{m+n+1}[\alpha].$$

(Since

$$H_n[\alpha^{-1}] = (-1)^n H_n[\alpha],$$

1963]

This content downloaded from 158.110.11.25 on Fri, 23 Jan 2015 08:40:38 AM All use subject to JSTOR Terms and Conditions

MATHEMATICS MAGAZINE

the right member of (9) remains unchanged when we interchange m and n and replace α by α^{-1}).

Combining (7) and (9) we get

(10)

$$H_{m}(x \mid \alpha) H_{n}(x \mid \alpha^{-1}) = -(1 - \alpha) \sum_{r=1}^{m} {m \choose r} H_{r}[\alpha] \frac{B_{m+n-r+1}(x)}{m+n-r+1}$$

$$-(1 - \alpha^{-1}) \sum_{s=1}^{n} {n \choose s} H_{s}[\alpha^{-1}] \frac{B_{m+n-s+1}(x)}{m+n-s+1}$$

$$+(-1)^{n+1} \frac{m!n!}{(m+n+1)!} (1 - \alpha) H_{m+n+1}[\alpha],$$

where of course $\alpha \neq 1$.

In particular if we take $\alpha = -1$, (5) and (10) reduce to

(11)

$$B_{m}(x)E_{n}(x) = E_{m+n}(x) + \sum_{r=2}^{m} \binom{m}{r} B_{r}E_{m+n-r}(x)$$

$$-\frac{m}{2} \sum_{s=1}^{n} \binom{n}{s} 2^{-s}C_{s}E_{m+n-s-1}(x),$$

$$E_{m}(x)E_{n}(x) = -2 \sum_{r=1}^{m} \binom{m}{r} 2^{-r}C_{r} \frac{B_{m+n-r+1}(x)}{m+n-r+1}$$

$$-2 \sum_{s=1}^{n} \binom{n}{s} 2^{-s}C_{s} \frac{B_{m+n-s+1}(x)}{m+n-s+1}$$

$$+ (-1)^{n+1}2^{-m-n} \frac{m!n!}{(m+n+1)!} C_{m+n+1},$$

where [4, p. 28]

$$C_n = 2^n E_n(0) = (2 - 2^{-n}) \frac{B_{n+1}}{n+1}$$

The formulas (11) and (12) may be compared with [3, p. 77, formulas (12), (16)].

We note also that since

$$\int_0^1 B_m(x) dx = \frac{B_{m+1}(1) - B_{m+1}(0)}{m+1} = 0 \qquad (m \ge 1),$$

(10) yields

(13)
$$\int_{0}^{1} H_{m}(x \mid \alpha) H_{n}(x \mid \alpha^{-1}) dx = (-1)^{n+1} \frac{m! n!}{(m+n+1)!} (1-\alpha) H_{m+n+1}[\alpha]$$
$$(m \ge 1, n \ge 1).$$

Finally we remark that (4), (5) and (10) imply the following special formulas: 1963]

(14)
$$H_m(x \mid \alpha) = H_m(x \mid \beta) + \frac{\alpha - \beta}{1 - \beta} \sum_{r=1}^m \binom{m}{r} H_r[\alpha] H_{m-r}(x \mid \beta) \qquad (\alpha \neq 1, \beta \neq 1),$$

(15)
$$B_m(x) = mH_{m-1}(x \mid \alpha) + \sum_{r=0}^m \binom{m}{r} B_r H_{m-r}(x \mid \alpha) \qquad (\alpha \neq 1),$$

(16)
$$H_m(x \mid \alpha) = -\frac{1-\alpha}{m+1} \sum_{r=1}^{m+1} {m+1 \choose r} H_r[\alpha] B_{m-r+1}(x) \qquad (\alpha \neq 1).$$

It is not difficult to prove these formulas directly. For example (14) follows easily from the identity

$$\frac{(1-\alpha)e^{xu}}{e^u-\alpha} = \frac{1}{1-\beta} \left\{ 1-\alpha + (\alpha-\beta)\frac{1-\alpha}{e^u-\alpha} \right\} \frac{(1-\beta)e^{xu}}{e^x-\beta}$$

References

1. L. Carlitz, Eulerian numbers and polynomials, MATHEMATICS MAGAZINE, 32 (1959) 247-260.

2. L. Carlitz, Note on the integral of the product of several Bernoulli polynomials, Journal of the London Mathematical Society, **34** (1959) 361–363.

3. N. Nielsen, Traité élémentaire des nombres de Bernoulli, Paris, 1923.

4. N. E. Norlund, Vorlesungen über Differenzenrechnung, Berlin, 1924.

NUMBER THEORY

A pump's a composite of handle and spout That has to be primed, or nothing comes out. A gun's a composite of barrel and butt That has to be primed, or nothing will sput. In the arts, composition is carefully timed And one doesn't begin till the surface is primed. You will find composition is easy to do When you start with a primer and carry it thru.

MARLOW SHOLANDER