



# Fundamentos lógicos de bases de datos

(Logical foundations of databases)

Diego Figueira

Gabriele Puppis

CNRS

LaBRI



# Recap

- **Relational model** (tables)
- **Relational Algebra** (union, product, difference, selection, projection)
- **SQL** (SELECT ... FROM ... WHERE ...)
- **First-order logic** (syntax, semantics, active domain)
- **Expressiveness** ( $\text{FO}^{\text{act}} = \text{RA} = \text{basic SQL}$ )
- **Undecidable problems** (Halting  $\leq$  Domino  $\leq$  Satisfiability  $\leq$  Equivalence)
- **Data complexity / Combined complexity**
- **Complexity of evaluation** (LOGSPACE / PSPACE complexity)

# Definability in FO

**Goal:** check which properties / queries are *expressible in FO*

# Definability in FO

**Goal:** check which properties / queries are *expressible in FO*

Example.  $Q(G) = \{ (u, v) \mid G \text{ contains a path from } u \text{ to } v \}$

Is  $Q$  expressible as a first-order formula?

# Definability in FO

Definition. **Quantifier rank** of  $\phi$  = max number of nested quantifiers in  $\phi$ .

# Definability in FO

Definition. **Quantifier rank** of  $\phi$  = max number of nested quantifiers in  $\phi$ .

Example.  $\phi = \forall x \forall y \left( \neg E(x,y) \vee \exists z \left( (E(x,z) \vee E(z,x)) \wedge (E(y,z) \vee E(z,y)) \right) \right)$   
has quantifier rank 3.

# Definability in FO

Definition. **Quantifier rank** of  $\phi$  = max number of nested quantifiers in  $\phi$ .

Example.  $\phi = \forall x \forall y \left( \neg E(x,y) \vee \exists z \left( (E(x,z) \vee E(z,x)) \wedge (E(y,z) \vee E(z,y)) \right) \right)$   
has quantifier rank 3.

Quantifier rank  $\neq$  quantity of quantifiers

# Definability in FO

Definition. **Quantifier rank** of  $\phi$  = max number of nested quantifiers in  $\phi$ .

Example.  $\phi = \forall x \forall y \left( \neg E(x,y) \vee \exists z \left( (E(x,z) \vee E(z,x)) \wedge (E(y,z) \vee E(z,y)) \right) \right)$   
has quantifier rank 3.

Quantifier rank  $\neq$  quantity of quantifiers

*Eg*, in  $d_0(x, y) = E(x, y)$ , and

$$d_k(x, y) = \exists z ( d_{k-1}(x, z) \wedge d_{k-1}(z, y) )$$

$qr(d_k) = k$  but # quantifiers of  $d_k$  is  $2^k$

# Definability in FO

Definition. **Quantifier rank** of  $\phi$  = max number of nested quantifiers in  $\phi$ .


Example.  $\phi = \forall x \forall y \left( \neg E(x,y) \vee \exists z \left( (E(x,z) \vee E(z,x)) \wedge (E(y,z) \vee E(z,y)) \right) \right)$   
has quantifier rank 3.

Quantifier rank  $\neq$  quantity of quantifiers

*Eg*, in  $d_0(x,y) = E(x,y)$ , and

$$d_k(x,y) = \exists z ( d_{k-1}(x,z) \wedge d_{k-1}(z,y) )$$

$qr(d_k) = k$  but # quantifiers of  $d_k$  is  $2^k$



What does it  
define?

# Definability in FO

Definition. **Quantifier rank** of  $\phi$  = max number of nested quantifiers in  $\phi$ .

Example.  $\phi = \forall x \forall y \left( \neg E(x,y) \vee \exists z \left( (E(x,z) \vee E(z,x)) \wedge (E(y,z) \vee E(z,y)) \right) \right)$   
has quantifier rank 3.

Quantifier rank  $\neq$  quantity of quantifiers

*Eg*, in  $d_0(x,y) = E(x,y)$ , and

$$d_k(x,y) = \exists z ( d_{k-1}(x,z) \wedge d_{k-1}(z,y) )$$

$qr(d_k) = k$  but # quantifiers of  $d_k$  is  $2^k$

What does it  
define?

Quantifier rank is a **measure of complexity** of a formula

# Definability in FO

Definition. **Quantifier rank** of  $\phi$  = max number of nested quantifiers in  $\phi$ .


Example.  $\phi = \forall x \forall y \left( \neg E(x,y) \vee \exists z \left( (E(x,z) \vee E(z,x)) \wedge (E(y,z) \vee E(z,y)) \right) \right)$   
has quantifier rank 3.

Quantifier rank  $\neq$  quantity of quantifiers

*Eg*, in  $d_0(x,y) = E(x,y)$ , and

$$d_k(x,y) = \exists z ( d_{k-1}(x,z) \wedge d_{k-1}(z,y) )$$

$qr(d_k) = k$  but # quantifiers of  $d_k$  is  $2^k$



What does it  
define?

Quantifier rank is a **measure of complexity** of a formula

**Sub-goal:** Given a property  $P$  and a number  $n$ ,  
tell whether  $P$  is expressible by a sentence of quantifier rank at most  $n$ .

# Definability in FO

Definition. Two structures  $S_1$  and  $S_2$  are  **$n$ -equivalent**  
iff

they satisfy the same FO sentences of quantifier rank  $\leq n$   
( i.e.  $S_1 \models \phi$  iff  $S_2 \models \phi$  for all  $\phi \in \text{FO}$  with  $\text{qr}(\phi) \leq n$ )

[Tarski '30]

# Definability in FO

Definition. Two structures  $S_1$  and  $S_2$  are  **$n$ -equivalent**

[Tarski '30]

iff

they satisfy the same FO sentences of quantifier rank  $\leq n$

( i.e.  $S_1 \models \phi$  iff  $S_2 \models \phi$  for all  $\phi \in \text{FO}$  with  $\text{qr}(\phi) \leq n$ )

Consider a property (i.e. a set of structures)  $P$ .

Suppose that there are  $S_1 \in P, S_2 \notin P$  s.t.

$S_1$  and  $S_2$  are  $n$ -equivalent.

Then  $P$  is *not expressible* by any sentence of quantifier rank  $n$ .

# Definability in FO

Definition. Two structures  $S_1$  and  $S_2$  are  **$n$ -equivalent**  
iff  
they satisfy the same FO sentences of quantifier rank  $\leq n$   
( i.e.  $S_1 \models \phi$  iff  $S_2 \models \phi$  for all  $\phi \in \text{FO}$  with  $\text{qr}(\phi) \leq n$ )

[Tarski '30]

Consider a property (i.e. a set of structures)  $P$ .

Suppose that there are  $S_1 \in P, S_2 \notin P$  s.t.

$S_1$  and  $S_2$  are  $n$ -equivalent.

Then  $P$  is *not expressible* by any sentence of quantifier rank  $n$ .

Note: if the above happens  
 $\forall n$ , then  $P$  is not  
expressible by *any* FO  
sentence.

# Definability in FO

Definition. Two structures  $S_1$  and  $S_2$  are *n-equivalent* iff they satisfy the same FO sentences of quantifier rank  $\leq n$  ( i.e.  $S_1 \models \phi$  iff  $S_2 \models \phi$  for all  $\phi \in \text{FO}$  with  $\text{qr}(\phi) \leq n$ )

[Tarski '30]

Consider a property (i.e. a set of structures)  $P$ .

Suppose that there are  $S_1 \in P, S_2 \notin P$  s.t.

$S_1$  and  $S_2$  are *n-equivalent*.

Then  $P$  is *not expressible* by any sentence of quantifier rank  $n$ .

Note: if the above happens  $\forall n$ , then  $P$  is not expressible by *any* FO sentence.

Example.  $P = \{ \text{finite structures} \}$  seems to be not FO-definable.

One could then aim at proving that

for all  $n$  there are  $S_1 \in P$  and  $S_2 \notin P$  s.t.  $S_1, S_2$  *n-equivalent*...

# Expressive power via games

# Expressive power via games

Characterization of the expressive power of FO in terms of **Games**

# Expressive power via games

Characterization of the expressive power of FO in terms of **Games**

Idea: For every two structures  $(S, S')$  there is a game where

a player of the game has a **winning strategy**

iff

$S, S'$  are **indistinguishable**

# Ehrenfeucht-Fraïssé games

A game between two players

$S_1$  and  $S_2$  are  
 $n$ -equivalent!



Duplicator

No they're  
**NOT!!!!**



Spoiler

Board:  $(S_1, S_2)$

One player plays in one structure, the other player answers in the other structure.

If after  $n$  rounds Duplicator doesn't lose:  $S_1, S_2$  are  $n$ -equivalent

# Ehrenfeucht-Fraïssé games

Definition. Partial isomorphism between  $S_1$  and  $S_2$  = injective partial map

$f$ : nodes of  $S_1 \rightarrow$  nodes of  $S_2$

so that

$E(x,y)$  iff  $E(f(x),f(y))$

# Ehrenfeucht-Fraïssé games

Definition. Partial isomorphism between  $S_1$  and  $S_2$  = **injective** partial map

$f$ : nodes of  $S_1 \rightarrow$  nodes of  $S_2$

so that

$E(x,y)$  iff  $E(f(x),f(y))$

*Spoiler*

and

*Duplicator*

play for  $n$  rounds on the board  $S_1, S_2$

# Ehrenfeucht-Fraïssé games

Definition. Partial isomorphism between  $S_1$  and  $S_2$  = **injective** partial map

$$f: \text{nodes of } S_1 \rightarrow \text{nodes of } S_2$$

so that

$$E(x,y) \text{ iff } E(f(x),f(y))$$

**Spoiler** and **Duplicator** play for  $n$  rounds on the board  $S_1, S_2$

At each round  $i$ :

1. **Spoiler** chooses a node  $x_i$  from  $S_1$   
and **Duplicator** answers with a node  $y_i$  from  $S_2$ ,

# Ehrenfeucht-Fraïssé games

Definition. Partial isomorphism between  $S_1$  and  $S_2$  = **injective** partial map

$$f: \text{nodes of } S_1 \rightarrow \text{nodes of } S_2$$

so that

$$E(x,y) \text{ iff } E(f(x),f(y))$$

**Spoiler** and **Duplicator** play for  $n$  rounds on the board  $S_1, S_2$

At each round  $i$ :

1. **Spoiler** chooses a node  $x_i$  from  $S_1$   
and **Duplicator** answers with a node  $y_i$  from  $S_2$ ,

or

2. **Spoiler** chooses a node  $y_i$  from  $S_2$   
and **Duplicator** answers with a node  $x_i$  from  $S_1$ ,

# Ehrenfeucht-Fraïssé games

Definition. Partial isomorphism between  $S_1$  and  $S_2$  = injective partial map

$$f: \text{nodes of } S_1 \rightarrow \text{nodes of } S_2$$

so that

$$E(x,y) \text{ iff } E(f(x),f(y))$$

*Spoiler* and *Duplicator* play for  $n$  rounds on the board  $S_1, S_2$

At each round  $i$ :

1. **Spoiler** chooses a node  $x_i$  from  $S_1$   
and **Duplicator** answers with a node  $y_i$  from  $S_2$ ,


or

2. **Spoiler** chooses a node  $y_i$  from  $S_2$   
and **Duplicator** answers with a node  $x_i$  from  $S_1$ ,


or **Spoiler** wins if  $\{x_i \mapsto y_i \mid 1 \leq i \leq n\}$  is not a partial isomorphism between  $S_1$  and  $S_2$ .


# Ehrenfeucht-Fraïssé games


$S_1$



$S_2$




 = Spoiler


 = Duplicator

# Ehrenfeucht-Fraïssé games


$S_1$



$S_2$




 = Spoiler


 = Duplicator


# Ehrenfeucht-Fraïssé games


$S_1$



$S_2$




 = Spoiler


 = Duplicator

# Ehrenfeucht-Fraïssé games


$S_1$



$S_2$




 = Spoiler


 = Duplicator


# Ehrenfeucht-Fraïssé games


$S_1$



$S_2$




 = Spoiler


 = Duplicator

# Ehrenfeucht-Fraïssé games


$S_1$



$S_2$




 = Spoiler

 = Duplicator


# Ehrenfeucht-Fraïssé games


Question: Can Spoiler win in 3 rounds ?


$S_1$



$S_2$




 = Spoiler

 = Duplicator


# Ehrenfeucht-Fraïssé games


Question: Can Spoiler win in 3 rounds ?

$S_1$



$S_2$




 = Spoiler

 = Duplicator


# Ehrenfeucht-Fraïssé games


Question: Can Spoiler win in 3 rounds ?

$S_1$



$S_2$




 = Spoiler

 = Duplicator


# Ehrenfeucht-Fraïssé games


Question: Can Spoiler win in 3 rounds ?

$S_1$



$S_2$




 = Spoiler

 = Duplicator


# Ehrenfeucht-Fraïssé games


Question: Can Spoiler win in 3 rounds ?

$S_1$



$S_2$




 = Spoiler

 = Duplicator


# Ehrenfeucht-Fraïssé games


Question: Can Spoiler win in 3 rounds ?

$S_1$



$S_2$



 = Spoiler

 = Duplicator

# Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* structures, Spoilers wins eventually...



**Why?**

# Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* structures, Spoilers wins eventually...

Why?

...and he often wins very quickly:





# Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* structures, Spoilers wins eventually...

Why?

...and he often wins very quickly:





# Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* structures, Spoilers wins eventually...

Why?

...and he often wins very quickly:





# Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* structures, Spoilers wins eventually...

Why?

...and he often wins very quickly:





# Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* structures, Spoilers wins eventually...

Why?

...and he often wins very quickly:





# Ehrenfeucht-Fraïssé games

On non-isomorphic *finite* structures, Spoilers wins eventually...

Why?

...and he often wins very quickly:



But there are non-isomorphic *infinite* structures where Duplicator can survive for *arbitrarily many rounds* (not necessarily forever!)



Any idea?

# Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* structures, Spoilers wins eventually...

Why?

...and he often wins very quickly:



But there are non-isomorphic *infinite* structures where Duplicator can survive for *arbitrarily many rounds* (not necessarily forever!)





# Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* structures, Spoilers wins eventually...

Why?

...and he often wins very quickly:



But there are non-isomorphic *infinite* structures where Duplicator can survive for *arbitrarily many rounds* (not necessarily forever!)





Given  $n$ ,  
at each round  $i = 1, \dots, n$ ,  
pairs of marked nodes in  $S_1$  and  $S_2$   
must be either at *equal distance*  
or at *distance  $\geq 2^{n-i}$*

# Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* structures, Spoilers wins eventually...

Why?

...and he often wins very quickly:



But there are non-isomorphic *infinite* structures where Duplicator can survive for *arbitrarily many rounds* (not necessarily forever!)





Given  $n$ ,  
at each round  $i = 1, \dots, n$ ,  
pairs of marked nodes in  $S_1$  and  $S_2$   
must be either at *equal distance*  
or at *distance  $\geq 2^{n-i}$*

# Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* structures, Spoilers wins eventually...

Why?

...and he often wins very quickly:



But there are non-isomorphic *infinite* structures where Duplicator can survive for *arbitrarily many rounds* (not necessarily forever!)





Given  $n$ ,  
 at each round  $i = 1, \dots, n$ ,  
 pairs of marked nodes in  $S_1$  and  $S_2$   
 must be either at *equal distance*  
 or at *distance  $\geq 2^{n-i}$*

# Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* structures, Spoilers wins eventually...

Why?

...and he often wins very quickly:



But there are non-isomorphic *infinite* structures where Duplicator can survive for *arbitrarily many rounds* (not necessarily forever!)





Given  $n$ ,  
at each round  $i = 1, \dots, n$ ,  
pairs of marked nodes in  $S_1$  and  $S_2$   
must be either at *equal distance*  
or at *distance  $\geq 2^{n-i}$*

# Ehrenfeucht-Fraïssé games



On non-isomorphic *finite* structures, Spoilers wins eventually...

Why?

...and he often wins very quickly:



But there are non-isomorphic *infinite* structures where Duplicator can survive for *arbitrarily many rounds* (not necessarily forever!)



Given  $n$ ,  
 at each round  $i = 1, \dots, n$ ,  
 pairs of marked nodes in  $S_1$  and  $S_2$   
 must be either at *equal distance*  
 or at *distance  $\geq 2^{n-i}$*

# Ehrenfeucht-Fraïssé games

**Theorem.**

$S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]

iff

Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

# Ehrenfeucht-Fraïssé games

**Theorem.**

$S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]

iff

Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

**Exercise:** prove that the following **corollary** of the theorem.

If  $\mathbf{P}$  is a property and

for all  $n$  there are  $S_1 \in \mathbf{P}$ ,  $S_2 \notin \mathbf{P}$  s.t. Duplicator can survive  $n$  rounds on  $S_1$  and  $S_2$

Then  $\mathbf{P}$  is *not definable in FO*.

# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent [Fraïssé '50, Ehrenfeucht '60]  
iff  
Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

**Exercise:** prove that the following **corollary** of the theorem.

If  $\mathbf{P}$  is a property and  
for all  $n$  there are  $S_1 \in \mathbf{P}, S_2 \notin \mathbf{P}$  s.t. Duplicator can survive  $n$  rounds on  $S_1$  and  $S_2$   $\rangle^*$

Then  $\mathbf{P}$  is *not definable in FO*.

**Proof:** Suppose that  $\mathbf{P}$  is defined by an FO sentence  $\phi$  and  $\ast$  holds.

From  $\ast$  and the above theorem:  $\forall n \exists S_1 \in \mathbf{P} \exists S_2 \notin \mathbf{P} . S_1, S_2$  are  $n$ -equivalent.

In particular, when  $n = \text{quantifier rank of } \phi$ ,  $S_1 \models \phi$  iff  $S_2 \models \phi$ .

A contradiction!

# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

Consider  $\phi$  with quantifier rank  $n$ .

Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .

We need to prove that  $S_2 \models \phi$ .

# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

Consider  $\phi$  with quantifier rank  $n$ .

Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .

We need to prove that  $S_2 \models \phi$ .



A new game to evaluate formulas....

# The semantics game

Assume w.l.o.g. that  $\phi$  is in **negation normal form**.

# The semantics game

Assume w.l.o.g. that  $\phi$  is in **negation normal form**.

push negations inside:

$$\neg \forall \phi \rightsquigarrow \exists \neg \phi$$

$$\neg \exists \phi \rightsquigarrow \forall \neg \phi$$

$$\neg(\phi \wedge \psi) \rightsquigarrow \neg \phi \vee \neg \psi$$

...

# The semantics game

Assume w.l.o.g. that  $\phi$  is in **negation normal form**.

push negations inside:

$$\neg \forall \phi \rightsquigarrow \exists \neg \phi$$

$$\neg \exists \phi \rightsquigarrow \forall \neg \phi$$

$$\neg(\phi \wedge \psi) \rightsquigarrow \neg \phi \vee \neg \psi$$

...

Whether  $S \models \phi$  can be decided by a *new game* between two players, **True** and **False**:

- $\phi = E(x,y)$   $\rightarrow$  **True** wins if nodes marked  $x$  and  $y$  are connected by an edge, otherwise he loses
- $\phi = \exists x \phi'(x)$   $\rightarrow$  **True** moves by marking a node  $x$  in  $S$ , the game continues with  $\phi'$
- $\phi = \forall y \phi'(y)$   $\rightarrow$  **False** moves by marking a node  $y$  in  $S$ , the game continues with  $\phi'$
- $\phi = \phi_1 \vee \phi_2$   $\rightarrow$  **True** moves by choosing  $\phi_1$  or  $\phi_2$ , the game continues with what he chose
- $\phi = \phi_1 \wedge \phi_2$   $\rightarrow$  **False** moves by choosing  $\phi_1$  or  $\phi_2$ , the game continues with what he chose
- ...

# The semantics game

Assume w.l.o.g. that  $\phi$  is in **negation normal form**.

push negations inside:

$$\neg \forall \phi \rightsquigarrow \exists \neg \phi$$

$$\neg \exists \phi \rightsquigarrow \forall \neg \phi$$

$$\neg(\phi \wedge \psi) \rightsquigarrow \neg \phi \vee \neg \psi$$

...

Whether  $S \models \phi$  can be decided by a *new game* between two players, **True** and **False**:

- $\phi = E(x,y)$   $\rightarrow$  **True** wins if nodes marked  $x$  and  $y$  are connected by an edge, otherwise he loses
- $\phi = \exists x \phi'(x)$   $\rightarrow$  **True** moves by marking a node  $x$  in  $S$ , the game continues with  $\phi'$
- $\phi = \forall y \phi'(y)$   $\rightarrow$  **False** moves by marking a node  $y$  in  $S$ , the game continues with  $\phi'$
- $\phi = \phi_1 \vee \phi_2$   $\rightarrow$  **True** moves by choosing  $\phi_1$  or  $\phi_2$ , the game continues with what he chose
- $\phi = \phi_1 \wedge \phi_2$   $\rightarrow$  **False** moves by choosing  $\phi_1$  or  $\phi_2$ , the game continues with what he chose
- ...

**Lemma.**  $S \models \phi$  iff **True** wins the semantics game.

# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

Consider  $\phi$  with quantifier rank  $n$ .

Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .

We need to prove that  $S_2 \models \phi$ .

# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)




True wins the game on  $S_1$

Consider  $\phi$  with quantifier rank  $n$ .

Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .

We need to prove that  $S_2 \models \phi$ .



True wins the game on  $S_2$

# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

**True** wins the game on  $S_1$

Consider  $\phi$  with quantifier rank  $n$ .

Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .

We need to prove that  $S_2 \models \phi$ .

**True** wins the game on  $S_2$



Turn winning strategy for **True** in  $S_1$  into winning strategy for **True** in  $S_2$ ....

# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .



Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

Consider  $\phi$  with quantifier rank  $n$ .


Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .

We need to prove that  $S_2 \models \phi$ .

$S_1$



$S_2$



# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .


Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

Consider  $\phi$  with quantifier rank  $n$ .

Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .


We need to prove that  $S_2 \models \phi$ .

$S_1$



F

$S_2$



# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .


Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

Consider  $\phi$  with quantifier rank  $n$ .


Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .

We need to prove that  $S_2 \models \phi$ .

$S_1$



$S_2$



# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]


iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

Consider  $\phi$  with quantifier rank  $n$ .

Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .

We need to prove that  $S_2 \models \phi$ .



# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]


iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

Consider  $\phi$  with quantifier rank  $n$ .

Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .

We need to prove that  $S_2 \models \phi$ .



# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]


iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

Consider  $\phi$  with quantifier rank  $n$ .

Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .

We need to prove that  $S_2 \models \phi$ .



# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]


iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

Consider  $\phi$  with quantifier rank  $n$ .

Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .

We need to prove that  $S_2 \models \phi$ .



# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]


iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

Consider  $\phi$  with quantifier rank  $n$ .

Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .

We need to prove that  $S_2 \models \phi$ .



# Ehrenfeucht-Fraïssé games

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]


iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

Proof ideas for the if-direction (from Duplicator's winning strategy to  $n$ -equivalence)

Consider  $\phi$  with quantifier rank  $n$ .

Suppose  $S_1 \models \phi$  and Duplicator survives  $n$  rounds on  $S_1, S_2$ .

We need to prove that  $S_2 \models \phi$ .



# Definability in FO

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent

[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

**Corollary.** A property  $P$  is *not definable in FO*

iff  $\forall n \exists S_1 \in P \exists S_2 \notin P$  Duplicator can survive  $n$  rounds on  $S_1$  and  $S_2$ .

# Definability in FO

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent



[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

**Corollary.** A property  $P$  is *not definable in FO*

iff  $\forall n \exists S_1 \in P \exists S_2 \notin P$  Duplicator can survive  $n$  rounds on  $S_1$  and  $S_2$ .

Example:  $P = \{ \text{connected graphs} \}$ . Given  $n$ , take  $S_1 \in P$  and  $S_2 = S_1 \uplus S_1 \notin P$



# Definability in FO

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent



[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

**Corollary.** A property  $P$  is *not definable in FO*

iff  $\forall n \exists S_1 \in P \exists S_2 \notin P$  Duplicator can survive  $n$  rounds on  $S_1$  and  $S_2$ .

Example:  $P = \{ \text{connected graphs} \}$ . Given  $n$ , take  $S_1 \in P$  and  $S_2 = S_1 \uplus S_1 \notin P$



# Definability in FO

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent



[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

**Corollary.** A property  $P$  is *not definable in FO*

iff  $\forall n \exists S_1 \in P \exists S_2 \notin P$  Duplicator can survive  $n$  rounds on  $S_1$  and  $S_2$ .

Example:  $P = \{ \text{connected graphs} \}$ . Given  $n$ , take  $S_1 \in P$  and  $S_2 = S_1 \uplus S_1 \notin P$



# Definability in FO

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent



[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

**Corollary.** A property  $P$  is *not definable in FO*

iff  $\forall n \exists S_1 \in P \exists S_2 \notin P$  Duplicator can survive  $n$  rounds on  $S_1$  and  $S_2$ .

Example:  $P = \{ \text{connected graphs} \}$ . Given  $n$ , take  $S_1 \in P$  and  $S_2 = S_1 \uplus S_1 \notin P$



# Definability in FO

**Theorem.**  $S_1$  and  $S_2$  are  $n$ -equivalent



[Fraïssé '50, Ehrenfeucht '60]

iff Duplicator has a strategy to survive  $n$  rounds in the EF game on  $S_1$  and  $S_2$ .

**Corollary.** A property  $P$  is *not definable in FO*

iff  $\forall n \exists S_1 \in P \exists S_2 \notin P$  Duplicator can survive  $n$  rounds on  $S_1$  and  $S_2$ .

Example:  $P = \{ \text{connected graphs} \}$ . Given  $n$ , take  $S_1 \in P$  and  $S_2 = S_1 \uplus S_1 \notin P$



# Ehrenfeucht-Fraïssé games

Several properties can be proved to be *not FO-definable*:

- connectivity ( previous slide )

# Ehrenfeucht-Fraïssé games

Several properties can be proved to be *not FO-definable*:

- connectivity ( previous slide )

- even / odd size      **Your turn now!**      ...given  $n$ , take  $S_1 =$  large even structure  
 $S_2 =$  large odd structure...



# Ehrenfeucht-Fraïssé games

Several properties can be proved to be *not FO-definable*:

- connectivity (previous slide)

- even / odd size **Your turn now!** ...given  $n$ , take  $S_1 =$  large even structure  
 $S_2 =$  large odd structure...

- 2-colorability Given  $n$ , take  $S_1 =$  large even cycle  $S_2 =$  large odd cycle



# Ehrenfeucht-Fraïssé games

Several properties can be proved to be *not FO-definable*:

- connectivity (previous slide)



- even / odd size **Your turn now!** ...given  $n$ , take  $S_1 =$  large even structure  
 $S_2 =$  large odd structure...

- 2-colorability Given  $n$ , take  $S_1 =$  large even cycle  $S_2 =$  large odd cycle

- finiteness

- acyclicity

...



# Some Bibliography

- Libkin, “Elements of Finite Model Theory”, Springer, 2004.
- Otto, “Finite Model Theory”, Springer, 2005  
(freely available at [www.mathematik.tu-darmstadt.de/~otto/LEHRE/FMT0809.ps](http://www.mathematik.tu-darmstadt.de/~otto/LEHRE/FMT0809.ps))
- Väänänen, “A Short course on Finite Model Theory”, 1994.  
(available at [www.math.helsinki.fi/logic/people/jouko.vaanananen/shortcourse.pdf](http://www.math.helsinki.fi/logic/people/jouko.vaanananen/shortcourse.pdf))