

where all the right-hand side parameters are positive, and

$$(3.5) \quad t^2 = 2(xu + yv + zw + zu + zv), \quad t \text{ even.}$$

Similar extensions follow for the n -tids.

An alternate method of generating infinite numbers of Pythagorean n -tids from a given n -tid is discussed in [7].

REFERENCES

1. M. Feinberg. "Fibonacci-Tribonacci." *The Fibonacci Quarterly* 1, No. 3 (1963):71-74.
2. A. F. Horadam. "A Generalized Fibonacci Sequence." *American Math. Monthly* 68 (1961): 455-456.
3. A. F. Horadam. "Special Properties of the Sequence $W_n(a, b, p, q)$." *The Fibonacci Quarterly* 5 (1967):424-434.
4. A. G. Shannon and A. F. Horadam. "A Generalized Pythagorean Theorem." *The Fibonacci Quarterly* 9 (1971):307-312.
5. A. G. Shannon and A. F. Horadam. "Generalized Fibonacci Number Triples." *American Math. Monthly* 80 (1973):178-190.
6. M. G. Teigen and D. W. Hadwin. "On Generating Pythagorean Triples." *American Math. Monthly* 78 (1971):378-379.
7. H. V. Krishna. "On Generating Pythagorean Pentids." *Proceedings of the 42nd I.M.S. Conference*, 1976.

A TRIANGLE FOR THE BELL NUMBERS

JEFFREY SHALLIT

University of California, Berkeley, California

The Bell, or exponential, numbers B_n are defined by

$$(1) \quad B_n = \frac{1}{e} \sum_{k=0}^{\infty} \frac{k^n}{k!} = \frac{1}{e} \left(\frac{0^n}{0!} + \frac{1^n}{1!} + \frac{2^n}{2!} + \dots \right)$$

The first twelve Bell numbers are given in the following table:

TABLE 1. Bell Numbers

n	B_n
0	1
1	1
2	2
3	5
4	15
5	52
6	203
7	877
8	4140
9	21147
10	115975
11	678570

The Bell numbers also appear in the Maclaurin expansion of e^{e^x} :

$$(2) \quad e^{e^x} = e \sum_{k=0}^{\infty} \frac{B_k x^k}{k!} = e \left(1 + \frac{x}{1!} + \frac{2x^2}{2!} + \frac{5x^3}{3!} + \frac{15x^4}{4!} + \dots \right)$$

The Bell numbers can be generated recursively by an interesting method described in [2]. If we take the array described in this article and "flip" it about and then reorient it, the following triangle appears. This triangle is similar in form to Pascal's triangle. We shall call it the "Bell Triangle," and denote each element by $B'(n,r)$. This notation is similar to $C(n,r)$ for Pascal's triangle. There are three rules of formation for this triangle.

- (3) $B'(0,0) = 1$
 (4) $B'(n,0) = B'(n-1, n-1)$ ($n \geq 1$)
 (5) $B'(n,r) = B'(n,r-1) + B'(n-1, r-1)$ ($1 \leq r \leq n$)

Row										
0				1						
1			1	2						
2			2	3	5					
3			5	7	10	15				
4			15	20	27	37	52			
5			52	67	87	114	151	203		
6			203	255	322	409	523	674	877	
7			877	1080	1335	1657	2066	2589	3263	4140

$B'(3,2) + B'(4,2) = B'(4,3)$

The Bell numbers form the left and right sides of the triangle. In fact,

- (6) $B'(n,n) = B_{n+1}$
 (7) $B'(n,0) = B_n$

Equations (6) and (7) follow from the two equivalent identities for Bell numbers:

- (8) $B_n = \binom{n}{0}B_{n+1} - \binom{n}{1}B_n + \binom{n}{2}B_{n-1} - \dots \pm \binom{n}{n}B_1$
 (9) $B_n = nB_{n-1} - \binom{n-1}{2}B_{n-2} + \binom{n-1}{3}B_{n-3} - \dots \pm \binom{n-1}{n-1}B_1$

The Bell triangle has many interesting properties. Here we present several new identities:

(10)
$$\sum_{k=a}^b B'(n,k) = B'(n+1, b+1) - B'(n+1, a).$$

For $a = 0$ and $b = n$, this reduces to

(11)
$$\sum_{k=0}^n B'(n,k) = B'(n+1, n+1) - B'(n+1, 0) = B'(n+1, n) = B_{n+2} - B_{n+1},$$

(12)
$$\sum_{k=x}^n B'(k+a, k) = B'(n+a, n+1) - B'(x+a-1, x),$$

(13)
$$\sum_{k=x}^n (-1)^{n-k} \binom{n-x}{k-x} B'(k+a, k) = B'(n+a, x).$$

For $a = 0$ and $x = 0$, equation (13) reduces to (8).

(14)
$$\sum_{k=x}^n \binom{n-x}{k-x} B'(k, a) = B'(n, a+n-x).$$

For $a = 0$ and $x = 0$, the following identity results:

(15)
$$\sum_{k=0}^n \binom{n}{k} B'(k, 0) = B'(n, n).$$

This is equivalent to

(16)
$$\sum_{k=0}^n \binom{n}{k} B_k = B_{n+1}.$$

To my knowledge, identities (10)-(14) were heretofore unknown.

If we ignore the restricting inequality in (4), and substitute $n = 0$, we get $1 = B'(0,0) = B'(-1,-1)$. From this value, we may obtain values of $B'(n,-1)$ for $n \geq -1$ (see Table 2).

Note the following identity:

(17) $B'(n-1, -1) + B'(n, -1) = B'(n, 0) = B_n.$

TABLE 2. Values of $B'(n,-1)$

n	$B'(n,-1)$
-1	1
0	0
1	1
2	1
3	4
4	11
5	41
6	162
7	715

Apparently the Bell triangle cannot be extended further because $B(-1,0) = B_{-1}$ which is undefined, by equation (1). Epstein [3] drops the term $0^n/0!$ in equation (1) without explanation and therefore gets $B_0 = 1 - 1/e$, in contradiction with Williams [5], Bell [1], and Rota [4].

The Bell numbers have combinatoric significance in that B_n is the number of ways of factoring a product of n distinct primes. Whether the rest of the numbers in the Bell triangle have any such significance remains to be seen.

REFERENCES

1. E. T. Bell. "Exponential Numbers." *American Math. Monthly* 41 (1934):411-419.
2. Martin Cohn, et al. "On the Number of Partitionings of a Set of n Distinct Objects." *American Math. Monthly* 69 (1962):782-785.
3. Leo Epstein. "A Function Related to the Series e^{e^x} ." *J. Math. and Physics* 18 (1939):153-173.
4. Gian-Carlo Rota. "The Number of Partitions of a Set." *American Math. Monthly* 71 (1964):498-504.
5. G. T. Williams. "Numbers Generated by the Function $e^{e^{x-1}}$." *American Math. Monthly* 52 (1945):323-327.

THE EQUATIONS $z^2 - 3y^2 = -2$ AND $z^2 - 6x^2 = -5$

MANORANJITHAM VELUPPILLAI

Royal Holloway College, Egham, Surrey, England

The four numbers 2, 4, 12, 420 have the property that the product of any two increased by 1 is a perfect square. The object of this paper is to prove that no positive integer can replace 420.

Any integer N which can replace 420 while preserving this property must satisfy the equations

$$2N + 1 = x^2, \quad 4N + 1 = y^2, \quad 12N + 1 = z^2.$$

Eliminating N , we have

$$z^2 - 3y^2 = -2 \quad \text{and} \quad z^2 - 6x^2 = -5.$$

Now, the equation $z^2 - 3y^2 = -2$ can be written in the form

$$(1) \quad u^2 - 3v^2 = 1$$

where $u = z^2 + 1$, $v = zy$.

Substituting for z^2 in $z^2 - 6x^2 = -5$, we have

$$(2) \quad X^2 = 6u + 24$$

where $X = 6x$.

Hence, to solve the equations of the title, it is sufficient to solve (1) and (2) simultaneously.

Now, all the positive integral solutions of (1) are given by the formula: