

A q -analogue of Catalan Hankel determinants

By

Masao ISHIKAWA,*Hiroyuki TAGAWA,**and Jiang ZENG***

Abstract

In this article we shall survey the various methods of evaluating Hankel determinants and as an illustration we evaluate some Hankel determinants of a q -analogue of Catalan numbers. Here we consider $\frac{(aq;q)_n}{(abq^2;q)_n}$ as a q -analogue of Catalan numbers $C_n = \frac{1}{n+1} \binom{2n}{n}$, which is known as the moments of the little q -Jacobi polynomials. We also give several proofs of this q -analogue, in which we use lattice paths, the orthogonal polynomials, or the basic hypergeometric series. We also consider a q -analogue of Schröder Hankel determinants, and give a new proof of Moztkin Hankel determinants using an addition formula for ${}_2F_1$.

§ 1. Introduction

Given a sequence a_0, a_1, a_2, \dots , we set the Hankel matrix of the sequence to be

$$(1.1) \quad A_n^{(t)} = (a_{i+j+t})_{0 \leq i, j \leq n-1} = \begin{pmatrix} a_t & a_{t+1} & \dots & a_{t+n-1} \\ a_{t+1} & a_{t+2} & \dots & a_{t+n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{t+n-1} & a_{t+n} & \dots & a_{t+2n-2} \end{pmatrix}.$$

For $n = 0, 1, 2, \dots$, let

$$(1.2) \quad C_n = \frac{1}{n+1} \binom{2n}{n},$$

which are called the *Catalan numbers*. The generating function for the Catalan numbers is given by

$$\sum_{n \geq 0} C_n t^n = \frac{1 - \sqrt{1 - 4t}}{2t}.$$

Received May 8, 2008. Accepted July 6, 2008.

2000 Mathematics Subject Classification(s): Primary 05A30; Secondary 05A10, 05E35, 15E15, 33D15

Key Words: Catalan numbers, determinants, Dyck paths, orthogonal polynomials, continued fractions

*Faculty of Education, Tottori University, Koyama, Tottori, Japan.

**Faculty of Education, Wakayama University, Sakaedani, Wakayama, Japan.

***Institut Camille Jordan, Université Claude Bernard Lyon I, 43, boulevard du 11 novembre 1918, 69622 Villeurbanne Cedex, France.

If we put $a_n = C_n$ in (1.1), then the following identity is well-known and several proofs are known [5, 6, 14, 17, 19]:

$$(1.3) \quad \det A_n^{(t)} = \det (C_{i+j+t})_{0 \leq i, j \leq n-1} = \prod_{1 \leq i \leq j \leq t-1} \frac{i+j+2n}{i+j}.$$

If we put $B_n = \binom{2n+1}{n}$ and $D_n = \binom{2n}{n}$, then the following variations are also known [17]:

$$(1.4) \quad \det (B_{i+j+t})_{0 \leq i, j \leq n-1} = \prod_{1 \leq i \leq j \leq t-1} \frac{i+j-1+2n}{i+j-1},$$

$$(1.5) \quad \det (D_{i+j+t})_{0 \leq i, j \leq n-1} = 2^n \prod_{1 \leq i < j \leq t-1} \frac{i+j+2n}{i+j}.$$

As a generalization of (1.3), Krattenthaler [12] has obtained

$$(1.6) \quad \det (C_{k_i+j})_{0 \leq i, j \leq n-1} = \prod_{0 \leq i < j \leq n-1} (k_j - k_i) \prod_{i=0}^{n-1} \frac{(i+n)!(2k_i)!}{(2i)!k_i!(k_i+n)!}$$

for a positive integer n and non-negative integers k_0, k_1, \dots, k_{n-1} .

In this article we shall survey the various methods of evaluating Hankel determinants and as an illustration we give a q -analogue of the above results. We first recall some terminology in q -series (see Gasper-Rahman's book [9]) before stating the main theorem. Next some terminology is defined before stating the main theorem. We use the notation:

$$(a; q)_\infty = \prod_{k=0}^{\infty} (1 - aq^k), \quad (a; q)_n = \prod_{k=0}^{n-1} (1 - aq^k)$$

for a nonnegative integer $n \geq 0$. Usually $(a; q)_n$ is called the q -shifted factorial, and we frequently use the compact notation:

$$(a_1, a_2, \dots, a_r; q)_\infty = (a_1; q)_\infty (a_2; q)_\infty \cdots (a_r; q)_\infty,$$

$$(a_1, a_2, \dots, a_r; q)_n = (a_1; q)_n (a_2; q)_n \cdots (a_r; q)_n.$$

If we put $a = q^\alpha$ and $q \rightarrow 1$, then we have

$$\lim_{q \rightarrow 1} \frac{(q^\alpha; q)_n}{(1-q)^n} = (\alpha)_n,$$

where $(\alpha)_n = \prod_{k=0}^{n-1} (\alpha + k)$ is called the *raising factorial*. We shall define the ${}_{r+1}\phi_r$ basic hypergeometric series by

$${}_{r+1}\phi_r \left[\begin{matrix} a_1, a_2, \dots, a_{r+1} \\ b_1, \dots, b_r \end{matrix}; q, z \right] = \sum_{n=0}^{\infty} \frac{(a_1, a_2, \dots, a_{r+1}; q)_n}{(q, b_1, \dots, b_r; q)_n} z^n.$$

If we put $a_i = q^{\alpha_i}$ and $b_i = q^{\beta_i}$ in the above series and let $q \rightarrow 1$, then we obtain the ${}_{r+1}F_r$ hypergeometric series

$${}_{r+1}F_r \left[\begin{matrix} \alpha_1, \alpha_2, \dots, \alpha_{r+1} \\ \beta_1, \dots, \beta_r \end{matrix}; z \right] = \sum_{n=0}^{\infty} \frac{(\alpha_1)_n (\alpha_2)_n \cdots (\alpha_{r+1})_n}{n! (\beta_1)_n \cdots (\beta_r)_n} z^n.$$

The Motzkin number M_n is defined to be

$$M_n = {}_2F_1 \left[\begin{matrix} (1-n)/2, -n/2 \\ 2 \end{matrix}; 4 \right].$$

The generating function for the Motzkin numbers is given by

$$\sum_{n=0}^{\infty} M_n x^n = \frac{1-x-\sqrt{1-2x-3x^2}}{2x^2}.$$

It is known [1] that

$$(1.7) \quad \det (M_{i+j})_{0 \leq i, j \leq n-1} = 1$$

for $n \geq 1$, and

$$(1.8) \quad \det (M_{i+j+1})_{0 \leq i, j \leq n-1} = 1, 0, -1$$

for $n \equiv 0, 1 \pmod{6}$, $n \equiv 2, 5 \pmod{6}$, $n \equiv 3, 4 \pmod{6}$, respectively.

The large Schröder number S_n is defined to be

$$S_n = {}_2F_1 \left[\begin{matrix} -n+1, n+2 \\ 2 \end{matrix}; -1 \right]$$

for $n \geq 1$ ($S_0 = 1$). The generating function for the large Schröder numbers is

$$(1.9) \quad \sum_{n=0}^{\infty} S_n x^n = \frac{1-x-\sqrt{1-6x+x^2}}{2x}.$$

Eu and Fu [8] have proved

$$(1.10) \quad \det (S_{i+j})_{0 \leq i, j \leq n-1} = 2^{\binom{n}{2}}, \quad \det (S_{i+j+1})_{0 \leq i, j \leq n-1} = 2^{\binom{n+1}{2}}$$

for $n \geq 1$ (see [4, 8, 16]). We can also prove that

$$(1.11) \quad \det (S_{i+j+2})_{0 \leq i, j \leq n-1} = 2^{\binom{n+1}{2}} (2^{n+1} - 1)$$

holds for $n \geq 1$.

In this article, as a generalization of (1.2), we choose

$$(1.12) \quad \mu_n = \frac{(aq; q)_n}{(abq^2; q)_n}$$

for a nonnegative integer n . The aim of this article is to give three different proofs of the following theorem:

Theorem 1.1. *Let n be a positive integer. Then we have*

$$(1.13) \quad \det (\mu_{i+j})_{0 \leq i, j \leq n-1} = a^{\frac{1}{2}n(n-1)} q^{\frac{1}{6}n(n-1)(2n-1)} \prod_{k=1}^n \frac{(q, aq, bq; q)_{n-k}}{(abq^{n-k+1}; q)_{n-k} (abq^2; q)_{2(n-k)}}.$$

As a corollary of this theorem we can get the following more general identity.

Corollary 1.2. *Let n be a positive integer, and t a nonnegative integer. Then we have*

$$(1.14) \quad \det(\mu_{i+j+t})_{0 \leq i, j \leq n-1} = a^{\frac{1}{2}n(n-1)} q^{\frac{1}{6}n(n-1)(2n-1) + \frac{1}{2}n(n-1)t} \left\{ \frac{(aq; q)_t}{(abq^2; q)_t} \right\}^n \\ \times \prod_{k=1}^n \frac{(q, aq^{t+1}, bq; q)_{n-k}}{(abq^{n-k+t+1}; q)_{n-k} (abq^{t+2}; q)_{2(n-k)}}.$$

Proof. If we use

$$\mu_{n+t} = \frac{(aq; q)_{n+t}}{(abq^2; q)_{n+t}} = \frac{(aq; q)_t}{(abq^2; q)_t} \cdot \frac{(aq^{t+1}; q)_n}{(abq^{t+2}; q)_n},$$

then we have

$$\det(\mu_{i+j+t})_{0 \leq i, j \leq n-1} = \det \left(\frac{(aq; q)_t}{(abq^2; q)_t} \cdot \frac{(aq^{t+1}; q)_{i+j}}{(abq^{t+2}; q)_{i+j}} \right)_{0 \leq i, j \leq n-1} \\ = \left\{ \frac{(aq; q)_t}{(abq^2; q)_t} \right\}^n \det \left(\frac{(a'q; q)_{i+j}}{(a'bq^2; q)_{i+j}} \right)_{0 \leq i, j \leq n-1},$$

where $a' = aq^t$. If we use (1.13), then we obtain (1.14) by a straightforward computation. \square

We can prove (1.3), (1.4) and (1.5) as a corollary of Corollary 1.2.

Proof of (1.3), (1.4) and (1.5). If we substitute $a = q^\alpha$ and $b = q^\beta$ into ν_n , and we put $q \rightarrow 1$, then we obtain $\mu_n \rightarrow \frac{(\alpha+1)_n}{(\alpha+\beta+2)_n}$, which we write ν_n . Thus (1.14), leads to

$$\det(\nu_{i+j+t})_{0 \leq i, j \leq n-1} = \nu_t^n \prod_{k=1}^n \frac{(n-k)!(\alpha+t+1)_{n-k}(\beta+1)_{n-k}}{(\alpha+\beta+t+n-k+1)_{n-k}(\alpha+\beta+t+2)_{2(n-k)}}.$$

Note that

$$\nu_n = \begin{cases} C_n/2^{2n} & \text{if } \alpha = -\frac{1}{2} \text{ and } \beta = \frac{1}{2}, \\ B_n/2^{2n} & \text{if } \alpha = \frac{1}{2} \text{ and } \beta = -\frac{1}{2}, \\ D_n/2^{2n} & \text{if } \alpha = -\frac{1}{2} \text{ and } \beta = -\frac{1}{2}. \end{cases}$$

Hence we obtain

$$2^{2n(n+t-1)} \det(\nu_{i+j+t})_{0 \leq i, j \leq n-1} = \begin{cases} \det(C_{i+j+t})_{0 \leq i, j \leq n-1} & \text{if } \alpha = -\frac{1}{2} \text{ and } \beta = \frac{1}{2}, \\ \det(B_{i+j+t})_{0 \leq i, j \leq n-1} & \text{if } \alpha = \frac{1}{2} \text{ and } \beta = -\frac{1}{2}, \\ \det(D_{i+j+t})_{0 \leq i, j \leq n-1} & \text{if } \alpha = -\frac{1}{2} \text{ and } \beta = -\frac{1}{2}. \end{cases}$$

Thus we can prove (1.3), (1.4) and (1.5) by direct computations from the above identity. \square

In fact we can also obtain the following generalization of (1.6).

Theorem 1.3. *Let n be a positive integer, and k_0, \dots, k_{n-1} nonnegative integers. Then we have*

$$(1.15) \quad \det(\mu_{k_i+j})_{0 \leq i, j \leq n-1} = a^{\binom{n}{2}} q^{\binom{n+1}{3}} \prod_{i=0}^{n-1} \frac{(aq; q)_{k_i}}{(abq^2; q)_{k_i+n-1}} \prod_{0 \leq i < j \leq n-1} (q^{k_i} - q^{k_j}) \prod_{i=0}^{n-1} (bq; q)_i.$$

§ 2. Non-intersecting lattice paths

In this section we give our first proof of Theorem 1.1 using non-intersecting lattice paths.

Let m and n be nonnegative integers. A *Dyck path* is, by definition, a lattice path in the plane lattice \mathbb{Z}^2 consisting of two types of steps: rise vector $(1, 1)$ and fall vector $(1, -1)$, which never passes below the x -axis. We say a rise vector (resp. fall vector) whose origin is (x, y) and ends at $(x + 1, y + 1)$ (resp. $(x + 1, y - 1)$) has *height* y . For example, Figure 1 presents




Figure 1. A Dyck Path starting from $(0, 0)$ and ending at $(8, 2)$

a Dyck path starting from $(0, 0)$ and ending at $(8, 2)$, in which each red number stands for the height of the step. Let $\mathcal{D}_{m,n}$ denote the set of Dyck paths starting from $(0, 0)$ and ending at (m, n) . Especially, the cardinality of $\mathcal{D}_{2n,0}$ is known to be the Catalan number C_n .

A *Motzkin path* is, by definition, a lattice path in \mathbb{Z}^2 consisting of three types of steps: rise vectors $(1, 1)$, fall vectors $(1, -1)$, and (short) level vectors $(1, 0)$ which never passes below the x -axis. We say a rise vector, fall vector and level vector whose origin is (x, y) and ends at $(x + 1, y + 1)$, $(x + 1, y - 1)$ and $(x + 1, y)$ has *height* y , respectively. Figure 2 presents a




Figure 2. A Motzkin path starting from $(0, 0)$ and ending at $(9, 2)$

Motzkin path starting from $(0, 0)$ and ending at $(9, 2)$, in which each red number stands for the height of the step. Let $\mathcal{M}_{m,n}$ denote the set of Motzkin paths starting from $(0, 0)$ and ending at (m, n) . Note that the cardinality of $\mathcal{M}_{n,0}$ is known to be the Motzkin number M_n . We define the height of each step similarly as before.




Figure 3. A Schröder path starting from $(0,0)$ and ending at $(10,0)$

A *Schröder path* is, by definition, a lattice path in \mathbb{Z}^2 consisting of three types of steps: rise vectors $(1, 1)$, fall vectors $(1, -1)$, and long level vectors $(2, 0)$ which never passes below the x -axis. Figure 3 presents a Schröder path starting from $(0,0)$ and ending at $(10,0)$, in which each red number stands for the height of the step. Let $\mathcal{S}_{m,n}$ denote the set of Schröder paths starting from $(0,0)$ and ending at (m,n) . Note that the cardinality of $\mathcal{S}_{2n,0}$ is known to be the large Schröder number S_n .

Assign the weight a_h, b_h, c_h to each rise vector, fall vector, (short or long) level vector of height h , respectively. Set the weight of a path P to be the product of the weights of its edges and denote it by $w(P)$. Given any family \mathcal{F} of paths, we write the generating function of \mathcal{F} as

$$\text{GF} [\mathcal{F}] = \sum_{P \in \mathcal{F}} w(P).$$

Proposition 2.1. (Flajolet [7]) *The generating function for the Dyck paths is given by the following Stieltjes type continued fraction:*

$$\sum_{n \geq 0} \text{GF} [\mathcal{D}_{(2n,0)}] t^{2n} = \frac{1}{1 - \frac{a_0 b_1 t^2}{1 - \frac{a_1 b_2 t^2}{1 - \frac{a_2 b_3 t^2}{\ddots}}}}.$$

Meanwhile, the generating function for the Motzkin paths is given by the following Jacobi type continued fraction:

$$\sum_{n \geq 0} \text{GF} [\mathcal{M}_{(n,0)}] t^n = \frac{1}{1 - c_0 t - \frac{a_0 b_1 t^2}{1 - c_1 t - \frac{a_1 b_2 t^2}{1 - c_2 t - \frac{a_2 b_3 t^2}{\ddots}}}}.$$

It is also easy to see the following proposition holds.

Proposition 2.2. *Let n be a positive integer. Then the generating function for Schröder*

paths is given by the following continued fraction:

$$\sum_{n \geq 0} \text{GF} [\mathcal{S}_{(2n,0)}] t^{2n} = \frac{1}{1 - c_0 t^2 - \frac{a_0 b_1 t^2}{1 - c_1 t^2 - \frac{a_1 b_2 t^2}{1 - c_2 t^2 - \frac{a_2 b_3 t^2}{\ddots}}}}.$$

Next we recall notation and definitions used for the lattice path method due to Gessel and Viennot [10]. Let $D = (V, E)$ be an acyclic digraph without multiple edges. If u and v are any pair of vertices, let $\mathcal{P}(u, v)$ denote the set of all directed paths from u to v . For a fixed positive integer n , an n -vertex is an n -tuple of vertices of D . If $\mathbf{u} = (u_1, \dots, u_n)$ and $\mathbf{v} = (v_1, \dots, v_n)$ are n -vertices, an n -path from \mathbf{u} to \mathbf{v} is an n -tuple $\mathbf{P} = (P_1, \dots, P_n)$ such that $P_i \in \mathcal{P}(u_i, v_i)$, $i = 1, \dots, n$. The n -path $\mathbf{P} = (P_1, \dots, P_n)$ is said to be *non-intersecting* if any two different paths P_i and P_j have no vertex in common. We will write $\mathcal{P}(\mathbf{u}, \mathbf{v})$ for the set of all n -paths from \mathbf{u} to \mathbf{v} , and write $\mathcal{P}_0(\mathbf{u}, \mathbf{v})$ for the subset of $\mathcal{P}(\mathbf{u}, \mathbf{v})$ consisting of non-intersecting n -paths. If $\mathbf{u} = (u_1, \dots, u_m)$ and $\mathbf{v} = (v_1, \dots, v_n)$ are linearly ordered sets of vertices of D , then \mathbf{u} is said to be D -compatible with \mathbf{v} if every path $P \in \mathcal{P}(u_i, v_l)$ intersects with every path $Q \in \mathcal{P}(u_j, v_k)$ whenever $i < j$ and $k < l$. Let S_n denote the symmetric group on $\{1, 2, \dots, n\}$. Then for $\pi \in S_n$, by \mathbf{v}^π we mean the n vertex $(v_{\pi(1)}, \dots, v_{\pi(n)})$.

The weight $w(\mathbf{P})$ of an n -path \mathbf{P} is defined to be the product of the weights of its components. Thus, if $\mathbf{u} = (u_1, \dots, u_n)$ and $\mathbf{v} = (v_1, \dots, v_n)$ are n -vertices, we define the generating functions $F(\mathbf{u}, \mathbf{v}) = \text{GF} [\mathcal{P}(\mathbf{u}, \mathbf{v})] = \sum_{\mathbf{P} \in \mathcal{P}(\mathbf{u}, \mathbf{v})} w(\mathbf{P})$ and $F_0(\mathbf{u}, \mathbf{v}) = \text{GF} [\mathcal{P}_0(\mathbf{u}, \mathbf{v})] = \sum_{\mathbf{P} \in \mathcal{P}_0(\mathbf{u}, \mathbf{v})} w(\mathbf{P})$. In particular, if u and v are any pair of vertices, we write

$$h(u, v) = \text{GF} [\mathcal{P}(u, v)] = \sum_{P \in \mathcal{P}(u, v)} w(P).$$

The following lemma is called the Gessel-Viennot formula for counting lattice paths in terms of determinants. (See [10].)

Lemma 2.3. (*Lidström-Gessel-Viennot*)

Let $\mathbf{u} = (u_1, \dots, u_n)$ and $\mathbf{v} = (v_1, \dots, v_n)$ be two n -vertices in an acyclic digraph D . Then

$$(2.1) \quad \sum_{\pi \in S_n} \text{sgn} \pi F_0(\mathbf{u}^\pi, \mathbf{v}) = \det[h(u_i, v_j)]_{1 \leq i, j \leq n}.$$

In particular, if \mathbf{u} is D -compatible with \mathbf{v} , then

$$(2.2) \quad F_0(\mathbf{u}, \mathbf{v}) = \det[h(u_i, v_j)]_{1 \leq i, j \leq n}.$$

If we apply Lemma 2.3 to Dyck paths, then we obtain the following proposition:

Proposition 2.4. Let $G_m = \text{GF} [\mathcal{D}_{2m,0}]$ for non-negative integer m .

(i) If $t = 0$, then we have

$$(2.3) \quad \det (G_{i+j})_{0 \leq i, j \leq n-1} = \prod_{i=1}^n (a_{2i-2} b_{2i-1} a_{2i-1} b_{2i})^{n-i}.$$

(ii) If $t = 1$, then we have

$$(2.4) \quad \det (G_{i+j+1})_{0 \leq i, j \leq n-1} = \prod_{i=1}^n (a_{2i-2} b_{2i-1})^{n-i+1} (a_{2i-1} b_{2i})^{n-i}.$$

(iii) If $t = 2$, then we have $\det (G_{i+j+2})_{0 \leq i, j \leq n-1}$ equals

$$(2.5) \quad \sum_{k=0}^n \prod_{i=1}^k (a_0 a_1 \cdots a_{2i-3} a_{2i-2}^2 b_1 b_2 \cdots b_{2i-1} b_{2i-1}^2) \cdot \prod_{i=1}^k (a_0 a_1 \cdots a_{2i-1} b_1 b_2 \cdots b_{2i}).$$

(iv) If $t = 3$, then we have $\det (G_{i+j+3})_{0 \leq i, j \leq n-1}$ equals

$$(2.6) \quad \sum_{k=0}^n \left\{ \sum_{l=0}^k \prod_{i=1}^l (a_0 a_1 \cdots a_{2i-3} a_{2i-2}^2 b_{2i-1}) \prod_{i=l+1}^k (a_0 a_1 \cdots a_{2i-3} a_{2i-2} a_{2i-1} b_{2i}) \right\} \\ \times \left\{ \sum_{l=0}^k \prod_{i=1}^l (b_1 b_2 \cdots b_{2i-2} b_{2i-1}^2 a_{2i-2}) \prod_{i=l+1}^k (b_1 b_2 \cdots b_{2i-2} b_{2i-1} b_{2i} a_{2i-1}) \right\} \\ \times \prod_{i=k+1}^n (a_0 a_1 \cdots a_{2i-1} a_{2i} b_1 b_2 \cdots b_{2i} b_{2i+1}).$$

(i) and (ii) of this proposition are originally appeared in [18, Ch. 4, §3].

Proof. We consider the digraph (V, E) , in which V is the plane lattice \mathbb{Z}^2 and E the set of rise vectors and fall vectors in the above half plane. Let $u_i = (x_0 - 2(i-1), 0)$ and $v_j = (x_0 + 2(j+t-1), 0)$ for $i, j = 1, 2, \dots, n$, $t = 0, 1, 2, 3$ and a fixed integer x_0 . It is easy to see that the n -vertex $\mathbf{u} = (u_1, \dots, u_n)$ is D -compatible with the n -vertex $\mathbf{v} = (v_1, \dots, v_n)$. If $t = 0$, then there is always a unique n -path $\mathbf{P} = (P_1, \dots, P_n)$ that connect \mathbf{u} to \mathbf{v} as in





Figure 4. $t = 0$ and $n = 4$

Figure 4. By multiplying the weights of all edges in \mathbf{P} , we obtain the right-hand side of (2.3). On the other hand, applying Lemma 2.3, we obtain the left-hand side of (2.3).

The other identities can be proven similarly. For example, if $t = 1$, there is only one n -path $\mathbf{P} = (P_1, \dots, P_n)$ that connect \mathbf{u} to \mathbf{v} as in Figure 5. As the product of the weights of all edges in \mathbf{P} we obtain (2.4). If $t = 2$, there are $(n+1)$ ways to connect \mathbf{u} to \mathbf{v} with n -path $\mathbf{P} = (P_1, \dots, P_n)$. As an example, we show one way in Figure 6. A similar reasoning leads to (2.5). One can also derive (2.6) by a similar argument. \square

We assign the following weight to each step: the weight of a rise vector is 1, while the weight of a fall vector of height h is

$$(2.7) \quad \lambda_h = \begin{cases} \frac{q^k(1-aq^{k+1})(1-abq^{k+1})}{(1-abq^{2k+1})(1-abq^{2k+2})} & \text{if } h = 2k+1 \text{ is odd,} \\ \frac{aq^k(1-q^k)(1-bq^k)}{(1-abq^{2k})(1-abq^{2k+1})} & \text{if } h = 2k \text{ is even.} \end{cases}$$


Figure 5. $t = 1$ and $n = 4$

For example, we have $\lambda_1 = \frac{1-aq}{1-abq^2}$, $\lambda_2 = \frac{aq(1-q)(1-bq)}{(1-abq^2)(1-abq^3)}$, $\lambda_3 = \frac{q(1-aq^2)(1-abq^2)}{(1-abq^3)(1-abq^4)}$, and an example of the weight of a path is Figure 7.

Lemma 2.5. *Let m and n be a non-negative integers such that $m \equiv n \pmod{2}$. Then the generating function of $\mathcal{D}_{m,n}$ is given by*

$$(2.8) \quad \text{GF}(\mathcal{D}_{m,n}) = \left[\begin{matrix} \lfloor \frac{m}{2} \rfloor \\ \lfloor \frac{n}{2} \rfloor \end{matrix} \right]_q \frac{(aq^{1+\lceil \frac{n}{2} \rceil}; q)_{\frac{m-n}{2}}}{(abq^{2+n}; q)_{\frac{m-n}{2}}}.$$

Here $\lfloor x \rfloor$ (resp. $\lceil x \rceil$) stands for the greatest integer that does not exceed x (resp. the smallest integer that is not smaller than x). Especially, we have

$$(2.9) \quad \text{GF}(\mathcal{D}_{2n,0}) = \frac{(aq; q)_n}{(abq^2; q)_n}.$$

Proof. We prove (2.8) by induction on m . If $m = 0$, then it is obvious that $\text{GF}(\mathcal{D}_{0,n})$ equals 1 if $n = 0$, and 0 otherwise. Assume that (2.8) holds up to $m - 1$. Then we have

$$\text{GF}[\mathcal{D}_{m,n}] = \text{GF}[\mathcal{D}_{m-1,n-1}] + \lambda_{n+1} \text{GF}[\mathcal{D}_{m-1,n+1}].$$

If $m = 2r$ and $n = 2s$, then, by induction hypothesis and the above recursion, we obtain $\text{GF}[\mathcal{D}_{2r,2s}]$ equals

$$\begin{aligned} & \left[\begin{matrix} r-1 \\ s-1 \end{matrix} \right]_q \frac{(aq^{s+1}; q)_{r-s}}{(abq^{2s+1}; q)_{r-s}} + \frac{q^s(1-aq^{s+1})(1-abq^{s+1})}{(1-abq^{2s+1})(1-abq^{2s+2})} \left[\begin{matrix} r-1 \\ s \end{matrix} \right]_q \frac{(aq^{s+2}; q)_{r-s-1}}{(abq^{2s+3}; q)_{r-s-1}} \\ &= \frac{(q; q)_{r-1} (aq^{s+1}; q)_{r-s}}{(q; q)_s (q; q)_{r-s} (abq^{2s+1}; q)_{r-s+1}} \{ (1-q^s)(1-abq^{r+s+1}) + q^s(1-q^{r-s})(1-abq^{s+1}) \} \\ &= \left[\begin{matrix} r \\ s \end{matrix} \right]_q \frac{(aq^{s+1}; q)_{r-s}}{(abq^{2s+2}; q)_{r-s}}. \end{aligned}$$

This equals the right-hand side of (2.8) with $m = 2r$ and $n = 2s$. Hence (2.8) holds when $m = 2r$. One can prove (2.8) similarly when $m = 2r + 1$ and $n = 2s + 1$. \square

For example, if $m = 4$ and $n = 0$, then $\mathcal{D}_{4,0}$ has the two Dyck paths shown in Figure 9. Thus, the generating function of $\mathcal{D}_{4,0}$ equals

$$\text{GF}(\mathcal{D}_{4,0}) = \lambda_1^2 + \lambda_1\lambda_2 = \frac{(1-aq)(1-aq^2)}{(1-abq^2)(1-abq^3)}.$$

Proof of Theorem 1.1. If we use (2.3), (2.7) and (2.9), then we conclude that $\det(\mu_{i+j})_{0 \leq i, j \leq n-1}$ equals

$$\prod_{i=1}^n (\lambda_{2i-1}\lambda_{2i})^{n-i} = \prod_{i=1}^n \left\{ \frac{aq^{2i-1}(1-q^i)(1-aq^i)(1-bq^i)(1-abq^i)}{(1-abq^{2i-1})(1-abq^{2i})^2(1-abq^{2i+1})} \right\}^{n-i}.$$

An easy computation leads to (1.13). □

Remark. One can also prove Theorem 1.1 by using Motzkin paths and giving the weight λ_{2h+1} to rise vector of height h , λ_{2h} to fall vector of height h and $\lambda_{2h} + \lambda_{2h+1}$ to level vector of height h . Then one can prove

$$(2.10) \quad \text{GF}(\mathcal{M}_{m,n}) = q^{\binom{m}{n}} \begin{bmatrix} m \\ n \end{bmatrix}_q \frac{(aq; q)_m (1-abq^{2n+1})}{(abq^{n+1}; q)_{m+1}}.$$

for nonnegative integers m and n .

§ 3. Orthogonal Polynomials

In this section we give our second proof of Theorem 1.1 using the little q -Jacobi polynomials. We use the notation $S(t; \lambda_1, \lambda_2, \dots)$ for the Stieltjes-type continued fraction

$$(3.1) \quad \frac{1}{1 - \frac{\lambda_1 t}{1 - \frac{\lambda_2 t}{1 - \dots}}},$$

and $J(t; b_0, b_1, b_2, \dots; \lambda_1, \lambda_2, \dots)$ for the Jacobi-type continued fraction

$$(3.2) \quad \frac{1}{1 - b_0 x - \frac{\lambda_1 x^2}{1 - b_1 x - \frac{\lambda_2 x^2}{1 - b_2 x - \dots}}}$$

Given a moment sequence $\{\mu_n\}$, we define the linear functional $\mathcal{L} : x^n \mapsto \mu_n$ on the vector space of polynomials $\mathbb{C}[x]$. Then the monic polynomials $p_n(x)$ orthogonal with respect to \mathcal{L} and of $\deg p_n(x) = n$ satisfy a three term recurrence relation (Favard's theorem), say

$$(3.3) \quad p_{n+1}(x) = (x - b_n)p_n(x) - \lambda_n p_{n-1}(x),$$

where $p_{-1}(x) = 0$ and $p_0(x) = 1$. The moment sequence $\{\mu_n\}$ is related to the coefficients b_n and λ_n by the identity:

$$(3.4) \quad 1 + \sum_{n \geq 1} \mu_n x^n = J(t; b_0, b_1, b_2, \dots; \lambda_1, \lambda_2, \dots).$$

Hereafter we assume $\lambda_0 = \mu_0 = 1$ for simplicity of arguments.

Define Δ_n and $D_n(x)$ by

$$\Delta_n = \begin{vmatrix} \mu_0 & \mu_1 & \dots & \mu_n \\ \mu_1 & \mu_2 & \dots & \mu_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_n & \mu_{n+1} & \dots & \mu_{2n} \end{vmatrix}, \quad D_n(x) = \begin{vmatrix} \mu_0 & \mu_1 & \dots & \mu_n \\ \mu_1 & \mu_2 & \dots & \mu_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_{n-1} & \mu_n & \dots & \mu_{2n-1} \\ 1 & x & \dots & x^n \end{vmatrix}.$$

Then $p_n(x) = (\Delta_{n-1})^{-1} D_n(x)$ is the monic OPS for \mathcal{L} .

It is easy to see that

$$(3.5) \quad \mathcal{L}(x^n p_n(x)) = \frac{\Delta_n}{\Delta_{n-1}} = \lambda_n \lambda_{n-1} \dots \lambda_1 \mu_0,$$

$$(3.6) \quad \mathcal{L}(x^{n+1} p_n(x)) = \frac{\chi_n}{\Delta_{n-1}} = \lambda_n \lambda_{n-1} \dots \lambda_1 \mu_0 (b_0 + \dots + b_n),$$

where

$$\chi_n = \begin{vmatrix} \mu_0 & \mu_1 & \dots & \mu_n \\ \mu_1 & \mu_2 & \dots & \mu_{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ \mu^{n-1} & \mu^n & \dots & \mu_{2n-1} \\ \mu_{n+1} & \mu_{n+2} & \dots & \mu_{2n+1} \end{vmatrix}.$$

Therefore

$$(3.7) \quad \lambda_n = \frac{\mathcal{L}[p_n^2(x)]}{\mathcal{L}[p_{n-1}^2(x)]} = \frac{\Delta_{n-2} \Delta_n}{\Delta_{n-1}^2},$$

and

$$(3.8) \quad b_n = \frac{\mathcal{L}[x p_n^2(x)]}{\mathcal{L}[p_n^2(x)]} = \frac{\chi_n}{\Delta_n} - \frac{\chi_{n-1}}{\Delta_{n-1}}.$$

Theorem 3.1 (The Stieltjes-Rogers addition formula). *The formal power series $f(x) = \sum_{i \geq 0} a_i x^i / i!$ ($a_0 = 1$) has the property that*

$$f(x+y) = \sum_{m \geq 0} \alpha_m f_m(x) f_m(y),$$

where α_m is independent of x and y and

$$f_m(x) = \frac{x^m}{m!} + \beta_m \frac{x^{m+1}}{(m+1)!} + O(x^{m+2}),$$

if and only if the formal power series $\hat{f}(x) = \sum_{i \geq 0} a_i x^i$ has the J -continued fraction expansion $J(x; b_0, b_1, b_2, \dots; \lambda_1, \lambda_2, \dots)$ with the parameters

$$b_m = \beta_{m+1} - \beta_m \quad \text{and} \quad \lambda_m = \frac{\alpha_{m+1}}{\alpha_m}, \quad m \geq 0.$$

From (3.5), one can compute the Hankel determinants

$$(3.9) \quad \det (\mu_{i+j})_{0 \leq i, j \leq n-1} = \Delta_{n-1} = \mu_0^n \lambda_1^{n-1} \lambda_2^{n-2} \cdots \lambda_{n-2}^2 \lambda_{n-1},$$

of (1.13) by taking appropriate orthogonal polynomials $p_n(x)$. Recall the definition of Heine's q -hypergeometric series

$${}_2\phi_1(a, b; c; q; x) = \sum_{n=0}^{\infty} \frac{(a; q)_n (b; q)_n}{(c; q)_n} \frac{x^n}{(q; q)_n}.$$

The following is one of Heine's three-term contiguous relations for ${}_2\phi_1$:

$${}_2\phi_1(a, b; c; q; x) = {}_2\phi_1(a, bq; cq; q; x) + \frac{(1-a)(c-b)}{(1-c)(1-cq)} x {}_2\phi_1(aq, bq; cq^2; q; x).$$

It follows that

$$\frac{{}_2\phi_1(a, bq; cq; q; x)}{{}_2\phi_1(a, b; c; q; x)} = S \left(x; \frac{(1-a)(b-c)}{(1-c)(1-cq)}, \frac{(1-bq)(a-cq)}{(1-cq)(1-cq^2)}, \frac{(1-aq)(bq-cq^2)}{(1-cq^2)(1-cq^3)}, \dots \right).$$

Hence, by induction, we can prove that

$$\frac{{}_2\phi_1(a, bq; cq; q; x)}{{}_2\phi_1(a, b; c; q; x)} = S(x; \lambda_1, \lambda_2, \dots),$$

where

$$\lambda_{2n+1} = \frac{(1-aq^n)(b-cq^n)q^n}{(1-cq^{2n})(1-cq^{2n+1})}, \quad \lambda_{2n} = \frac{(1-bq^n)(a-cq^n)q^{n-1}}{(1-cq^{2n-1})(1-cq^{2n})}.$$

Making the substitution $b \leftarrow 1$, $a \leftarrow aq$ and $c \leftarrow abq$ into the above equation, we obtain

$$\sum_{n \geq 0} \frac{(aq; q)_n}{(abq^2; q)_n} x^n = S(x; \lambda_1, \lambda_2, \dots),$$

where

$$\lambda_{2n+1} = \frac{(1-aq^{n+1})(1-abq^{n+1})q^n}{(1-aq^{2n+1})(1-abq^{2n+2})}, \quad \lambda_{2n} = \frac{(1-q^n)(1-bq^n)aq^n}{(1-abq^{2n})(1-abq^{2n+1})}.$$

This corresponds to the little q -Jacobi polynomials. Indeed, the little q -Jacobi polynomials

$$(3.10) \quad p_n(x; a, b; q) = \frac{(aq; q)_n}{(abq^{n+1}; q)_n} (-1)^n q^{\binom{n}{2}} {}_2\phi_1 \left[\begin{matrix} q^{-n}, abq^{n+1} \\ aq \end{matrix}; q, xq \right]$$

are introduced in [2]. The polynomials satisfy the recurrence equation

$$(3.11) \quad xp_n(x) = p_{n+1}(x) + (A_n + C_n)p_n(x) + A_{n-1}C_{n-1}p_{n-1}(x)$$

where $p_{-1}(x) = 0$, $p_0(x) = 1$ and

$$(3.12) \quad A_n = \frac{q^n(1-aq^{n+1})(1-abq^{n+1})}{(1-abq^{2n+1})(1-abq^{2n+2})}, \quad C_n = \frac{aq^n(1-q^n)(1-bq^n)}{(1-abq^{2n})(1-abq^{2n+1})}.$$

They are orthogonal with respect to the moment sequence $\{\mu_n\}_{n \geq 0}$ where

$$(3.13) \quad \mu_n = \frac{(aq; q)_n}{(abq^2; q)_n}.$$

For the passage from the Stieltjes-type continued fraction to the Jacobi-type continued fraction we use the following contraction formula:

$$S(x, \lambda_1, \lambda_2, \dots) = \frac{1}{1 - \lambda_1 t - \frac{\lambda_1 \lambda_2 t^2}{1 - (\lambda_2 + \lambda_3)t - \frac{\lambda_3 \lambda_4 t^2}{1 - \dots}}}.$$

Thus, by the same computation as in the former section, we conclude that the determinant (3.13) is equal to (1.13). This proof gives us an insight to the determinant (1.13) from the point of view of the classical orthogonal polynomial theory.

§ 4. q -Dougall's formula

In this section we give our third proof of Theorem 1.1 using q -Dougall's formula and LU-decomposition of the Hankel matrix.

First the following formula is known as q -Dougall's formula: We have

$$(4.1) \quad {}_6\phi_5 \left[\begin{matrix} a, qa^{\frac{1}{2}}, -qa^{\frac{1}{2}}, b, c, d \\ a^{\frac{1}{2}}, -a^{\frac{1}{2}}, aq/b, aq/c, aq/d \end{matrix}; q, \frac{aq}{bcd} \right] = \frac{(qa, aq/bc, aq/bd, aq/cd; q)_\infty}{(aq/b, aq/c, aq/d, aq/bcd; q)_\infty}.$$

provided $|aq/bcd| < 1$ (see [9, (2.7.1)]). If we perform the substitution $a \leftarrow abq$, $b \leftarrow bq$, $c \leftarrow q^{-i}$ and $d \leftarrow q^{-j}$ in (4.1), then we obtain

$$(4.2) \quad {}_6\phi_5 \left[\begin{matrix} abq, a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{3}{2}}, -a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{3}{2}}, bq, q^{-i}, q^{-j} \\ a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{1}{2}}, -a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{1}{2}}, aq, abq^{i+2}, abq^{j+2} \end{matrix}; q, aq^{i+j+1} \right] = \frac{\mu_{i+j}}{\mu_i \mu_j},$$

where $\mu_n = \frac{(aq; q)_n}{(abq^2; q)_n}$ as before. If we use

$$(q^{-n}; q)_k = \frac{(q; q)_n}{(q; q)_{n-k}} (-1)^k q^{\binom{k}{2} - nk}$$

then this identity can be rewritten as

$$(4.3) \quad \sum_{k=0}^{\infty} a^k q^{k^2} \begin{bmatrix} i \\ k \end{bmatrix}_q \begin{bmatrix} j \\ k \end{bmatrix}_q \frac{(q, abq, a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{3}{2}}, -a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{3}{2}}, bq; q)_k}{(a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{1}{2}}, -a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{1}{2}}, aq, abq^{i+2}, abq^{j+2}; q)_k} = \frac{\mu_{i+j}}{\mu_i \mu_j}.$$

If we put

$$(4.4) \quad l_{ij} = \frac{\mu_i (abq^{j+2}; q)_j}{\mu_j (abq^{i+2}; q)_j} \begin{bmatrix} i \\ j \end{bmatrix}_q,$$

$$(4.5) \quad u_{ij} = a^i q^{i^2} \mu_i \mu_j \begin{bmatrix} j \\ i \end{bmatrix}_q \frac{(q, abq, a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{3}{2}}, -a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{3}{2}}, bq; q)_i}{(a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{1}{2}}, -a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{1}{2}}, aq, abq^{i+2}, abq^{j+2}; q)_i},$$

then (4.3) implies

$$(4.6) \quad \sum_{k=0}^{\infty} l_{ik} u_{kj} = \mu_{i+j}$$

Note that $L_n = (l_{ij})_{0 \leq i, j \leq n-1}$ is a lower triangular matrix such that all main-diagonal entries are 1, and $U_n = (u_{ij})_{0 \leq i, j \leq n-1}$ is an upper-triangular matrix with diagonal entries

$$(4.7) \quad \begin{aligned} u_{ii} &= a^i q^{i^2} \mu_i^2 \frac{(q, abq, a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{3}{2}}, -a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{3}{2}}, bq; q)_i}{(a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{1}{2}}, -a^{\frac{1}{2}} b^{\frac{1}{2}} q^{\frac{1}{2}}, aq, abq^{i+2}, abq^{i+2}; q)_i} \\ &= a^i q^{i^2} \frac{1 - abq^{2i+1}}{1 - abq} \frac{(q, aq, bq, abq; q)_i}{(abq^2; q)_{2i}^2}. \end{aligned}$$

Since $(\mu_{i+j})_{0 \leq i, j \leq n-1} = L_n U_n$, $\det (\mu_{i+j})_{0 \leq i, j \leq n-1}$ is the product of the diagonal entries, i.e.,

$$\det (\mu_{i+j})_{0 \leq i, j \leq n-1} = \prod_{i=0}^{n-1} u_{ii}.$$

Using (4.7), one can easily prove (1.13) by a direct computation.

Remark. We should note that Corollary 1.2 can be proven by induction using the following Desnanot-Jacobi adjoint matrix theorem: If M is an $n \times n$ matrix, then we have

$$(4.8) \quad \det M \det M_{1,n}^{1,n} = \det M_1^1 \det M_n^n - \det M_n^1 \det M_1^n,$$

where $M_{j_1, \dots, j_r}^{i_1, \dots, i_r}$ denotes the $(n-r) \times (n-r)$ submatrix obtained by removing rows i_1, \dots, i_r and columns j_1, \dots, j_r from M .

Corollary 1.2 can be also proven as a special case of Theorem 1.3, which will be proven in Section 5.1. In fact, if one puts $k_i = i + t$ in (1.15), then he obtains (1.14).

§ 5. Miscellany

§ 5.1. A proof of Theorem 1.3

In this subsection we give a proof of Theorem 1.3. Before we prove the formula, we need to cite a lemma from [11, 12].

Lemma 5.1 (Krattenthaler [11]). *Let $X_0, \dots, X_n, A_1, \dots, A_{n-1}$, and B_1, \dots, B_{n-1} be indeterminates. Then there holds*

$$(5.1) \quad \det \left[\prod_{l=1}^j (X_i + B_l) \prod_{l=j+1}^{n-1} (X_i + A_l) \right]_{0 \leq i, j \leq n-1} = \prod_{0 \leq i < j \leq n-1} (X_i - X_j) \prod_{1 \leq i \leq j \leq n-1} (B_i - A_j).$$

Proof of Theorem 1.3. Using

$$\mu_n = \frac{(aq; q)_n}{(abq^2; q)_n},$$

we can write

$$\begin{aligned} \det (\mu_{k_i+j})_{0 \leq i, j \leq n-1} &= \prod_{i=0}^{n-1} \frac{(aq; q)_{k_i}}{(abq^2; q)_{k_i+n-1}} \det \left(\prod_{l=1}^j (1 - aq^{k_i+l}) \prod_{l=j+1}^{n-1} (1 - abq^{k_i+l+1}) \right)_{0 \leq i, j \leq n-1} \\ &= \prod_{i=0}^{n-1} \frac{q^{(n-1)k_i} (aq; q)_{k_i}}{(abq^2; q)_{k_i+n-1}} \det \left(\prod_{l=1}^j (q^{-k_i} - aq^l) \prod_{l=j+1}^{n-1} (q^{-k_i} - abq^{l+1}) \right)_{0 \leq i, j \leq n-1}. \end{aligned}$$

If we substitute $X_i = q^{-k_i}$, $B_l = -aq^l$ and $A_l = -abq^{l+1}$ into (5.1), then we see that

$$\det (\mu_{k_i+j})_{0 \leq i, j \leq n-1} = \prod_{i=0}^{n-1} \frac{q^{(n-1)k_i} (aq; q)_{k_i}}{(abq^2; q)_{k_i+n-1}} \prod_{0 \leq i < j \leq n-1} (q^{-k_i} - q^{-k_j}) \prod_{1 \leq i \leq j \leq n-1} (abq^{j+1} - aq^i)$$

One can derive (1.15) easily by a direct computation. \square

§ 5.2. An addition formula for ${}_2F_1$

In this subsection we give a new proof of (1.7) using an addition formula for ${}_2F_1$ and LU-decomposition of Motzkin Hankel matrices. First, we shall prove the following identity.

Lemma 5.2. *If i and j are nonnegative integers, then we have*

$$(5.2) \quad \sum_{k \geq 0} \binom{i}{k} \binom{j}{k} {}_2F_1 \left[\begin{matrix} \frac{k-i+1}{2}, \frac{k-i}{2} \\ k+2 \end{matrix}; 4 \right] {}_2F_1 \left[\begin{matrix} \frac{k-j+1}{2}, \frac{k-j}{2} \\ k+2 \end{matrix}; 4 \right] = {}_2F_1 \left[\begin{matrix} \frac{1-i-j}{2}, \frac{-i-j}{2} \\ 2 \end{matrix}; 4 \right].$$

Proof. Recall the quadratic transformation formula (see [9, (3.1.5)]):

$$(5.3) \quad (1-z)^a {}_2F_1(a, b; 2b; 2z) = {}_2F_1 \left(\frac{a}{2}, \frac{a+1}{2}; b + \frac{1}{2}; \frac{z^2}{(1-z)^2} \right).$$

Applying (5.3) with $a = k - i$, $b = k + 3/2$ and $z = 2$ we obtain

$${}_2F_1 \left[\begin{matrix} \frac{k-i+1}{2}, \frac{k-i}{2} \\ k+2 \end{matrix}; 4 \right] = (-1)^{k-i} {}_2F_1 \left[\begin{matrix} k-i, k+3/2 \\ 2k+3 \end{matrix}; 4 \right].$$

Substituting i by j yields

$${}_2F_1 \left[\begin{matrix} \frac{k-j+1}{2}, \frac{k-j}{2} \\ k+2 \end{matrix}; 4 \right] = (-1)^{k-j} {}_2F_1 \left[\begin{matrix} k-j, k+3/2 \\ 2k+3 \end{matrix}; 4 \right].$$

Now, applying (5.3) with $a = -i - j$, $b = 3/2$ and $z = 2$ we obtain

$${}_2F_1 \left[\begin{matrix} \frac{1-i-j}{2}, \frac{-i-j}{2} \\ 2 \end{matrix}; 4 \right] = (-1)^{i+j} {}_2F_1 \left[\begin{matrix} -i-j, \frac{3}{2} \\ 3 \end{matrix}; 4 \right].$$

Therefore we can rewrite (5.2) as follows:

$$(5.4) \quad \sum_{k \geq 0} \binom{i}{k} \binom{j}{k} {}_2F_1 \left[\begin{matrix} k-i, k+3/2 \\ 2k+3 \end{matrix}; 4 \right] {}_2F_1 \left[\begin{matrix} k-j, k+3/2 \\ 2k+3 \end{matrix}; 4 \right] = {}_2F_1 \left[\begin{matrix} -i-j, \frac{3}{2} \\ 3 \end{matrix}; 4 \right].$$

Now we recall a formula of Burchnall and Chaundy [3, (43)]:

$$(5.5) \quad {}_2F_1 \left[\begin{matrix} c-a, c-b \\ c \end{matrix} ; x \right] = \sum_{k \geq 0} \frac{(c-a)_k (a)_k (d)_k (c-b-d)_k}{k! (c+k-1)_k (c)_{2k}} x^{2k} \\ \times {}_2F_1 \left[\begin{matrix} c-a+k, c-b-d+k \\ c+2k \end{matrix} ; x \right] {}_2F_1 \left[\begin{matrix} c-a+k, d+k \\ c+2k \end{matrix} ; x \right].$$

It is then easy to check that the specialization of (5.5) with

$$a = \frac{3}{2}, \quad b = 3 + i + j, \quad c = 3, \quad d = -j, \quad x = 4$$

yields (5.4). □

Proof of (1.7). Define l_{ij} and u_{ij} by

$$(5.6) \quad l_{ij} = \binom{i}{j} {}_2F_1 \left[\begin{matrix} \frac{j-i+1}{2}, \frac{j-i}{2} \\ j+2 \end{matrix} ; 4 \right],$$

$$(5.7) \quad u_{ij} = \binom{j}{i} {}_2F_1 \left[\begin{matrix} \frac{i-j+1}{2}, \frac{i-j}{2} \\ i+2 \end{matrix} ; 4 \right].$$

Then $L_n = (l_{ij})_{0 \leq i, j \leq n-1}$ is a lower triangular matrix with all diagonal entries 1, and $U_n = (u_{ij})_{0 \leq i, j \leq n-1}$ is an upper triangular matrix with all diagonal entries 1. The formula (5.2) gives the LU-decomposition of Motzkin Hankel matrix:

$$(M_{ij}) = L_n U_n.$$

Hence we conclude that $\det(M_{ij}) = 1$. □

§ 5.3. A q -analogue of Schröder numbers

We define $S_n(q)$ ($n \geq 0$) by the following recurrence:

$$S_0(q) = 1, \quad S_n(q) = q^{2n-1} S_{n-1}(q) + \sum_{k=0}^{n-1} q^{2(k+1)(n-1-k)} S_{n-1-k}(q) S_k(q).$$

In fact one can show that

$$S_n(q) = \sum_{P \in \mathcal{S}_{2n,0}} \omega(P),$$

where $\omega(P)$ is the number of triangles below the path P (see Figure 10), and the sum runs over all Schröder paths from the origin to $(2n, 0)$. As a q -analogue of (1.10) and (1.11) we consider the matrix

$$(5.8) \quad S_n^{(t)}(q) = \left(q^{(i-j)(i-j-1)} S_{i+j+t}(q) \right)_{0 \leq i, j \leq n-1}.$$

Note that this matrix is not a Hankel matrix, but as a q -analogue of (1.10) and (1.11), the following theorem holds:

Theorem 5.3. *Let n be a positive integer.*

(i) *If $t = 0$ or 1 , then we have*

$$(5.9) \quad \det S_{n-1}^{(1)}(q) = \det S_n^{(0)}(q) = \prod_{k=1}^{n-1} (q^{2k-1} + 1)^{n-k}.$$

(ii) *If $t = 2$, then we have*

$$(5.10) \quad \det S_n^{(2)}(q) = q^{-1} \prod_{k=1}^n (q^{2k-1} + 1)^{n+1-k} \left(\prod_{k=1}^{n+1} (q^{2k-1} + 1) - 1 \right).$$

To prove this theorem, we define the matrices

$$\begin{aligned} \widehat{S}_n^{(t)}(q) &= \left(q^{2(n-i)(t+i+j-2)} S_{t+i+j-2}(q) \right)_{1 \leq i, j \leq n}, \\ \widetilde{S}_n^{(t)}(q) &= \left(q^{-(t+i+j)(t+i+j-1)} S_{t+i+j-2}(q) \right)_{1 \leq i, j \leq n}, \end{aligned}$$

then the following lemma can be easily proven by direct computations:

Lemma 5.4. *Let n be a positive integer. Then*

$$(5.11) \quad \det \widehat{S}_n^{(t)}(q) = q^{\frac{n(n-1)(2n+3t-4)}{3}} \det S_n^{(t)}(q),$$

$$(5.12) \quad \det \widetilde{S}_n^{(t)}(q) = q^{-\frac{n(4n^2+3(2t+1)n+3t^2+3t-1)}{3}} \det S_n^{(t)}(q).$$

Lemma 5.5. *Let n be a positive integer.*

(i) *If $n \geq 2$, then we have*

$$(5.13) \quad \det \widehat{S}_n^{(0)}(q) = q^{(n-1)(n-2)} \det \widehat{S}_{n-1}^{(1)}(q).$$

(ii) *If $n \geq 2$, then we have*

$$(5.14) \quad \det \widehat{S}_n^{(1)}(q) = q^{n^2-n+1} \det \widehat{S}_{n-1}^{(2)}(q) + q^{2(n-1)^2} \det \widehat{S}_{n-1}^{(1)}(q).$$

(iii) *If $n \geq 3$, then we have*

$$(5.15) \quad \det \widetilde{S}_n^{(0)}(q) \det \widetilde{S}_{n-2}^{(2)}(q) = \det \widetilde{S}_{n-1}^{(0)}(q) \det \widetilde{S}_{n-1}^{(2)}(q) - \left\{ \det \widetilde{S}_{n-1}^{(1)}(q) \right\}^2.$$

Proof. We consider the digraph (V, E) , in which V is the plane lattice \mathbb{Z}^2 and E the set of rise vectors, fall vectors and long level vectors in the above half plane. Let $u_i = (x_0 - 2(i-1), 0)$ and $v_j^{(t)} = (x_0 + 2(j+t-1), 0)$ for $i, j = 1, 2, \dots, n$, $t = 0, 1, 2$ and a fixed integer x_0 . It is easy to see that the n -vertex $\mathbf{u} = (u_1, \dots, u_n)$ is D -compatible with the n -vertex $\mathbf{v}^{(t)} = (v_1^{(t)}, \dots, v_n^{(t)})$. We assign the weight of each edge as a rise vector, a fall vector and a long level vector whose origin is (x, y) and ends at $(x+1, y+1)$, $(x+1, y-1)$ and $(x+2, y)$ has weight $q^{x-y-x_0+2(n-1)}$, 1 and q^{x-y-x_0+2n-1} , respectively, which is visualized in Figure 11. Then, by applying Lemma 2.3, we can obtain

$$(5.16) \quad \text{GF} \left[\mathcal{P}_0 \left(\mathbf{u}, \mathbf{v}^{(t)} \right) \right] = \det \widehat{S}_n^{(t)}(q).$$

This is important to prove the following.

(i) Assume $t = 0$ and let \mathbf{u} and \mathbf{v} be as above. Put $\tilde{u}_i = (x_0 - 2i + 3, 1)$ and $\tilde{v}_j^{(1)} = (x_0 + 2j - 3, 1)$ for $i, j = 2, \dots, n$, and let $\tilde{\mathbf{u}} = (\tilde{u}_2, \dots, \tilde{u}_n)$ and $\tilde{\mathbf{v}}^{(1)} = (\tilde{v}_2^{(1)}, \dots, \tilde{v}_n^{(1)})$. Then each n -path $\mathbf{P} = (P_1, P_2, \dots, P_n)$ from \mathbf{u} to $\mathbf{v}^{(0)}$ corresponds to an $(n-1)$ -path $\tilde{\mathbf{P}} = (\tilde{P}_2, \dots, \tilde{P}_n)$ from $\tilde{\mathbf{u}}$ to $\tilde{\mathbf{v}}^{(1)}$ by regarding $\tilde{\mathbf{P}}$ as the subpath of \mathbf{P} . In fact, note that P_1 is always the path composed of a single vertex $u_1 = v_1$, each P_i always starts from the rise vector $u_i \rightarrow \tilde{u}_i$ and ends at the fall vector $\tilde{v}_i^{(1)} \rightarrow v_i^{(0)}$ for $i = 2, \dots, n$. Hence this gives a bijection, and the product of the weight of the rise vectors $u_i \rightarrow \tilde{u}_i$ and the fall vectors $\tilde{v}_i^{(1)} \rightarrow v_i^{(0)}$ for $i = 2, \dots, n$ is $q^{(n-1)(n-2)}$. This proves (5.13).

(ii) Assume $t = 1$ and let \mathbf{u} and \mathbf{v} be as above, i.e., $u_i = (x_0 - 2(i-1), 0)$ and $v_j^{(1)} = (x_0 + 2j, 0)$ for $1 \leq i, j \leq n$ (see Figure 12). Put $\tilde{u}_i = (x_0 - 2i + 3, 1)$ ($2 \leq i \leq n$) and $\tilde{v}_j^{(2)} = (x_0 + 2j - 1, 1)$ ($2 \leq j \leq n$), and let $\tilde{\mathbf{u}} = (\tilde{u}_2, \dots, \tilde{u}_n)$ and $\tilde{\mathbf{v}}^{(2)} = (\tilde{v}_2^{(2)}, \dots, \tilde{v}_n^{(2)})$. Further, put $\check{u}_i = (x_0 - 2i + 4, 2)$ ($2 \leq i \leq n$) and $\check{v}_j^{(1)} = (x_0 + 2j - 2, 2)$ ($2 \leq j \leq n$), and let $\check{\mathbf{u}} = (\check{u}_2, \dots, \check{u}_n)$ and $\check{\mathbf{v}}^{(1)} = (\check{v}_2^{(1)}, \dots, \check{v}_n^{(1)})$. Let $\mathbf{P} = (P_1, P_2, \dots, P_n)$ be any non-intersecting n -paths from \mathbf{u} to $\mathbf{v}^{(1)}$. Then, it is easy to see that \mathbf{P} must satisfy one of the following two conditions:

- (1) P_1 is the long level vector whose origin is u_1 and ends at $v_1^{(1)}$, and P_i goes through the vertices \tilde{u}_i and $\tilde{v}_i^{(2)}$ for $i = 2, 3, \dots, n$.
- (2) P_1 is a path which goes through only three vertices $u_1, u_1 + (1, 1) (= v_1^{(1)} - (1, -1))$ and $v_1^{(1)}$, and P_i goes through the vertices $\tilde{u}_i, \check{u}_i, \check{v}_i^{(1)}$ and $\tilde{v}_i^{(2)}$ for $i = 2, 3, \dots, n$.

By a similar argument as in the proof of (i), we can deduce that

$$\text{GF} \left[\mathcal{P}_0(\mathbf{u}_n, \mathbf{v}_n^{(1)}) \right] = q^{n^2 - n + 1} \text{GF} \left[\mathcal{P}_0(\mathbf{u}_{n-1}, \mathbf{v}_{n-1}^{(2)}) \right] + q^{2(n-1)^2} \text{GF} \left[\mathcal{P}_0(\mathbf{u}_{n-1}, \mathbf{v}_{n-1}^{(1)}) \right]$$

holds. By the equality (5.16), we obtain the identity (5.14).

(iii) This identity can be proven by applying the Desnanot-Jacobi adjoint matrix theorem (4.8) to $\tilde{S}_n^{(t)}(q)$. \square

Proof of Theorem 5.3. (i) The first equality of (5.9) is easily obtained from (5.11) and (5.13). By applying the equalities (5.11) and (5.12) to (5.14) and (5.15), we have

$$(5.17) \quad \det S_n^{(1)}(q) = q \det S_{n-1}^{(2)}(q) + \det S_{n-1}^{(1)}(q)$$

for $n \geq 2$, and we have

$$(5.18) \quad \det S_n^{(0)}(q) \det S_{n-2}^{(2)}(q) = \det S_{n-1}^{(0)}(q) \det S_{n-1}^{(2)}(q) - q^{2(n-1)} \{\det S_{n-1}^{(1)}(q)\}^2$$

for $n \geq 3$. By the equalities (5.17) and (5.18), for $n \geq 3$, the following identity holds:

$$(5.19) \quad \begin{aligned} & \det S_n^{(0)}(q) (\det S_{n-1}^{(1)}(q) - \det S_{n-2}^{(1)}(q)) \\ &= \det S_{n-1}^{(0)}(q) (\det S_n^{(1)}(q) - \det S_{n-1}^{(1)}(q)) - q^{2n-1} \{\det S_{n-1}^{(1)}(q)\}^2. \end{aligned}$$

Moreover, by applying the first equality of (5.9) to (5.19) and replacing n with $n-1$, we obtain

$$(5.20) \quad (1 + q^{2n-3}) \{\det S_{n-1}^{(0)}(q)\}^2 = \det S_{n-2}^{(0)}(q) \det S_n^{(0)}(q)$$

for $n \geq 4$. We prove the second equality of (5.9) by induction on n . If $n = 1, 2, 3$, then it is easily obtained by direct computations. Assume that (5.9) holds up to $n-1$. Then, by (5.20) and induction hypothesis, we can obtain the second equality (5.9).

(ii) It follows from our result of (i) and the equality (5.17) that (5.10) holds. \square

By applying the Desnanot-Jacobi adjoint matrix theorem (4.8) to $S_{n+1}^{(1)}(1)$, then we have

$$(5.21) \quad \det S_{n+1}^{(1)}(1) \det S_{n-1}^{(3)}(1) = \det S_n^{(1)}(1) \det S_n^{(3)}(1) - \{\det S_n^{(2)}(1)\}^2$$

for $n \geq 2$. Therefore the following identity is easily obtained by induction on n and the formula (5.21):

Remark. For positive integer n , we have

$$(5.22) \quad \det S_n^{(3)}(1) = 2^{\binom{n+3}{2}} - (2n+3)2^{\binom{n+2}{2}} - 2^{\binom{n+1}{2}}.$$

Note that there is a relation between domino tilings of the Aztec diamonds and Schröder paths (see [8, 15]). It might be an interesting problem to consider what this weight means.

§ 5.4. Delannoy numbers

The *Delannoy numbers* $D(a, b)$ are the number of lattice paths from $(0, 0)$ to (b, a) in which only east $(1, 0)$, north $(0, 1)$, and northeast $(1, 1)$ steps are allowed. They are given by the recurrence relation

$$(5.23) \quad \begin{aligned} D(a, 0) &= D(0, b) = 1, \\ D(a, b) &= D(a-1, b) + D(a-1, b-1) + D(a, b-1). \end{aligned}$$

The first few terms of $D(n, n)$ ($n = 0, 1, 2, \dots$) are given by 1, 3, 13, 63, 321, \dots . By a similar argument we can derive the following result. We may give a proof in another occasion.

Proposition 5.6. *Let n be a positive integers. Then the following identities would hold:*

$$(5.24) \quad \det (D(i+j, i+j))_{0 \leq i, j \leq n-1} = 2^{\binom{n+1}{2}-1},$$

$$(5.25) \quad \det (D(i+j+1, i+j+1))_{0 \leq i, j \leq n-1} = 2^{\binom{n+2}{2}-2} + 2^{\binom{n+1}{2}-1},$$

$$(5.26) \quad \det (D(i+j+2, i+j+2))_{0 \leq i, j \leq n-1} = 2^{\binom{n+3}{2}-3} + (2n+1)2^{\binom{n+2}{2}-2} - 2^{\binom{n+1}{2}-1}.$$

§ 6. Concluding remarks

Since a hyperpfaffian version of (1.3) is obtained in [13], we believe it will be interesting problem to consider a hyperpfaffian version of Theorem 1.1 and Theorem 1.3. We shall argue on it in another chance. The authors also would like to express their gratitude to the anonymous referee for his (her) constructive comments.

References

- [1] M. Aigner, "Motzkin Numbers", *Europ. J. Combinatorics* **19** (1998), 663 – 675.
- [2] G. Andrews and R. Askey, "Enumeration of partitions: The role of Eulerian series and q -orthogonal polynomials", in *Higher Combinatorics* (M. Aigner, ed.), Reidel Publications, Boston, 1977, pp. 3 – 26.

- [3] J. L. Burchnall and T. W. Chaundy, “The hypergeometric identities of Cayley, Orr, and Bailey”, *Proc. London Math. Soc.* (2) **50**, (1948). 56–74.
- [4] R. A. Brualdi and S. Kirkland, “Aztec diamonds and digraphs, and Hankel determinants of Schröder numbers”, *J. Combin. Theory Ser. B* **94**, (2005). 334–351.
- [5] A. Benjamin, N. Cameron, J. Quinn and C. Yerger, “Catalan Determinants — A Combinatorial Approach”, preprint.
- [6] A. Cvetković, P. Rajković and M. Ivković, “Catalan numbers, the Hankel transform, and Fibonacci numbers”, *J. Integer Seq.* **5** (2002), Article 02.1.3.
- [7] P. Flajolet, “Combinatorial aspects of continued fractions”, *Discrete Math.* **32** (1980) 125 – 161.
- [8] S. Eu and T. Fu, “A Simple Proof of the Aztec Diamond Theorem”, *Electron. J. Combin.* **12** (2005), #R18.
- [9] G. Gasper and M. Rahman, *Basic Hypergeometric Series*, Cambridge University Press, Cambridge, Second edition, 2004.
- [10] I. Gessel and G. Viennot, *Determinants, Paths, and Plane Partitions*, preprint (1989).
- [11] C. Krattenthaler, “Advanced Determinant Calculus”, *Sem. Lothar. Combin.* **42** (“The Andrews Festschrift”) (1999), Article B42q.
- [12] C. Krattenthaler, “Determinants of (generalized) Catalan numbers”, [arXiv:math.CO/0709.3044](https://arxiv.org/abs/math/0709.3044).
- [13] J-G. Luque and J-Y. Thibon, “Hankel hyperdeterminants and Selberg integrals [arXiv:math-ph/0211044](https://arxiv.org/abs/math-ph/0211044).”
- [14] M.E. Mays and J. Wojciechowski, “A determinant property of Catalan numbers”, *Discrete Math.* **211** (2000), 125 – 133.
- [15] R. Stanley, *Enumerative Combinatorics*, Volume 1, 2, Cambridge University Press, 1997.
- [16] R. Sulanke and G. Xin, “Hankel Determinants for Some Common Lattice Paths” 18th International Conference on Formal Power Series and Algebraic Combinatorics, San Diego, California 2006.
- [17] U. Tamm, “Some Aspects of Hankel Matrices in Coding Theory and Combinatorics”, *Electron. J. Combin.* **8** (2001), #A1.
- [18] X. Viennot, *Une théorie combinatoire des polynômes orthogonaux*, in Lecture Notes at UQAM, 1984.
- [19] X. Viennot, “A combinatorial theory for general orthogonal polynomials with extensions and applications”, in *Orthogonal Polynomials and Applications*, Lecture Notes in Mathematics, Vol. 1171, Springer, Berlin, 1985, 139 – 157.




Figure 6. $t = 1$ and $n = 4$




Figure 7. A Dyck Path of weight $\lambda_1 \lambda_3^2$




Figure 8. $\text{GF} [\mathcal{D}_{m,n}] = \text{GF} [\mathcal{D}_{m-1,n-1}] + \lambda_{n+1} \text{GF} [\mathcal{D}_{m-1,n+1}]$




Figure 9. Dyck Paths in $\mathcal{D}_{4,0}$




Figure 10. Weight of Schröder paths in $\mathcal{S}_{4,0}$




Figure 11. Weight of each edge




Figure 12. $t = 1$ case