

LINEAR RECURRENCES AND CHEBYSHEV POLYNOMIALS

Sergey Kitaev

Matematik, Chalmers tekniska högskola och Göteborgs universitet, S-412 96 Göteborg, Sweden
e-mail: kitaev@math.chalmers.se

Toufik Mansour

Matematik, Chalmers tekniska högskola och Göteborgs universitet, S-412 96 Göteborg, Sweden
e-mail: toufik@math.chalmers.se

(Submitted December 2002-Final Revision May 2003)

1. INTRODUCTION AND THE MAIN RESULT

As usual, Fibonacci polynomials $F_n(x)$, Lucas polynomials $L_n(x)$, and Pell polynomials $P_n(x)$ are defined by the second-order linear recurrence

$$t_{n+2} = at_{n+1} + bt_n, \tag{1}$$

with given a, b, t_0, t_1 and $n \geq 0$. This sequence was introduced by Horadam [3] in 1965, and it generalizes many sequences (see [1, 4]). Examples of such sequences are Fibonacci polynomials sequence $(F_n(x))_{n \geq 0}$, Lucas polynomials sequence $(L_n(x))_{n \geq 0}$, and Pell polynomials sequence $(P_n(x))_{n \geq 0}$, when one has $a = x, b = t_1 = 1, t_0 = 0$; $a = t_1 = x, b = 1, t_0 = 2$; and $a = 2x, b = t_1 = 1, t_0 = 0$; respectively.

Chebyshev polynomials of the second kind (in this paper just Chebyshev polynomials) are defined by

$$U_n(\cos \theta) = \frac{\sin(n+1)\theta}{\sin \theta}$$

for $n \geq 0$. Evidently, $U_n(x)$ is a polynomial of degree n in x with integer coefficients. For example, $U_0(x) = 1, U_1(x) = 2x, U_2(x) = 4x^2 - 1$, and in general (see Recurrence 1 for $a = 2x, b = -1, t_0 = 1$, and $t_1 = 2x, U_{n+2}(x) = 2xU_{n+1}(x) - U_n(x)$). Chebyshev polynomials were invented for the needs of approximation theory, but are also widely used in various other branches of mathematics, including algebra, combinatorics, and number theory (see [5]).

Lemma 1.1: *Let $(t_n)_{n \geq 0}$ be any sequence that satisfies $t_{n+2} = 2x \cdot t_{n+1} - t_n$ with given t_0, t_1 , and $n \geq 0$. Then for all $n \geq 0$,*

$$t_n = t_1 \cdot U_{n-1}(x) - t_0 \cdot U_{n-2}(x),$$

where U_m is the m^{th} Chebyshev polynomial of the second kind.

Proof: A proof is straightforward using the relation $U_{n+2}(x) = 2xU_{n+1}(x) - U_n(x)$ and induction on n . \square

Let A be a tile of size 1×1 and B be a tile of size 1×2 . We denote by \mathcal{L}_n the set of all *tilings* of a $1 \times n$ rectangle with tiles A and B . An element of \mathcal{L}_n can be written as a sequence of the letters A and B . For example, $\mathcal{L}_1 = \{A\}$, $\mathcal{L}_2 = \{AA, B\}$, and $\mathcal{L}_3 = \{AAA, AB, BA\}$. We denoted by $|\alpha|$ the number of tiles A and B in α . For example, $|AAA| = 3$ and $|AB| = 2$.

Proposition 1.2: *The number of tilings of a $1 \times n$ rectangle with tiles A and B is the Fibonacci number F_{n+1} , that is, $|\mathcal{L}_n| = F_{n+1}$.*

Proof: The result is immediate for $n \leq 1$, so it is sufficient to show that the number of such tilings satisfies the recurrence $F_m = F_{m-1} + F_{m-2}$. To do this, we observe that there is a one-to-one correspondence between the tilings of a $1 \times (n - i)$ rectangle and the tilings of a $1 \times n$ rectangle in which the rightmost tile has length i , where $i = 1, 2$. Therefore, if we count tilings of a $1 \times n$ rectangle according to the length of the rightmost tile, we find the number of such tilings satisfies the recurrence $F_m = F_{m-1} + F_{m-2}$, as desired. \square

Let α be any element of \mathcal{L}_n , we define β by $\beta_i = 1$ if $\alpha_i = A$; otherwise $\beta_i = 2$, and we write $\beta = \chi(\alpha)$. For example, $\chi(AAABAB) = 111212$.

Now, let us fix an integer s and a natural number q such that $q \geq 1$. Let $a_0, a_1, \dots, a_{q-1}, b_0, b_1, \dots, b_{q-1}$ be $2q$ constants and $\mathbf{a} = (a_0, a_1, \dots, a_{q-1})$, $\mathbf{b} = (b_0, b_1, \dots, b_{q-1})$. For any $\alpha \in \mathcal{L}_n$, we define $v(n; s) = v_{\mathbf{a}, \mathbf{b}}(n; \alpha, q, s) = \prod_{i=1}^{|\alpha|} k(\beta_i)$ where

$$k(\beta_i) = \begin{cases} a_{(s+\beta_1+\dots+\beta_i) \bmod q}, & \text{if } \beta_i = 1, \\ b_{(s+\beta_1+\dots+\beta_i) \bmod q}, & \text{if } \beta_i = 2, \end{cases}$$

and $\beta = \chi(\alpha)$. For example, if $q = 3, a_n = n$ and $b_n = 1$ for $n = 0, 1, 2, s = 0$, and $\alpha = AABAB$, then we have that

$$v_{\mathbf{a}, \mathbf{b}}(n; \alpha, q, s) = a_{1 \bmod 3} a_{2 \bmod 3} a_{4 \bmod 3} a_{5 \bmod 3} b_{7 \bmod 3} = a_1 a_2 b_1 a_2 b_1 = a_1 a_2^2 = 4.$$

We will be interested in the sum of all $v_{\mathbf{a}, \mathbf{b}}(n; \alpha, q, s)$ over all $\alpha \in \mathcal{L}_n$, which is denoted by $V(n; s) = V_{\mathbf{a}, \mathbf{b}}(n; q, s)$, that is, $V(n; s) = \sum_{\alpha \in \mathcal{L}_n} v_{\mathbf{a}, \mathbf{b}}(n; \alpha, q, s)$. For example, $V(1; s) = a_{(s+1) \bmod q}$ and $V(2; s) = a_{(s+1) \bmod q} a_{(s+2) \bmod q} + b_{(s+2) \bmod q}$. We extend the definition of $V(n; s)$ as $V(0; s) = 1$ and $V(n; s) = 0$ for $n < 0$. We remark that $V(n; q, s)$ can be given by a combinatorial interpretation as follows: $V(n; q, s)$ counts the number of ways to tile a boards of length n , with cells numbers $s + 1$ through $s + n$, using colored tiles of size 1×1 and tiles of size 1×2 . For a tile of size 1×1 on cell i , we have $a_{i \bmod q}$ color choices; for a tile of size 1×2 on cells $i - 1$ and i , we have $b_{i \bmod q}$ choices. The main result of this paper can be formulated as follows.

Theorem 1.3: *Let $(x_n)_{n \geq 0}$ be any sequence $(x_n = x_{n; q}(\mathbf{a}, \mathbf{b}))$ that satisfies*

$$x_{qn+d} = a_d \cdot x_{qn+d-1} + b_d \cdot x_{qn+d-2}, \tag{2}$$

for all $n \geq 1, 0 \leq d \leq q - 1$, with given x_0, x_1, \dots, x_{q-1} . Then for $n \geq 1, x_{qn+d}$ is given by

$$\sqrt{-J_{q;d}}^{n-2} \left(x_{q+d} \sqrt{-J_{q;d}} U_{n-1}(w_{q;d}) + (x_{2q+d} - I_{q;d} x_{q+d}) \cdot U_{n-2}(w_{q;d}) \right),$$

for all $n \geq 1$, where U_m is the m^{th} Chebyshev polynomial,

$$x_{q+d} = V(d + 1; -1)x_{q-1} + b_0 V(d; 0)x_{q-2}$$

$$x_{2q+d} = V(q + d + 1; -1)x_{q-1} + b_0 V(q + d; 0)x_{q-2},$$

and

$$w_{q;d} = \frac{I_{q;d}}{2\sqrt{-J_{q;d}}},$$

$$\begin{aligned} I_{q;d} &= b_{(d+1) \bmod q} \cdot V(q-2; d+1) + V(q; d), \\ J_{q;d} &= b_{(d+1) \bmod q} \cdot (V(q-1; d+1)V(q-1; d) - V(q; d)V(q-2; d+1)). \end{aligned} \quad (3)$$

The paper is organized as follows. In Section 2 we give a proof of Theorem 1.3, and in Section 3 we give some applications for Theorem 1.3.

2. PROOFS

Throughout this section, we assume that q is a natural number ($q \geq 1$) and s is an integer. Also, let $a_0, a_1, \dots, a_{q-1}, b_0, b_1, \dots, b_{q-1}$ be $2q$ constants and $\mathbf{a} = (a_0, a_1, \dots, a_{q-1})$, $\mathbf{b} = (b_0, b_1, \dots, b_{q-1})$. We start from the following lemma.

Lemma 2.1: *Let ℓ be an integer such that $\ell \geq s + 2$. Then*

$$V(\ell - s; s) = a_{\ell \bmod q} \cdot V(\ell - s - 1; s) + b_{\ell \bmod q} \cdot V(\ell - s - 2; s).$$

Proof: To verify this lemma, we observe that there is a one-to-one correspondence between the tilings of a $1 \times (\ell - s - i)$ rectangle and the tilings of a $1 \times (\ell - s)$ rectangle in which the rightmost tile has length i , where $i = 1, 2$. Hence $V(\ell - s; s) = a_{\ell \bmod q} \cdot V(\ell - s - 1; s) + b_{\ell \bmod q} \cdot V(\ell - s - 2, s)$, where the first term corresponds to the case $i = 1$ and the second one to the case $i = 2$. \square

Now, let us apply this lemma to find x_{qn+d+m} in terms of x_{qn+d} and x_{qn+d-1} .

Proposition 2.2: *Let $q - 1 \geq d \geq 0$ and $n \geq 1$. Then for all $m \geq 0$,*

$$x_{qn+d+m} = V(m; d) \cdot x_{qn+d} + b_{(d+1) \bmod q} \cdot V(m-1; d+1) \cdot x_{qn+d-1}.$$

Proof: Let us prove this proposition by induction on m . Since

$$x_{qn+d+0} = 1 \cdot x_{qn+d+0} + b_{(d+1) \bmod q} \cdot 0 \cdot x_{qn+d-1},$$

$V(0; d) = 1$ and $V(m; d) = 0$ for $m < 0$, we have that the proposition holds for $m = 0$. By Recurrence 2 we get

$$\begin{aligned} x_{qn+d+1} &= a_{(d+1) \bmod q} \cdot x_{qn+d} + b_{(d+1) \bmod q} \cdot x_{qn+d-1} \\ &= V(1; d) \cdot x_{qn+d} + b_{(d+1) \bmod q} \cdot V(0; d+1) \cdot x_{qn+d-1}, \end{aligned}$$

therefore the proposition holds for $m = 1$. Now, we assume that the proposition holds for $0, 1, \dots, m-1$, and prove that it holds for m . By induction hypothesis we have

$$x_{qn+d+m-2} = V(m-2; d) \cdot x_{qn+d} + b_{(d+1) \bmod q} \cdot V(m-3; d+1) \cdot x_{qn+d-1},$$

and

$$x_{qn+d+m-1} = V(m-1; d) \cdot x_{qn+d} + b_{(d+1) \bmod q} \cdot V(m-2; d+1) \cdot x_{qn+d-1},$$

hence, by Equation 2 we get

$$\begin{aligned} x_{qn+d+m} &= a_{(d+m) \bmod q} \cdot x_{qn+d+m-1} + b_{(d+m) \bmod q} \cdot x_{qn+m+d-2} \\ &= (a_{(d+m) \bmod q} \cdot V(m-1; d) + b_{(d+m) \bmod q} \cdot V(m-2; d)) x_{qn+d} \\ &\quad + b_{(d+1) \bmod q} (a_{(d+m) \bmod q} \cdot V(m-2; d+1) + b_{(d+m) \bmod q} \cdot V(m-3; d+1)) x_{qn+d-1}. \end{aligned}$$

Using Lemma 2.1 for $\ell = m + d, s = d$ and for $\ell = m + d, s = d + 1$, we get the desired result. \square

Now we introduce a recurrence relation that plays the crucial role in the proof of the Main Theorem.

Proposition 2.3: *Let $q - 1 \geq d \geq 0$. Then for all $n \geq 2$,*

$$\begin{aligned} x_{q(n+1)+d} &= (b_{(d+1) \bmod q} \cdot V(q-2; d+1) + V(q; d)) x_{qn+d} \\ &\quad + b_{(d+1) \bmod q} \cdot (V(q-1; d+1)V(q-1; d) - V(q; d)V(q-2; d+1)) x_{q(n-1)+d}. \end{aligned}$$

Proof: Using Proposition 2.2 for $m = q - 1$ we get

$$x_{q(n+1)+d-1} - b_{(d+1) \bmod q} \cdot V(q-2; d+1) \cdot x_{qn+d-1} = V(q-1; d) \cdot x_{qn+d}, \quad (4)$$

and for $m = q$ we have

$$x_{q(n+1)+d} = V(q; d) \cdot x_{qn+d} + b_{(d+1) \bmod q} \cdot V(q-1; d+1) \cdot x_{qn+d-1}. \quad (5)$$

Hence, Equation 4 yields

$$\begin{aligned} x_{q(n+1)+d} - b_{(d+1) \bmod q} \cdot V(q-2; d+1) \cdot x_{qn+d} &= \\ &= V(q; d) (x_{qn+d} - b_{(d+1) \bmod q} \cdot V(q-2; d+1) \cdot x_{qn+d}) \\ &\quad + b_{(q+1) \bmod q} \cdot V(q-1; d+1) (x_{qn+d-1} - b_{(d+1) \bmod q} \cdot V(q-2; d+1) \cdot x_{q(n-1)+d-1}), \end{aligned}$$

and by using Equation 4 we get the desired result. \square

Proof of Theorem 1.3: Recall the definitions in 3. Now we are ready to prove the main result of this paper. Using Proposition 2.3 we have for $n \geq 2$,

$$x_{q(n+1)+d} = I_{q;d} \cdot x_{qn+d} + J_{q;d} \cdot x_{q(n-1)+d}.$$

If we define $t_n = x_{qn+d}$ for $n \geq 1$, then we get

$$t_{n+1} = I_{q;d} \cdot t_n + J_{q;d} \cdot t_{n-1},$$

therefore, by defining $(-J_{q;d})^{n/2} t'_n = t'_n$ we have for $n \geq 2$, $t'_{n+1} = 2w_{q;d} t'_n - t'_{n-1}$.

Let us find expressions for t'_0 and t'_1 . By the recurrence for t_n we can define t_0 as $t_2 = I_{q;d} t_1 + J_{q;d} t_0$, which means that $t'_0 = t_0 = \frac{1}{J_{q;d}} (x_{2q+d} - I_{q;d} x_{q+d})$. By definitions, $t'_1 = \frac{x_{q+d}}{\sqrt{-J_{q;d}}}$.

Using Proposition 2.2, we get $x_{q+d} = V(d+1; -1)x_{q-1} + b_0V(d; 0)x_{q-2}$ and $x_{2q+d} = V(q+d+1; -1)x_{q-1} + b_0V(q+d; 0)x_{q-2}$. Hence, using Lemma 1.1 we get the desired result.

3. APPLICATIONS

There is a connection between the sequences which are defined by Recurrence 2, and the sequences which are defined by Recurrence 1. Indeed, from Theorem 1.3 we get the following result.

Corollary 3.1: *For given x_0 and x_{-1} , and the recurrence $x_{n+2} = a_0x_{n+1} + b_0x_n$, an explicit solution for this recurrence is given by*

$$x_n = \sqrt{-b_0}^{n-2} \left[\sqrt{-b_0}(a_0x_0 + b_0x_{-1})U_{n-1} \left(\frac{a_0}{2\sqrt{-b_0}} \right) + b_0x_0U_{n-2} \left(\frac{a_0}{2\sqrt{-b_0}} \right) \right],$$

where U_m is the m^{th} Chebyshev polynomial.

Proof: Using Theorem 1.3 for $q = 1$ with the parameters $d = 0, I_{1;0} = a_0, J_{1;0} = b_0, x_1 = a_0x_0 + b_0x_{-1}, x_2 = (a_0^2 + b_0)x_0 + a_0b_0x_{-1}$, and $w_{1;0} = \frac{a_0}{2\sqrt{-b_0}}$, we get the explicit solution for the recurrence $x_{n+2} = a_0x_{n+1} + b_0x_n$, as requested. \square

The first interesting case is $q = 2$. Then Recurrence 2 gives

$$\begin{cases} x_{2n} &= a_0x_{2n-1} + b_0x_{2n-2} \\ x_{2n+1} &= a_1x_{2n} + b_1x_{2n-1}, \end{cases} \quad (6)$$

with given x_0 and x_1 . In this case we have two possibilities: either $d = 0$ or $d = 1$. Let $d = 0$, so the parameters of the problem are given by $I_{2;0} = a_0a_1 + b_0 + b_1, J_{2;0} = -b_0b_1, w_{2;0} = \frac{a_0a_1 + b_0 + b_1}{2\sqrt{b_0b_1}}, x_2 = a_0x_1 + b_0x_0$, and $x_4 = (a_0^2a_1 + a_0b_1 + a_0b_0)x_1 + (a_0b_0a_1 + b_0^2)x_0$. Hence, Theorem 1.3 gives the following result.

Corollary 3.2: *The solution x_{2n} for Recurrence 6 is given by*

$$\sqrt{b_0b_1}^{n-2} \left[\sqrt{b_0b_1}(a_0x_1 + b_0x_0)U_{n-1} \left(\frac{a_0a_1 + b_0 + b_1}{2\sqrt{b_0b_1}} \right) - b_0b_1x_0U_{n-2} \left(\frac{a_0a_1 + b_0 + b_1}{2\sqrt{b_0b_1}} \right) \right],$$

where U_m is the m^{th} Chebyshev polynomial.

Example 3.3: *If $x_0 = 0, x_1 = 1, a_0 = x, a_1 = xy$, and $b_0 = b_1 = 1$, then the explicit expression to x_{2n} for the Recurrence 6 is given by $xU_{n-1}(1 + \frac{1}{2}x^2y)$. Hence, by the definition it is easy to see that in the case $y = 1$, we have that the Fibonacci polynomial $F_{2n}(x)$ is given by $xU_{n-1}(1 + \frac{1}{2}x^2)$.*

If $x_0 = 2, x_1 = 1, a_0 = x, a_1 = xy$, and $b_0 = b_1 = 1$, then an explicit expression to x_{2n} for the Recurrence 6 is given by $(x+2)U_{n-1}(1 + \frac{1}{2}x^2y) - 2U_{n-2}(1 + \frac{1}{2}x^2y)$. Hence, in the case $y = 1$ we have that the Lucas polynomial $L_{2n}(x)$ is given by $(x+2)U_{n-1}(1 + \frac{1}{2}x^2) - 2U_{n-2}(1 + \frac{1}{2}x^2)$.

If $x_0 = 0, x_1 = 1, a_0 = 2x, a_1 = yx$, and $b_0 = b_1 = 1$, then an explicit expression to x_{2n} for the Recurrence 6 is given by $2xU_{n-1}(1 + x^2y)$. Hence, in the case $y = 2$ we have that the Pell polynomial $P_{2n}(x)$ is given by $2xU_{n-1}(1 + 2x^2)$.

Another example for Theorem 1.3 is when $q = 3$ and $d = 0$. In this case the parameters of the problem are given by $I_{3;0} = a_0a_1a_2 + b_0a_1 + b_1a_2 + a_0b_2$, $J_{3;0} = b_0b_1b_2$, $x_3 = a_0x_2 + b_0x_1$, and $x_6 = I_{3;0}x_3 = b_0b_1(x_2 - a_2x_1)$. Therefore, we get the following result.

Corollary 3.4: *The solution x_{2n} for Recurrence 2, when $q = 3$, is given by*

$$\sqrt{-b_0b_1b_2}^{n-2} \left(\sqrt{-b_0b_1b_2}(a_0x_2 + b_0x_1)U_{n-1}(w) + b_0b_1(x_2 - a_2x_1)U_{n-2}(w) \right),$$

for all $n \geq 1$, where $w = \frac{a_0a_1a_2 + a_0b_2 + b_0a_1 + b_1a_2}{2\sqrt{-b_0b_1b_2}}$, and U_m is the m^{th} Chebyshev polynomial.

For example, if we are interested in solving the recurrence

$$\begin{cases} x_{3n} &= x_{3n-1} + x_{3n-2} \\ x_{3n+1} &= x_{3n} + x_{3n-1} \\ x_{3n+2} &= yx_{3n+1} + x_{3n}, \end{cases}$$

with $x_0 = 0$ and $x_1 = x_2 = 1$, then by the above corollary we get that the solution x_{3n} for this recurrence is given by

$$2i^{n-1}U_{n-1}(-i(1+y)) + i^{n-2}(1-y)U_{n-2}(-i(1+y)),$$

where $i^2 = -1$. In particular, if $y = 1$ then we have that the $(3n)^{\text{th}}$ Fibonacci number, F_{3n} , is given by $2i^{n-1}U_{n-1}(-2i)$.

ACKNOWLEDGMENT

The authors would like to thank the anonymous referee for careful reading of the manuscript.

The second authors research financed by EC's IHRP Programme, within the Research training Network "Algebraic Combinatorics in Europe", grant HPRN-CT-2001-00272.

REFERENCES

- [1] G.H. Hardy and E.M. Wright. *An Introduction to the Theory of Numbers*. 4th Edition. London, Oxford Univresity Press, 1962.
- [2] P. Haukkanen. "A Note on Horadam's Sequence." *The Fibonacci Quarterly* **40.4** (2002): 358-361.
- [3] A.F. Horadam. "Generalization of a Result of Morgado." *Portugaliae Math.* **44** (1987): 131-136.
- [4] A.F. Horadam and J.M. Mahon. "Pell and Pell-Lucas Polynomials." *The Fibonacci Quarterly* **23.1** (1985): 7-20.
- [5] Th. Rivlin. *Chebyshev Polynomials. From Approximation Theory to Algebra and Number Theory*. John Wiley, New York, 1990.

AMS Classification Numbers: 11B83, 42C05, 11K31

