

ON THE FIBONACCI NUMBERS WHOSE SUBSCRIPT IS A POWER

Piero Filippini

Fondazione Ugo Bordoni, Via B. Castiglione 59, I-00142 Rome, Italy

(Submitted October 1994)

1. AIM OF THE PAPER

The well-known identity (e.g., see [1], p. 127)

$$F_{2^n} = \prod_{i=1}^{n-1} L_{2^i} \tag{1.1}$$

led us to investigate the Fibonacci numbers of the form F_{k^n} with k and n positive integers. The principal aim of this note is to present some new identities involving F_{k^n} , some of which generalize (1.1). This is done in Sections 2 and 3. In Section 4, a first-order recurrence relation for F_{k^n} is established which involves certain combinatorial quantities whose properties will be investigated in a future paper. A glimpse of analogous results concerning the Lucas numbers L_{k^n} is caught in Section 5.

The formulas established in this note encompass the trivial case $n = 1$ under the usual assumptions

$$\prod_{i=a}^b f(i) = 1 \text{ if } b < a \tag{1.2}$$

and

$$\sum_{i=a}^b f(i) = 0 \text{ if } b < a. \tag{1.3}$$

2. MAIN RESULT

Proposition 1: If k is even and $n \geq 1$, then

$$F_{k^n} = F_k \prod_{i=1}^{n-1} \left[\sum_{j=1}^{k/2} L_{(2^{j-1})k^i} \right]. \tag{2.1}$$

We can immediately observe that, if $k = 2$, identity (2.1) reduces to (1.1).

Proof of Proposition 1: Write

$$F_{k^n} = F_k \cdot \frac{F_{k^2}}{F_k} \cdot \frac{F_{k^3}}{F_{k^2}} \cdot \dots \cdot \frac{F_{k^n}}{F_{k^{n-1}}} = F_k \prod_{i=1}^{n-1} \frac{F_{k^{i+1}}}{F_{k^i}} \tag{2.2}$$

whence, following the notation used in [2] [namely, $R_s(t) = F_{st} / F_t$], we can rewrite (2.2) as

$$F_{k^n} = F_k \prod_{i=1}^{n-1} R_k(k^i). \tag{2.3}$$

Using (2.3) along with (2.1) of [2] yields

$$F_{k^n} = F_k \prod_{i=1}^{n-1} \left\{ \sum_{j=1}^{k/2} [F_{k'(2j-1)-1} + F_{k'(2j-1)+1}] \right\}. \tag{2.4}$$

The right-hand side of (2.4) clearly equals that of (2.1). Q.E.D.

Proposition 2: If k is odd and $n \geq 1$, then

$$F_{k^n} = (-1)^{(n-1)(k-1)/2} F_k \prod_{i=1}^{n-1} \left[1 + \sum_{j=1}^{(k-1)/2} (-1)^j L_{2jk^i} \right]. \tag{2.5}$$

Observe that, if $k = 3$, identity (2.5) reduces beautifully to

$$F_{3^n} = 2 \prod_{i=1}^{n-1} (L_{2 \cdot 3^i} - 1). \tag{2.6}$$

In order to prove Proposition 2, we need the identity

$$\sum_{i=1}^r (-1)^i L_{ai} = \frac{(-1)^{a+r} L_{ar} + (-1)^r L_{a(r+1)} - L_a - 2(-1)^a}{L_a + 1 + (-1)^a}, \tag{2.7}$$

which can readily be proved by using the Binet form for Lucas numbers and the geometric series formula. We also need the identity

$$L_{a+b} - (-1)^b L_{a-b} = 5F_a F_b, \tag{2.8}$$

which can be obtained from identities $I_{21} - I_{24}$ of [3].

Proof of Proposition 2: The proof has to be split into two cases according to the residue of k modulo 4.

Case 1. $k \equiv 1 \pmod{4}$ [i.e., $(k-1)/2$ is even]

By (2.2), we can write

$$\begin{aligned} F_{k^n} &= F_k \prod_{i=1}^{n-1} \frac{F_{k^{i+1}}}{F_{k^i}} = F_k \prod_{i=1}^{n-1} \left[1 + \frac{5F_{k^{i+1}}F_{k^i}}{5F_{k^i}^2} - 1 \right] \\ &= F_k \prod_{i=1}^{n-1} \left[1 + \frac{L_{k^{i+1}-k^i} + L_{k^{i+1}+k^i}}{L_{2k^i} + 2} - 1 \right] \quad (\text{by (2.8) and } I_{17} \text{ of [3]}) \\ &= F_k \prod_{i=1}^{n-1} \left[1 + \frac{L_{2k^i(k-1)/2} + L_{2k^i(k+1)/2} - L_{2k^i} - 2}{L_{2k^i} + 2} \right] \\ &= F_k \prod_{i=1}^{n-1} \left[1 + \sum_{h=1}^{(k-1)/2} (-1)^h L_{2hk^i} \right] \quad [\text{by (2.7)}]. \end{aligned}$$

Observe that, since $(k-1)/2$ is even by hypothesis, the above expression does not vary if we multiply it by $(-1)^{(n-1)(k-1)/2}$.

Case 2. $k \equiv 3 \pmod{4}$ [i.e., $(k-1)/2$ is odd]

Analogously, we can write

$$\begin{aligned} F_{k^n} &= (-1)^{n-1} F_k \prod_{i=1}^{n-1} \left[-\frac{5F_{k^{i+1}}F_{k^i}}{5F_{k^i}^2} - 1 + 1 \right] \\ &= (-1)^{n-1} F_k \prod_{i=1}^{n-1} \left[1 + \frac{-L_{2k^i(k-1)/2} - L_{2k^i(k+1)/2} - L_{2k^i} - 2}{L_{2k^i} + 2} \right] \\ &= (-1)^{n-1} F_k \prod_{i=1}^{n-1} \left[1 + \sum_{h=1}^{(k-1)/2} (-1)^h L_{2hk^i} \right]. \end{aligned}$$

Observe that, since $(k-1)/2$ is odd by hypothesis, the factor $(-1)^{n-1}$ in the above expression can be rewritten as $(-1)^{(n-1)(k-1)/2}$. Q.E.D.

3. RELATED RESULTS

Some results related to those established in Section 2 can be obtained readily. Observe that, if the exponent n in (2.1) is composite (say, $n = st$), then F_{k^n} (k even) can be expressed as

$$F_{k^n} = F_{k^{st}} = F_{k^s} \prod_{i=1}^{t-1} \left[\sum_{j=1}^{k^s/2} L_{(2j-1)k^{si}} \right] \quad (k \text{ even}), \tag{3.1}$$

where s and t can obviously be interchanged. For example, by (1.1) and (3.1), we have

$$F_{2^{2n}} = F_{4^n} = \prod_{i=1}^{2n-1} L_{2^i} = F_{2^n} \sum_{j=1}^{2^{n-1}} L_{(2j-1)2^n} = 3 \prod_{i=1}^{n-1} (L_{4^i} + L_{3 \cdot 4^i}). \tag{3.2}$$

For k odd, the analog of (3.1) can be obtained immediately.

An expression analogous to (2.1) can be established for F_{mk^n} when mk is even. If mk is odd, the corresponding expression is somewhat unattractive and its presentation is omitted.

Proposition 3: If $n \geq 1$,

$$F_{mk^n} = F_{mk} \prod_{i=1}^{n-1} \left[\sum_{j=1}^{k/2} L_{(2j-1)mk^i} \right] \quad (k \text{ even, } m \text{ arbitrary}) \tag{3.3}$$

and

$$F_{mk^n} = F_{mk} \prod_{i=1}^{n-1} \left[1 + \sum_{j=1}^{(k-1)/2} L_{2jmk^i} \right] \quad (k \text{ odd, } m \text{ even}). \tag{3.4}$$

Proof: Write

$$F_{mk^n} = F_{mk} \prod_{i=1}^{n-1} \frac{F_{mk^{i+1}}}{F_{mk^i}} = F_{mk} \prod_{i=1}^{n-1} R_k(mk^i)$$

and use (2.1) and (2.2) of [2]. Q.E.D.

If we let $m = k$ in (3.3), we see that an equivalent form for (2.1) is

$$F_{k^n} = F_{k^2} \prod_{i=1}^{n-2} \left[\sum_{j=1}^{k/2} L_{(2j-1)k^{i+1}} \right] \quad (k \text{ even}, n \geq 2). \quad (3.5)$$

Moreover, if we let $n = u + v$ ($u, v \geq 1$) and $m = k^u$ in (3.3), we get the relation

$$F_{k^n} = F_{k^{u+v}} = F_{k^u k^v} = F_{k^{u+1}} \prod_{i=1}^{v-1} \left[\sum_{j=1}^{k/2} L_{(2j-1)k^{u+i}} \right] \quad (k \text{ even}), \quad (3.6)$$

which generalizes (3.5).

4. A RECURRENCE RELATION FOR F_{k^n}

A problem [6] that appeared in this journal led us to discover the first-order nonlinear homogeneous recurrence relation

$$F_{3^{n+1}} = 5F_{3^n}^3 - 3F_{3^n} \quad (n \geq 0). \quad (4.1)$$

The aim of this section is to obtain an analogous relation valid for all positive subscripts k^{n+1} with k odd and n an arbitrary nonnegative integer.

Proposition 4: If k is a positive odd integer and n is a nonnegative integer, then

$$F_{k^{n+1}} = 5^{(k-1)/2} F_{k^n}^k - \sum_{i=0}^{(k-3)/2} 5^i C_{i,k} F_{k^n}^{2i+1}, \quad (4.2)$$

where the coefficients $C_{i,k}$ are given by

$$C_{i,k} = (-1)^{(k+1)/2+i} \binom{(k+1)/2+i}{2i+1} \frac{k}{(k+1)/2+i} \quad [0 \leq i \leq (k-3)/2]. \quad (4.3)$$

As an example, for $k = 3, 5, 7,$ and $9,$ (4.2) gives (4.1),

$$F_{3^{n+1}} = 25F_{3^n}^5 - 25F_{3^n}^3 + 5F_{3^n}, \quad (4.4)$$

$$F_{7^{n+1}} = 125F_{7^n}^7 - 175F_{7^n}^5 + 70F_{7^n}^3 - 7F_{7^n}, \quad (4.5)$$

and

$$F_{9^{n+1}} = 625F_{9^n}^9 - 1125F_{9^n}^7 + 675F_{9^n}^5 - 150F_{9^n}^3 + 9F_{9^n}, \quad (4.6)$$

respectively.

Proof of Proposition 4: First, let us write

$$F_{k^n}^k = \frac{1}{(\sqrt{5})^k} \sum_{j=0}^k \binom{k}{j} (-1)^j \beta^{jk^n} \alpha^{(k-j)k^n}, \quad (4.7)$$

where $\alpha = 1 - \beta = (1 + \sqrt{5})/2$. After several simple but tedious manipulations involving the use of the Binet forms for Fibonacci and Lucas numbers, and the relation $\alpha\beta = -1$, (4.7) yields the following identities which may be of some interest *per se*:

$$F_{k^n}^k = \frac{1}{5^{(k-1)/2}} \left[F_{k^{n+1}} + \sum_{j=1}^{(k-1)/2} \binom{k}{j} F_{(k-2j)k^n} \right] \quad (k \text{ odd}, n \geq 0), \quad (4.8)$$

$$F_{k^n}^k = \frac{1}{5^{k/2}} \left[(-1)^{k/2} \binom{k}{k/2} + \sum_{j=0}^{(k-2)/2} (-1)^j \binom{k}{j} L_{(k-2j)k^n} \right] \quad (k \text{ even}, n \geq 1). \quad (4.9)$$

By (4.8), we immediately obtain

$$F_{k^{n+1}} = 5^{(k-1)/2} F_{k^n}^k - \sum_{j=1}^{(k-1)/2} \binom{k}{j} F_{(k-2j)k^n} \quad (k \text{ odd}). \quad (4.10)$$

Then, using (4.10) along with Theorem 1 of [4] leads to the expression

$$F_{k^{n+1}} = 5^{(k-1)/2} F_{k^n}^k - \sum_{j=1}^{(k-1)/2} \sum_{i=0}^{(k-1)/2-j} (-1)^{(k-1)/2+i-j} \frac{k-2j}{(k+1)/2+i-j} 5^i \binom{k}{j} \binom{(k+1)/2+i-j}{2i+1} F_{k^n}^{2i+1} \quad (4.11)$$

which, after reversing the summation order, can be rewritten as

$$F_{k^{n+1}} = 5^{(k-1)/2} F_{k^n}^k - \sum_{i=0}^{(k-1)/2} 5^i A_{i,k} F_{k^n}^{2i+1}, \quad (4.12)$$

where

$$A_{i,k} = (-1)^i \sum_{j=1}^{(k-1)/2-i} (-1)^{(k-1)/2-j} \frac{k-2j}{(k+1)/2+i-j} \binom{k}{j} \binom{(k+1)/2+i-j}{2i+1}. \quad (4.13)$$

Since $A_{(k-1)/2,k} = 0$ by (1.3), expression (4.12) becomes

$$F_{k^{n+1}} = 5^{(k-1)/2} F_{k^n}^k - \sum_{i=0}^{(k-3)/2} 5^i A_{i,k} F_{k^n}^{2i+1}. \quad (4.14)$$

Now it remains to show that the numbers $A_{i,k}$ [defined by (4.13)] and the numbers $C_{i,k}$ [defined by (4.3)] coincide. To do this, consider the combinatorial identity

$$\begin{aligned} & \sum_{j=1}^m (-1)^j \frac{k-2j}{k-m-j} \binom{k}{j} \binom{k-m-j}{m-j} \\ &= -\frac{k}{k-m} \binom{k-m}{m} \quad [1 \leq m \leq (k-1)/2], \end{aligned} \quad (4.15)$$

which can be obtained by [5, p. 58], and replace m by $(k-1)/2-i$ in (4.15) to obtain the desired result $C_{i,k} = A_{i,k}$. Q.E.D.

5. CONCLUDING REMARKS

Some properties of the numbers F_{k^n} have been investigated in this note. In particular, expressions for these numbers in terms of products involving Lucas numbers have been established. Analogous expressions for L_{k^n} appeared to be rather unpleasant, so we confine ourselves to show some partial results whose proofs are left to the perseverance of the reader. In particular, we show the identity

$$L_{k^n} = 2 + (L_{k^2} - 2) \prod_{i=1}^{n-2} \left[\sum_{j=1}^{k/2} L_{(2j-1)k^{i+1}/2} \right]^2 \quad (k \text{ even, } n \geq 2) \quad (5.1)$$

which, for $k = 2$, reduces to

$$L_{2^n} = 2 + 5 \prod_{i=1}^{n-2} L_{2^i}^2 \quad (n \geq 2). \quad (5.2)$$

We also have

$$L_{3^n} = 4 \prod_{i=1}^{n-1} (L_{2 \cdot 3^i} + 1) \quad (n \geq 1). \quad (5.3)$$

Observe that the identity

$$F_{2 \cdot 3^n} = 8 \prod_{i=1}^{n-1} (L_{2 \cdot 3^i}^2 - 1) = 8 \prod_{i=1}^{n-1} (L_{4 \cdot 3^i} + 1) \quad (n \geq 1) \quad (5.4)$$

can be obtained either by (2.6) and (5.3), or by (3.4).

ACKNOWLEDGMENT

This work has been carried out in the framework of an agreement between the Italian PT Administration (Istituto Superiore PT) and the Fondazione Ugo Bordoni. The author wishes to express his gratitude to an anonymous referee whose suggestions have greatly improved this paper.

REFERENCES

1. R. M. Capocelli, ed. *Sequences*. New York: Springer-Verlag, 1990.
2. P. Filippini & H. T. Freitag. "The Zeckendorf Representation of F_{kn} / F_n ." In *Applications of Fibonacci Numbers* 5:217-19. Dordrecht: Kluwer, 1993.
3. V. E. Hoggatt, Jr. *Fibonacci and Lucas Numbers*. Boston: Houghton Mifflin, 1969; rpt. The Fibonacci Association, 1979.
4. D. Jennings. "Some Polynomial Identities for the Fibonacci and Lucas Numbers." *The Fibonacci Quarterly* 31.2 (1993):134-37.
5. J. Riordan. *Combinatorial Identities*. New York: Wiley, 1968.
6. Problem B-746. *The Fibonacci Quarterly* 31.3 (1993):278.

