

ON THE  $K^{\text{th}}$ -ORDER DERIVATIVE SEQUENCES OF FIBONACCI AND LUCAS POLYNOMIALS

Chizhong Zhou

Yueyang University, Yueyang, P.R. China, 414000

(Submitted December 1994)

1. INTRODUCTION

The Fibonacci polynomials  $u_n = u_n(x)$  and the Lucas polynomials  $v_n = v_n(x)$  are defined by the second-order linear recurrence relations

and 
$$\begin{aligned} u_n &= xu_{n-1} + u_{n-2} \quad (u_0 = 0, u_1 = 1), \\ v_n &= xv_{n-1} + v_{n-2} \quad (v_0 = 2, v_1 = x), \end{aligned} \tag{1.1}$$

where  $x$  is an indeterminate. Their  $k^{\text{th}}$ -order derivative sequences are defined as

$$u_n^{(k)} = u_n^{(k)}(x) = \frac{d^k}{dx^k} u_n(x) \quad \text{and} \quad v_n^{(k)} = v_n^{(k)}(x) = \frac{d^k}{dx^k} v_n(x).$$

Denote  $f_n = u_n(1)$ ,  $l_n = v_n(1)$ ,  $f_n^{(k)} = u_n^{(k)}(1)$ ,  $l_n^{(k)} = v_n^{(k)}(1)$ . P. Filipponi and A. F. Horadam ([1], [2]) considered  $f_n^{(k)}$  and  $l_n^{(k)}$  for  $k = 1, 2$  and obtained a series of results. By the end of [2], seven conjectures were presented for arbitrary  $k$ . In this paper we shall consider the more general cases,  $u_n^{(k)}$  and  $v_n^{(k)}$ , for arbitrary  $k$ . Our results will be generalizations of the results in [1] and [2]. As special cases of our results, the seven conjectures in [2] will be proved.

Following the symbols in [1] and [2], denote  $\Delta = \sqrt{x^2 + 4}$ ,  $\alpha = (x + \Delta) / 2$ ,  $\beta = (x - \Delta) / 2$ , so that  $\alpha + \beta = x$ ,  $\alpha\beta = -1$ ,  $\alpha - \beta = \Delta$ . It is well known that

$$u_n = (\alpha^n - \beta^n) / \Delta, \quad v_n = \alpha^n + \beta^n. \tag{1.2}$$

2. EXPRESSIONS FOR  $u_n^{(k)}$  AND  $v_n^{(k)}$  IN TERMS OF FIBONACCI AND LUCAS POLYNOMIALS

**Theorem 2.1:**

$$u_n^{(k)} = \frac{k!}{2\Delta^{2k}} (a_{n,k} u_n + b_{n,k} v_n), \tag{2.1}$$

where

$$a_{n,k} = \sum_{\substack{i=0 \\ 2|k-i}}^k \binom{k-i+n}{k-i} \Delta^{k-i} (c_{k,i} + d_{k,i}) + \sum_{\substack{i=0 \\ 2|k-i}}^k \binom{k-i+n}{k-i} \Delta^{k-i} (c_{k,i} - d_{k,i}), \tag{2.2}$$

and

$$b_{n,k} = \sum_{\substack{i=0 \\ 2|k-i}}^k \binom{k-i+n}{k-i} \Delta^{k-1-i} (c_{k,i} - d_{k,i}) + \sum_{\substack{i=0 \\ 2|k-i}}^k \binom{k-i+n}{k-i} \Delta^{k-1-i} (c_{k,i} + d_{k,i}), \tag{2.3}$$

where  $c_{k,i}$  and  $d_{k,i}$  ( $i = 0, 1, \dots, k$ ) satisfy the systems of linear equations

$$c_{k,i} + \binom{k+1}{1}\beta c_{k,i-1} + \dots + \binom{k+1}{i}\beta^i c_{k,0} = (-1)^i \binom{k+1}{i} \Delta^i \tag{2.4}$$

and

$$d_{k,i} + \binom{k+1}{1}\alpha d_{k,i-1} + \dots + \binom{k+1}{i}\alpha^i d_{k,0} = \binom{k+1}{i} \Delta^i. \tag{2.5}$$

Furthermore, for  $i = 0, 1, \dots, k$ , there exist polynomials  $p_{k,i}$  and  $q_{k,i}$  in  $x$ , with integer coefficients, which satisfy

$$c_{k,i} = p_{k,i}\alpha + q_{k,i} \quad \text{and} \quad d_{k,i} = p_{k,i}\beta + q_{k,i}. \tag{2.6}$$

**Proof:** Let the generating functions of  $\{u_n\}$  and  $\{u_n^{(k)}\}$  be  $U(t) = U(t, x) = \sum_{n=0}^{\infty} u_n t^n$  and  $U_k(t) = U_k(t, x) = \sum_{n=0}^{\infty} u_n^{(k)} t^n$ , respectively. It is well known that  $U(t) = t / (1 - xt - t^2)$ , hence,

$$U_k(t) = \frac{\partial^k}{\partial x^k} U(t) = k! t^{k+1} / (1 - xt - t^2)^{k+1}. \tag{2.7}$$

By partial fractions we have

$$t^{k+1} / (1 - xt - t^2)^{k+1} = \sum_{i=0}^k Q_{k,i} / (1 - \alpha t)^{k+1-i} + \sum_{i=0}^k R_{k,i} / (1 - \beta t)^{k+1-i}, \tag{2.8}$$

where  $Q_{k,i}$  and  $R_{k,i}$  are independent of  $t$ . Multiplying by  $\alpha^{k+1}(1 - \beta t)^{k+1}$ , we obtain

$$(\alpha t)^{k+1} / (1 - \alpha t)^{k+1} = (\alpha + t)^{k+1} \sum_{i=0}^k Q_{k,i} / (1 - \alpha t)^{k+1-i} + \varphi(t), \tag{2.9}$$

where the function  $\varphi(t)$  is analytic at the point  $t = \alpha^{-1}$  under the condition that  $t$  is considered as a complex variable (while  $x$  is a real constant). Since  $(\alpha t)^{k+1} / (1 - \alpha t)^{k+1} = [(1 - \alpha t)^{-1} - 1]^{k+1}$  and  $(\alpha + t)^{k+1} = [\Delta + \beta(1 - \alpha t)]^{k+1}$ , we can rewrite (2.9) as

$$\sum_{i=0}^{k+1} (-1)^i \binom{k+1}{i} (1 - \alpha t)^{-(k+1-i)} = \sum_{i=0}^{k+1} \binom{k+1}{i} \Delta^{k+1-i} \beta^i (1 - \alpha t)^i \cdot \sum_{i=0}^k Q_{k,i} (1 - \alpha t)^{-(k+1-i)} + \varphi(t).$$

Because of the uniqueness of the Laurent series [4] at the point  $t = \alpha^{-1}$  for the function  $(\alpha t)^{k+1} / (1 - \alpha t)^{k+1}$ , we can compare the coefficients of  $(1 - \alpha t)^{-(k+1-i)}$  ( $i = 0, 1, \dots, k$ ) of the two sides in the last equality to get

$$\sum_{j=0}^i \binom{k+1}{j} \Delta^{k+1-j} \beta^j Q_{k,i-j} = (-1)^i \binom{k+1}{i}. \tag{2.10}$$

Let

$$Q_{k,i} = \Delta^{-(k+1+i)} c_{k,i} \quad (i = 0, 1, \dots, k) \tag{2.11}$$

and substitute it into (2.10); then we get (2.4). For the same reason, it follows that

$$\sum_{j=0}^i \binom{k+1}{j} (-\Delta)^{k+1-j} \alpha^j R_{k,i-j} = (-1)^i \binom{k+1}{i}. \tag{2.12}$$

Let

$$R_{k,i} = (-\Delta)^{-(k+1+i)} d_{k,i} \quad (i = 0, 1, \dots, k) \tag{2.13}$$

and substitute it into (2.12); then we get (2.5).

Now we shall prove (2.6). From (2.4) and (2.5),  $c_{k,0} = d_{k,0} = 1$ ; hence, the conclusion holds for  $i = 0$ . Suppose the conclusion holds for  $0, 1, \dots, i-1$ . Then, from (2.4) and (2.5), we have

$$c_{k,i} = (-1)^i \binom{k+1}{i} \Delta^i - \sum_{j=1}^i \binom{k+1}{j} \beta^j (p_{k,i-j} \alpha + q_{k,i-j}) \tag{2.14}$$

and

$$d_{k,i} = \binom{k+1}{i} \Delta^i - \sum_{j=1}^i \binom{k+1}{j} \alpha^j (p_{k,i-j} \beta + q_{k,i-j}). \tag{2.15}$$

From (1.2), it is easy to show that  $\beta^j = -u_j \alpha + u_{j+1}$ ; hence,

$$\begin{aligned} \beta^j (p_{k,i-j} \alpha + q_{k,i-j}) &= -p_{k,i-j} \beta^{j-1} + q_{k,i-j} \beta^j \\ &= (p_{k,i-j} u_{j-1} - q_{k,i-j} u_j) \alpha + (q_{k,i-j} u_{j+1} - p_{k,i-j} u_j). \end{aligned}$$

For the same reason, we have

$$\alpha^j (p_{k,i-j} \beta + q_{k,i-j}) = (p_{k,i-j} u_{j-1} - q_{k,i-j} u_j) \beta + (q_{k,i-j} u_{j+1} - p_{k,i-j} u_j).$$

We can see that  $\Delta^i$  is a polynomial in  $x$  with integer coefficients for  $2|i$ , but  $\Delta^i = \Delta^{i-1}(x-2\beta)$  and  $(-\Delta)^i = \Delta^{i-1}(x-2\alpha)$  for  $2 \nmid i$ . By substituting the above results into (2.14) and (2.15), and by the inductive hypothesis, the conclusion is proved.

Now substituting (2.11), (2.13), and (2.6) into (2.8), then into (2.7), we get

$$\begin{aligned} U_k(t) &= \frac{k!}{\Delta^{2k}} \left[ \sum_{i=0}^k c_{k,i} \Delta^{k-1-i} / (1-\alpha t)^{k+1-i} + \sum_{i=0}^k d_{k,i} (-\Delta)^{k-1-i} / (1-\beta t)^{k+1-i} \right] \\ &= \frac{k!}{\Delta^{2k}} \left[ \sum_{2|k-i} (c_{k,i} / (1-\alpha t)^{k+1-i} - d_{k,i} / (1-\beta t)^{k+1-i}) \Delta^{k-1-i} \right. \\ &\quad \left. + \sum_{2 \nmid k-i} (c_{k,i} / (1-\alpha t)^{k+1-i} + d_{k,i} / (1-\beta t)^{k+1-i}) \Delta^{k-1-i} \right]. \end{aligned}$$

Expanding the right side of the last expression into power series in  $t$  and using (2.6), we obtain

$$u_n^{(k)} = \frac{k!}{\Delta^{2k}} \left[ \sum_{2|k-i} \binom{k-i+n}{k-i} \Delta^{k-i} (p_{k,i} u_{n+1} + q_{k,i} u_n) + \sum_{2 \nmid k-i} \binom{k-i+n}{k-i} \Delta^{k-1-i} (p_{k,i} v_{n+1} + q_{k,i} v_n) \right]. \tag{2.16}$$

It is easy to prove that  $u_{n+1} = (xu_n + v_n) / 2$ ,  $v_{n+1} = (\Delta^2 u_n + xv_n) / 2$ ; hence,

$$\begin{aligned} p_{k,i} u_{n+1} + q_{k,i} u_n &= ((p_{k,i} x + 2q_{k,i}) u_n + p_{k,i} v_n) / 2 \\ &= ((c_{k,i} + d_{k,i}) u_n + (c_{k,i} - d_{k,i}) \Delta^{-1} v_n) / 2, \end{aligned} \tag{2.17}$$

$$\begin{aligned} p_{k,i} v_{n+1} + q_{k,i} v_n &= (p_{k,i} \Delta^2 u_n + (p_{k,i} x + 2q_{k,i}) v_n) / 2 \\ &= ((c_{k,i} - d_{k,i}) \Delta u_n + (c_{k,i} + d_{k,i}) v_n) / 2. \end{aligned} \tag{2.18}$$

Substitute (2.17) and (2.18) into (2.16) and we are done.  $\square$

As an example, when  $k = 3$  and 4, Theorem 2.1 gives the following results:

$$\begin{aligned} c_{30} &= d_{30} = 1, & c_{31} &= -4\Delta - 4\beta, & d_{31} &= 4\Delta - 4\alpha, \\ c_{32} &= 6\Delta^2 + 16\beta\Delta + 10\beta^2, & d_{32} &= 6\Delta^2 - 16\alpha\Delta + 10\alpha^2, \\ c_{33} &= -4\Delta^3 - 24\beta\Delta^2 - 40\beta^2\Delta - 20\beta^3, & d_{33} &= 4\Delta^3 - 24\alpha\Delta^2 + 40\alpha^2\Delta - 20\alpha^3, \\ c_{30} + d_{30} &= 2, & c_{31} + d_{31} &= -4x, \\ c_{32} + d_{32} &= 6x^2 + 4, & c_{33} + d_{33} &= -4x^3 + 4x, \\ c_{30} - d_{30} &= 0, & c_{31} - d_{31} &= -4\Delta, \\ c_{32} - d_{32} &= 6x\Delta, & c_{33} - d_{33} &= (-4x^2 + 4)\Delta, \end{aligned}$$

$$\begin{aligned} a_{n3} &= \binom{2+n}{2}\Delta^2(-4x) + \binom{0+n}{0}(-4x^3+4x) + \binom{3+n}{3}\Delta^3 \cdot 0 + \binom{1+n}{1}\Delta \cdot 6x\Delta \\ &= -2(n^2+1)x^3 - 4(2n^2-3)x, \end{aligned}$$

$$\begin{aligned} b_{n3} &= \binom{2+n}{2}\Delta(-4\Delta) + \binom{0+n}{0}\Delta^{-1}(-4x^3+4)\Delta + \binom{3+n}{3}\Delta^2 \cdot 2 + \binom{1+n}{1}(6x^2+4) \\ &= \frac{1}{3}n(n^2+11)x^2 + \frac{4}{3}n(n^2-4), \end{aligned}$$

$$u_n^{(3)} = [-(6(n^2+1)x^3 + 12(2n^2-3)x)u_n + (n(n^2+11)x^2 + 4n(n^2-4))v_n] / \Delta^6; \tag{2.19}$$

in particular,

$$f_n^{(3)} = (n^2-1)(nl_n - 6f_n) / 25. \tag{2.20}$$

$$\begin{aligned} c_{40} &= d_{40} = 1, & c_{41} &= -5\Delta - 5\beta, & d_{41} &= 5\Delta - 5\alpha, \\ c_{42} &= 10\Delta^2 + 25\beta\Delta + 15\beta^2, & d_{42} &= 10\Delta^2 - 25\alpha\Delta + 15\alpha^2, \\ c_{43} &= -10\Delta^3 - 50\beta\Delta^2 - 75\beta^2\Delta - 35\beta^3, & d_{43} &= 10\Delta^3 - 50\alpha\Delta^2 + 75\alpha^2\Delta - 35\alpha^3, \\ c_{44} &= 5\Delta^4 + 50\beta\Delta^3 + 150\beta^2\Delta^2 + 175\beta^3\Delta + 70\beta^4, \\ d_{44} &= 5\Delta^4 - 50\alpha\Delta^3 + 150\alpha^2\Delta^2 - 175\alpha^3\Delta + 70\alpha^4, \\ c_{40} + d_{40} &= 2, & c_{41} + d_{41} &= -5x, \\ c_{42} + d_{42} &= 10x^2 + 10, & c_{43} + d_{43} &= -10x^3 - 5x, \\ c_{44} + d_{44} &= 5x^4 - 15x^2, & c_{40} - d_{40} &= 0, \\ c_{41} - d_{41} &= -5\Delta, & c_{42} - d_{42} &= 10x\Delta, \\ c_{43} - d_{43} &= (-10x^2 + 5)\Delta, & c_{44} - d_{44} &= (5x^3 - 15x)\Delta, \end{aligned}$$

$$\begin{aligned} a_{n4} &= \binom{4+n}{4}\Delta^4 \cdot 2 + \binom{2+n}{2}\Delta^2(10x^2+10) + \binom{0+n}{0}(5x^4-15x^2) \\ &\quad + \binom{3+n}{3}\Delta^3(-5\Delta) + \binom{1+n}{1}\Delta(-10x^2+5)\Delta \\ &= \frac{1}{12}(n^4+35n^2+24)x^4 + \frac{1}{3}(2n^4+25n^2-72)x^2 + \frac{4}{3}(n^4-10n^2+9), \end{aligned}$$

$$\begin{aligned}
 b_{n4} &= \binom{4+n}{4} \Delta^3 \cdot 0 + \binom{2+n}{2} \Delta(10x\Delta) + \binom{0+n}{0} \Delta^{-1}(5x^3 - 15x)\Delta \\
 &\quad + \binom{3+n}{3} \Delta^2(-5x) + \binom{1+n}{1} (-10x^3 - 5x) \\
 &= -\frac{5}{6}n(n^2 + 5)x^3 - \frac{5}{3}n(2n^2 - 11)x, \\
 u_n^{(4)} &= [(n^4 + 35n^2 + 24)x^4 + 4(2n^4 + 25n^2 - 72)x^2 + 16(n^4 - 10n^2 + 9)]u_n \\
 &\quad - (10n(n^2 + 5)x^3 + 20n(2n^2 - 11)x)v_n] / \Delta^8; \tag{2.21}
 \end{aligned}$$

in particular,

$$f_n^{(4)} = [(5n^4 - 5n^2 - 24)f_n - 2n(5n^2 - 17)\ell_n] / 125. \tag{2.22}$$

We observe that (2.6) can be verified by using the above results.

From  $v_n^{(k)} = mu_n^{(k-1)}$  (see 1° of Theorem 3.1 in the next section) and Theorem 2.1, we can obtain the expression for  $v_n^{(k)}$  in terms of  $u_n$  and  $v_n$ .

### 3. SOME IDENTITIES INVOLVING $u_n^{(k)}$ AND $v_n^{(k)}$

If we differentiate certain identities involving  $u_n$  and  $v_n$ , we can get the corresponding identities involving  $u_n^{(k)}$  and  $v_n^{(k)}$ .

**Theorem 3.1:**

$$1^\circ. v_n^{(k)} = mu_n^{(k-1)}, \tag{3.1}$$

$$2^\circ. u_n^{(k)} = xu_{n-1}^{(k)} + u_{n-2}^{(k)} + ku_{n-1}^{(k-1)}, \quad v_n^{(k)} = xv_{n-1}^{(k)} + v_{n-2}^{(k)} + kv_{n-1}^{(k-1)}, \tag{3.2}$$

$$3^\circ. v_n^{(k)} = u_{n+1}^{(k)} + u_{n-1}^{(k)}, \tag{3.3}$$

$$\Delta^2 u_n^{(k)} + 2kxu_n^{(k-1)} + k(k-1)u_n^{(k-2)} = v_{n+1}^{(k)} + v_{n-1}^{(k)}. \tag{3.4}$$

$$4^\circ. u_{m+n}^{(k)} = \sum_{i=0}^k \binom{k}{i} (u_{m+1}^{(k-i)} u_n^{(i)} + u_m^{(k-i)} u_{n-1}^{(i)}), \tag{3.5}$$

$$v_{m+n}^{(k)} = \sum_{i=0}^k \binom{k}{i} (v_{m+1}^{(k-i)} u_n^{(i)} + v_m^{(k-i)} u_{n-1}^{(i)}), \tag{3.6}$$

$$u_{m-n}^{(k)} = (-1)^n \sum_{i=0}^k \binom{k}{i} (u_m^{(k-i)} u_{n+1}^{(i)} - u_{m+1}^{(k-i)} u_n^{(i)}), \tag{3.7}$$

$$v_{m-n}^{(k)} = (-1)^n \sum_{i=0}^k \binom{k}{i} (v_{m+1}^{(k-i)} u_n^{(i)} - u_m^{(k-i)} v_{n+1}^{(i)}); \tag{3.8}$$

in particular,

$$u_{-n}^{(k)} = (-1)^{n-1} u_n^{(k)}, \tag{3.9}$$

$$v_{-n}^{(k)} = (-1)^n v_n^{(k)}, \tag{3.10}$$

$$u_{2n}^{(k)} = \sum_{i=0}^k \binom{k}{i} u_n^{(k-i)} v_n^{(i)}, \tag{3.11}$$

$$v_{2n}^{(k)} = 2 \sum_{i=0}^{k-1} \binom{k-1}{i} v_n^{(k-i)} v_n^{(i)}; \tag{3.12}$$

$$u_{2n+1}^{(k)} = \sum_{i=0}^k \binom{k}{i} u_{n+1}^{(k-i)} v_n^{(i)}; \tag{3.13}$$

$$v_{2n+1}^{(k)} = \sum_{i=0}^k \binom{k}{i} v_{n+1}^{(k-i)} v_n^{(i)} - (-1)^n \delta_{k,1} \quad (\delta \text{ is the Kronecker function}); \tag{3.14}$$

$$5^\circ. \quad u_{m+n}^{(k)} + (-1)^n u_{m-n}^{(k)} = \sum_{i=0}^k \binom{k}{i} u_m^{(k-i)} v_n^{(i)}; \tag{3.15}$$

$$u_{m+n}^{(k)} - (-1)^n u_{m-n}^{(k)} = \sum_{i=0}^k \binom{k}{i} v_m^{(k-i)} u_n^{(i)}; \tag{3.16}$$

$$v_{m+n}^{(k)} + (-1)^n v_{m-n}^{(k)} = \sum_{i=0}^k \binom{k}{i} v_m^{(k-i)} v_n^{(i)}; \tag{3.17}$$

$$v_{m+n}^{(k)} - (-1)^n v_{m-n}^{(k)} = \sum_{i=0}^k \binom{k}{i} u_m^{(k-i)} (v_{n+1}^{(i)} + v_{n-1}^{(i)}); \tag{3.18}$$

$$6^\circ. \quad xv_n^{(k)} = (n-k+1)v_n^{(k-1)} - 2(v_{n-1}^{(k)} + u_{n-1}^{(k-1)}). \tag{3.19}$$

*Proof:* 1°. This can be obtained by differentiating the identity  $v_n^{(1)} = nu_n$ , which had been proved in [1].

2°. By differentiating (1.1).

3° ~ 5°. By differentiating the following identities, which can be seen in [5] or can be derived from (1.2):

$$\begin{array}{ll} v_n = u_{n+1} + u_{n-1}, & \Delta^2 u_n = v_{n+1} + v_{n-1}, \\ u_{m+n} = u_{m+1} u_n + u_m u_{n-1}, & v_{m+n} = v_{m+1} u_n + v_m u_{n-1}, \\ u_{m-n} = (-1)^n (u_m u_{n+1} - u_{m+1} u_n), & v_{m-n} = (-1)^n (u_{m+1} v_n - u_m v_{n+1}), \\ u_{m+n} + (-1)^n u_{m-n} = u_m v_n, & u_{m+n} - (-1)^n u_{m-n} = v_m u_n, \\ v_{m+n} + (-1)^n v_{m-n} = v_m v_n, & v_{m+n} - (-1)^n v_{m-n} = \Delta^2 u_m u_n = u_m (v_{n+1} + v_{n-1}), \\ u_{-n} = (-1)^{n-1} u_n, & v_{-n} = (-1)^n v_n, \\ u_{2n} = u_n v_n, & v_{2n} = v_n^2 - 2(-1)^n, \\ u_{2n+1} = u_{n+1} v_n - (-1)^n, & v_{2n+1} = v_{n+1} v_n - (-1)^n x. \end{array}$$

6°. From the well-known identity  $v_n = xu_n + 2u_{n-1}$ , we get  $xmu_n = nv_n - 2((n-1)u_{n-1} + u_{n-1})$ , that is,  $xv_n^{(1)} = nv_n - 2(v_{n-1}^{(1)} + u_{n-1})$ , and the proof is finished by differentiating the last expression. □

Let  $x = 1$  in 1°, 2°, 3°, and 6° of Theorem 3.1; then Conjectures 1-5 in [2] and [3] are proved.

#### 4. SOME CONGRUENCE RELATIONS AND MODULAR PERIODICITIES

First, we introduce some concepts and lemmas. Set polynomials

$$g(t) = t^k - a_1 t^{k-1} - \dots - a_{k-1} t - a_k \tag{4.1}$$

and

$$\tilde{g}(t) = 1 - a_1 t - \dots - a_{k-1} t^{k-1} - a_k t^k. \tag{4.2}$$

Obviously,  $g(t) = t^k \tilde{g}(1/t)$  and  $\tilde{g}(t) = t^k g(1/t)$ . The set of homogeneous linear recurrence sequences  $\{g_n\}$  of order  $k$  [each of which has  $g(t)$  as its characteristic polynomial] defined by

$$g_{n+k} = a_1 g_{n+k-1} + \dots + a_{k-1} g_{n+1} + a_k g_n \tag{4.3}$$

is denoted by  $\Omega(g(t)) = \Omega(a_1, \dots, a_k)$ . The sequence  $\{w_n\} \in \Omega(g(t))$  is called the **principal sequence** in  $\Omega(g(t))$  if it has the initial values  $w_0 = w_1 = \dots = w_{k-2} = 0, w_{k-1} = 1$ .

**Lemma 4.1:** Let  $\{w_n\}$  be the principal sequence in  $\Omega(g(t))$ ; then its generating function is

$$W(t) = t^{k-1} / \tilde{g}(t) \tag{4.4}$$

(see [6], p. 137).

In the following discussions, we suppose that  $a_1, \dots, a_k$  are all integers. Let  $\{g_n\}$  be an integer sequence in  $\Omega(g(t))$  and  $m$  be an integer greater than one. Denote the period of  $\{g_n\}$  modulo  $m$  by  $P(m, g_n)$ . If there exists a positive integer  $\lambda$  such that

$$t^\lambda \equiv 1 \pmod{m, g(t)}, \tag{4.5}$$

then the least positive integer  $\lambda$  such that (4.5) holds is called the **period of  $g(t)$  modulo  $m$**  and is denoted by  $P(m, g(t))$ .

We point out that

$$P(m, g(t)) = P(m, \tilde{g}(t)) \text{ for } \gcd(m, a_k) = 1. \tag{4.6}$$

To show (4.6), it is sufficient to show that  $g(t)|(t^\lambda - 1) \pmod{m}$  iff  $\tilde{g}(t)|(t^\lambda - 1) \pmod{m}$ . Assume that  $\tilde{g}(t)|(t^\lambda - 1) \pmod{m}$ . Then we have  $t^\lambda - 1 = h(t)\tilde{g}(t) + m \cdot r(t)$ , where  $h(t)$  and  $r(t) \in Z(t)$  (the set of polynomials with integer coefficients). Replacing  $t$  with  $1/t$ , we obtain  $(1/t)^\lambda - 1 = h(1/t)\tilde{g}(1/t) + m \cdot r(1/t)$ . Multiplying by  $t^\lambda$ , we then have  $-(t^\lambda - 1) = t^{\lambda-k}h(1/t)g(t) + m \cdot t^\lambda r(1/t)$ . Since  $\gcd(m, a_k) = 1$ , the degree of  $\tilde{g}(t) \pmod{m}$  is  $k$ . This leads to  $t^{\lambda-k}h(1/t)$  and  $t^\lambda r(1/t) \in Z(t)$ . Hence,  $g(t)|(t^\lambda - 1) \pmod{m}$ . The converse can be proved in the same way.

Let  $B(t) = 1/\tilde{g}(t) = \sum_{n=0}^{\infty} b_n t^n$ . Let  $\{w_n\}$  be the principal sequence in  $\Omega(g(t))$ . Then, from (4.4), we have  $w_n = b_{n-k+1}$ ; and therefore,  $P(m, w_n) = P(m, b_n)$ . Corollary 2 in [7] means that  $P(m, b_n) = P(m, \tilde{g}(t))$ .\* Therefore,

$$P(m, w_n) = P(m, \tilde{g}(t)). \tag{4.7}$$

From (4.6) and (4.7), we obtain

**Lemma 4.2:** Let  $\{w_n\}$  be the principal sequence in  $\Omega(g(t)) = \Omega(a_1, \dots, a_k)$ ,  $\gcd(m, a_k) = 1$ . Then

$$P(m, w_n) = P(m, g(t)). \tag{4.8}$$

Using the footnote and (4.6), Theorems 17, 21, and 15 in [7] can be rewritten as Lemmas 4.3, 4.4, and 4.5, respectively.

\* In [7] the period of  $\{b_n\}$  modulo  $m$  is referred to as the period of its generating function  $B(t) = 1/\tilde{g}(t)$  modulo  $m$ . Hence, the concept "the period of  $1/\tilde{g}(t)$  modulo  $m$ " stated in [7] should be translated into " $P(m, \tilde{g}(t))$ " in this paper.

**Lemma 4.3:** Let  $\varphi(t)$  be a monic polynomial with integer coefficients,  $p$  be a prime,  $p \nmid \varphi(0)$ , and  $\varphi(t)$  be irreducible modulo  $p$ ; then, for  $p^{r-1} < s \leq p^r$  ( $r \geq 1$ ),

$$P(p^m, \varphi(t)^s) = p^{m+r-1} \cdot P(p, \varphi(t)). \tag{4.9}$$

**Lemma 4.4:** Let  $\varphi(t)$  be a monic polynomial with integer coefficients,  $p$  be an odd prime,  $p \nmid \varphi(0)$ , and  $\varphi(t)$  be irreducible modulo  $p$ . Assume  $h_r(t) = \prod_{i=1}^r \Psi_i(t)$ , where  $\Psi_i(t) \equiv \varphi(t)^s \pmod{p}$  ( $i = 1, \dots, r$ ). For fixed  $s, r \geq 1$ , if there exists an integer  $T > 1$  such that

$$(T-1)s \leq p^{r-1} < Ts < (T+1)s \leq p^r, \tag{4.10}$$

then, for every  $\tau$  satisfying  $p^{r-1} < \tau s \leq p^r$ , it follows that

$$P(p^m, h_r(t)) = P(p^m, \varphi(t)^{\tau s}) = p^{m+r-1} \cdot P(p, \varphi(t)). \tag{4.11}$$

**Lemma 4.5:** Let  $\varphi(t)$  be a monic polynomial with integer coefficients,  $p$  be an odd prime,  $p \nmid \varphi(0)$ . If  $P(p, \varphi(t)) = P(p^2, \varphi(t)) = \dots = P(p^i, \varphi(t)) \neq P(p^{i+1}, \varphi(t))$ , then  $m > i$  leads to

$$P(p^m, \varphi(t)) = p^{m-i} \cdot P(p^i, \varphi(t)). \tag{4.12}$$

**Lemma 4.6:** Let  $p$  be an odd prime, for  $j = 1, 2$ ,  $\varphi_j(t)$  be a monic polynomial with integer coefficients,  $p \nmid \varphi_j(0)$ , and  $\varphi_j(t)$  be irreducible modulo  $p$ . Assume  $h_r(t) = \prod_{i=1}^r \Psi_i(t)$ , where  $\Psi_i(t) \equiv \varphi_1(t)^s \varphi_2(t)^s \pmod{p}$  ( $i = 1, \dots, r$ ),  $\gcd(\varphi_1(t), \varphi_2(t)) = 1 \pmod{p}$ . For fixed  $s, r \geq 1$ , if there exists an integer  $T > 1$  such that (4.10) holds, then for every  $\tau$  satisfying  $p^{r-1} < \tau s \leq p^r$  it follows that

$$P(p^m, h_r(t)) = P(p^m, \varphi_1(t)^{\tau s} \varphi_2(t)^{\tau s}) = p^{m+r-1} \cdot \text{lcm}\{P(p, \varphi_1(t)), P(p, \varphi_2(t))\}. \tag{4.13}$$

**Proof:** Denote  $P(p, \varphi_j(t)) = \lambda_j$  ( $j = 1, 2$ ),  $\text{lcm}\{\lambda_1, \lambda_2\} = \lambda$ . Since  $h_r(t) \equiv \varphi_1(t)^{\tau s} \varphi_2(t)^{\tau s} \pmod{p}$ ,  $\gcd(\varphi_1(t), \varphi_2(t)) = 1 \pmod{p}$ , we have  $P(p, h_r(t)) = \text{lcm}\{P(p, \varphi_1(t)^{\tau s}), P(p, \varphi_2(t)^{\tau s})\}$ . By Lemma 4.3,  $P(p, \varphi_j(t)^{\tau s}) = p^r \lambda_j$ ; hence,  $P(p, h_r(t)) = p^r \lambda$ .

Because  $T$  is the least  $\tau$  satisfying  $p^{r-1} < \tau s \leq p^r$  from (4.10), we get  $h_T(t) | h_r(t)$ ; therefore,  $P(p^m, h_T(t)) | P(p^m, h_r(t))$ . By Lemma 4.5,  $P(p^m, h_r(t)) | p^{m-1} \cdot P(p, h_r(t)) = p^{m+r-1} \lambda$ . By the same lemma, if we can show  $P(p^2, h_T(t)) \neq P(p, h_T(t)) = p^r \lambda$ , then  $P(p^m, h_r(t)) = p^{m+r-1} \lambda$  and (4.13) holds.

Now we can rewrite  $\Psi_i(t) = \varphi_1(t)^s \varphi_2(t)^s - p \theta_i(t)$ ,  $i = 1, \dots, T$ . Hence,

$$h_T(t) \equiv \varphi_1(t)^{sT} \varphi_2(t)^{sT} - p \varphi_1(t)^{s(T-1)} \cdot \varphi_2(t)^{s(T-1)} \cdot \zeta(t) \pmod{p^2}, \text{ where } \zeta(t) = \sum_{i=0}^T \theta_i(t).$$

Then  $h_T(t)[\varphi_1(t)^s \varphi_2(t)^s + p \zeta(t)] \equiv \varphi_1(t)^{sT+s} \varphi_2(t)^{sT+s} \pmod{p^2}$ . Therefore,

$$\frac{t^{p^r \lambda} - 1}{h_T(t)} \equiv \frac{t^{p^r \lambda} - 1}{\varphi_1(t)^{sT} \varphi_2(t)^{sT}} + \frac{p(t^{p^r \lambda} - 1)\zeta(t)}{\varphi_1(t)^{sT+s} \varphi_2(t)^{sT+s}} \pmod{p^2}. \tag{4.14}$$

From (4.10) and Lemma 4.3, we know that  $P(p, \varphi_j(t)^{sT+s}) = p^r \cdot P(p, \varphi_j(t)) = p^r \lambda_j$ ; thus,  $\varphi_j(t)^{sT+s} \equiv (t^{p^r \lambda} - 1) \pmod{p}$ . From  $\gcd(\varphi_1(t), \varphi_2(t)) = 1 \pmod{p}$ , it follows that

$$\varphi_1(t)^{sT+s} \varphi_2(t)^{sT+s} \equiv (t^{p^r \lambda} - 1) \pmod{p},$$

and so



$$\varphi_1(t)^{sT+s} \varphi_2(t)^{sT+s} | p(t^{p^\lambda} - 1) \pmod{p^2}.$$

Assume that  $P(p^2, h_r(t)) = p^r \lambda$ , then  $h_r(t) | (t^{p^\lambda} - 1) \pmod{p^2}$ . From equation (4.14), we get  $\varphi_j(t)^{sT} | (t^{p^\lambda} - 1) \pmod{p^2}$ ; this leads to  $P(p^2, \varphi_j(t)^{sT}) | p^r \lambda$ . But from Lemma 4.3 we have  $P(p^2, \varphi_j(t)^{sT}) = p^{r+1} \lambda_j$ . This leads to the contradiction that  $p^{r+1} \lambda | p^r \lambda$ .  $\square$

In the following discussions of this section when the divisibilities of  $u_n^{(k)}$  and  $v_n^{(k)}$  are considered, we assume  $x$  takes integer values only.

**Theorem 4.1:**

$$u_n^{(k)} \equiv v_n^{(k)} \equiv 0 \pmod{k!}. \tag{4.15}$$

**Proof:** Denote

$$F_k(t) = (t^2 - xt - 1)^{k+1}. \tag{4.16}$$

Let  $\{w_n\}$  be the principal sequence in  $\Omega(F_k(t))$ . From Lemma 4.1, the generating function of  $\{w_n\}$  is

$$W(t) = t^{2k+1} / (1 - xt - t^2)^{k+1}. \tag{4.17}$$

Comparing (2.7) to (4.17), we get

$$u_n^{(k)} = k! w_{n+k}. \tag{4.18}$$

Because  $\{w_n\}$  is an integer sequence, we have  $u_n^{(k)} \equiv 0 \pmod{k!}$ , and from (3.3) we get  $v_n^{(k)} \equiv 0 \pmod{k!}$ .  $\square$

**Theorem 4.2:**

$$v_n^{(k)} \equiv 0 \pmod{n} \quad (k \geq 1). \tag{4.19}$$

This follows from (3.1).

The results of the last two theorems are generalizations of the results of Conjectures 6-7 in [2].

**Theorem 4.3:** Let  $p$  be an odd prime,  $p > k$ .

1°. If  $p \nmid \Delta^2$ , then

$$P(p^m, u_n^{(k)}) = P(p^m, v_n^{(k)}) = p^m \cdot P(p, u_n) = p^m \cdot P(p, v_n). \tag{4.20}$$

2°. If  $p \mid \Delta^2$  and  $p^{r-1} < 2k + 2 < p^r$  ( $r = 1$  or  $2$ ), then

$$P(p^m, u_n^{(k)}) = 4p^{m+r-1}. \tag{4.21}$$

3°. If  $p \mid \Delta^2$  and  $p^{r-1} < 2k < p^r$  ( $r = 1$  or  $2$ ), then

$$P(p^m, v_n^{(k)}) = 4p^{m+r-1}. \tag{4.22}$$

**Proof:** Denote  $f(t) = t^2 - xt - 1$ . From Lemma 4.2, (4.18), and (4.16), for  $p > k$ , we have  $P(p, u_n) = P(p, f(t))$  and  $P(p^m, u_n^{(k)}) = P(p^m, F_k(t))$ .

1°. Let  $p \nmid \Delta^2$ . From  $v_n = u_{n+1} + u_{n-1}$  and  $\Delta^2 u_n = v_{n+1} + v_{n-1}$ , it follows that  $P(p, u_n) = P(p, v_n) = \lambda$ .

When  $f(t)$  is irreducible modulo  $p$ , the conclusion  $P(p^m, u_n^{(k)}) = p^m \lambda$  can be proved by letting  $\varphi(t) = f(t)$ ,  $s = k + 1$ ,  $r = 1$  in Lemma 4.3. When  $f(t) \equiv (t - a)(t - b)$ ,  $a \not\equiv b \pmod{p}$ , the same conclusion can be proved by letting  $\varphi_1(t) = t - a$ ,  $\varphi_2(t) = t - b$ ,  $s = r = 1$ ,  $\tau = k + 1$  in Lemma 4.6.

We now prove  $P(p^m, v_n^{(k)}) = p^m \lambda$ . From (3.3), we can see that  $P(p^m, v_n^{(k)}) | P(p^m, u_n^{(k)})$ . On the other hand, from  $u_n = (v_{n+1} + v_{n-1}) / \Delta^2$ , by differentiating, we can obtain

$$u_n^{(k)} = \sum_{i=0}^k \binom{k}{i} (v_{n+1}^{(k-i)} + v_{n-1}^{(k-i)}) M_i(x) / \Delta^{2i+2}, \tag{4.23}$$

where  $M_i(x)$  is a polynomial in  $x$  with integer coefficients that are independent of  $n$ . We see that (3.2) implies  $P(p^m, v_n^{(i-1)}) | P(p^m, v_n^{(i)})$ . Hence, for  $i = 0, 1, \dots, k$ ,  $P(p^m, v_n^{(k-i)}) | P(p^m, v_n^{(k)})$ . From (4.23), it follows that  $P(p^m, u_n^{(k)}) | P(p^m, v_n^{(k)})$ . Thus,  $P(p^m, v_n^{(k)}) = P(p^m, u_n^{(k)}) = p^m \lambda$ .

2°. Let  $p | \Delta^2$ , then  $f(t) \equiv (t - x/2)^2 \pmod{p}$ . From  $x^2 \equiv -4$ , we get  $(x/2)^2 \equiv -1 \pmod{p}$ . Hence,  $P(p, t - x/2) = \text{ord}_p(x/2) = 4$ .\* In Lemma 4.4, if we take  $\varphi(t) = t - x/2$ ,  $h_\tau(t) = F_k(t) \equiv \varphi(t)^{2k+2} \pmod{p}$ ,  $s = 2$ ,  $r = 1$  or  $2$ ,  $\tau = k + 1$ , then we get the required result.

3°. Using the result of 2°, it follows that  $P(p^m, v_n^{(k)}) = P(p^m, mu_n^{(k-1)}) | \text{lcm}\{P(p^m, n), P(p^m, u_n^{(k-1)})\} = 4p^{m+r-1}$  when  $p^{r-1} < 2k < p^r$  ( $r = 1$  or  $2$ ). Since  $v_n = \alpha^n + \beta^n \equiv 2(x/2)^n \pmod{p}$ , then  $4 = P(p, v_n) | P(p^m, v_n^{(k)})$ , and we have  $P(p^m, v_n^{(k)}) = 4p^M$ . We want to show that  $M = m + r - 1 = m + 1$  for  $r = 2$ , or  $= m$  for  $r = 1$ . First, let  $r = 2$ . If it would not be the case, that is, if  $M \leq m$ , then if we replace  $n$  by  $n + 4p^m$  in (3.19) we have

$$xv_n^{(k)} \equiv (n + 4p^m - k + 1)v_n^{(k-1)} - 2[v_{n-1}^{(k)} + u_{n+4p^m-1}^{(k-1)}] \pmod{p^m}.$$

Subtracting this from  $xv_n^{(k)} \equiv (n - k + 1)v_n^{(k-1)} - 2[v_{n-1}^{(k)} + u_{n-1}^{(k-1)}] \pmod{p^m}$ , we get  $u_{n+4p^m-1}^{(k-1)} - u_{n-1}^{(k-1)} \equiv 2p^m v_n^{(k-1)} \equiv 0 \pmod{p^m}$ . This means that  $P(p^m, u_n^{(k-1)}) | 4p^m$  for  $r = 2$ . But, by 2°, we should have  $P(p^m, u_n^{(k-1)}) = 4p^{m+1}$  for  $r = 2$ . A contradiction!

Next, let  $r = 1$ . The least  $k$  satisfying  $1 < 2k < p$  is 1. Recalling that  $P(p^m, v_n^{(1)}) | P(p^m, v_n^{(k)})$ , we need only prove that  $M = m$  for  $k = 1$ . On the contrary, suppose  $M \leq m - 1$ . then

$$v_{n+4p^{m-1}}^{(1)} - v_n^{(1)} = (n + 4p^{m-1})u_{n+4p^{m-1}} - mu_n \equiv 0 \pmod{p^m}.$$

Expanding  $u_n$  in (1.2) into the polynomial in  $x$ ,  $\Delta$ , and noting  $p | \Delta^2$ , we obtain

$$mu_n = n \sum_{i=0}^{[(n-1)/2]} \binom{n}{2i+1} (x/2)^{n-2i-1} (\Delta/2)^{2i} \equiv n \sum_{i=0}^{m-1} \binom{n}{2i+1} (x/2)^{n-2i-1} (\Delta/2)^{2i} \pmod{p^m} \tag{4.24}$$

and

$$(n + 4p^{m-1})u_{n+4p^{m-1}} \equiv (n + 4p^{m-1}) \sum_{i=0}^{m-1} \binom{n + 4p^{m-1}}{2i+1} (x/2)^{n+4p^{m-1}-2i-1} (\Delta/2)^{2i} \pmod{p^m}. \tag{4.25}$$

When  $m > 1$ , since

\* Let  $m$  and  $a$  be integers greater than one,  $\text{gcd}(m, a) = 1$ . The least positive integer  $\lambda$  satisfying  $a^\lambda \equiv 1 \pmod{m}$  is called the order of  $a$  modulo  $m$  and is denoted by  $\text{ord}_m(a)$ . Since  $t^\lambda - 1 = [(t - a) + a]^\lambda - 1 \equiv a^\lambda - 1 \pmod{(t - a)}$ , we have  $P(m, t - a) = \text{ord}_m(a)$ .

$$(n + 4p^{m-1}) \binom{n + 4p^{m-1}}{2i+1} \equiv n \binom{n}{2i+1} \pmod{p^{m-1}} \quad \text{and} \quad p \mid \Delta^{2i} \text{ for } i \geq 1,$$

and furthermore,  $(x/2)^4 = 1 \pmod{p}$  implies  $(x/2)^{4p^{m-1}} \equiv 1 \pmod{p^m}$ , (4.25) can be reduced to

$$(n + 4p^{m-1})u_{n+4p^{m-1}} \equiv (n + 4p^{m-1})^2(x/2)^{n-1} + n \sum_{i=1}^{m-1} \binom{n}{2i+1} (x/2)^{n-2i-1} (\Delta/2)^{2i} \pmod{p^m}. \quad (4.26)$$

Subtract (4.24) from (4.26) to get

$$(n + 4p^{m-1})u_{n+4p^{m-1}} - nu_n \equiv 8np^{m-1}(x/2)^{n-1} \not\equiv 0 \pmod{p^m} \text{ for } p \nmid n.$$

This is a contradiction!

When  $m = 1$ , from (4.24) and (4.25), we obtain

$$(n + 4)u_{n+4} - nu_n \equiv 8(n + 2)(x/2)^{n-1} \not\equiv 0 \pmod{p}$$

for  $n \not\equiv -2 \pmod{p}$ . This is also a contradiction!  $\square$

From Theorem 4.3, we can obtain many specific congruences. For this, we introduce another concept. Let  $\{g_n\}$  be an integer sequence. If there exists a positive integer  $s$ , a nonnegative integer  $n_0$ , and an integer  $c$ ,  $\gcd(m, c) = 1$ , such that

$$g_{n+s} \equiv cg_n \pmod{m} \quad \text{iff } n \geq n_0, \quad (4.27)$$

then the least positive integer  $s$  satisfying (4.27) is called **the constrained period of  $\{g_n\}$  modulo  $m$**  and is denoted by  $s = P'(m, g_n)$ . The number  $c$  is called the **multiplier**.

**Lemma 4.7:** Let  $\{w_n\}$  be the principal sequence in  $\Omega(F_k(t))$ , where  $F_k(t)$  is denoted by (4.16). Then  $P'(m, w_n) = s$  exists and the multiplier  $c$  is equal to  $w_{s+2k+1} \pmod{m}$ . Furthermore, if  $r = \text{ord}_m(c)$ , then  $P(m, w_n) = sr$ , and the structure of  $\{w_n \pmod{m}\}$  in a period is as follows:

$$\begin{cases} 0, \dots, 0, 1, & w_{2k+2}, & w_{2k+3}, & \dots, & w_{s-1}, \\ 0, \dots, 0, c, & cw_{2k+2}, & cw_{2k+3}, & \dots, & cw_{s-1}, \\ \dots & & & & \\ 0, \dots, 0, c^{r-1}, & c^{r-1}w_{2k+2}, & c^{r-1}w_{2k+3}, & \dots, & c^{r-1}w_{s-1}. \end{cases} \quad (4.28)$$

**Proof:** Because  $\{w_n\}$  is periodic, it must be constrained periodic [in the most special case, the multiplier  $c$  may be equal to 1  $\pmod{m}$ ]. We have  $w_0 = \dots = w_{2k} = 0$  and  $w_{2k+1} = 1$ . Replacing  $n$  by  $2k + 1$  in the expression

$$w_{n+s} \equiv cw_n \pmod{m}, \quad (4.29)$$

we obtain  $c \equiv w_{s+2k+1} \pmod{m}$ . By induction, from (4.29), we can get

$$w_{n+js} \equiv c^j w_n \pmod{m}. \quad (4.30)$$

If  $j = r = \text{ord}_m(c)$ , then (4.30) becomes  $w_{n+rs} \equiv w_n \pmod{m}$ . This means that  $P(m, w_n) = sr$ . In (4.30), let  $j$  be  $0, 1, \dots, r - 1$  and  $n$  be  $0, 1, \dots, s - 1$ ; then (4.28) follows.  $\square$

From Lemma 4.7, (4.18), and (3.1), we obtain

**Theorem 4.4:** Let  $\{w_n\}$  be the principal sequence in  $\Omega(F_k(t))$ , where  $F_k(t)$  is denoted by (4.16), and let  $p$  be an odd prime,  $p > k$ ,  $P'(p^m, w_n) = s$ . If  $w_n \equiv 0 \pmod{p^m}$  for  $n \equiv i \pmod{s}$ , then

$$u_n^{(k)} \equiv 0 \pmod{p^m} \text{ for } n \equiv i - k \pmod{s}$$

and

$$v_n^{(k+1)} \equiv 0 \pmod{p^m} \text{ for } n \equiv i - k \pmod{s} \text{ or } n \equiv 0 \pmod{p^m}.$$

Furthermore, if  $\lambda p^r \equiv i - k \pmod{s}$ , then  $v_{\lambda p^r}^{(k+1)} \equiv 0 \pmod{p^{m+r}}$ .

**Example 1:** Let  $x = 1, p = 3$ . Then  $\Delta^2 = 5, p \nmid \Delta^2$ . Hence, from (4.20), we obtain  $P(3^m, f_n^{(k)}) = P(3^m, \ell_n^{(k)}) = 3^m \cdot P(3, f_n) = 8 \cdot 3^m$  for  $k = 1, 2$ .

**Example 2:** Let  $x = 1, p = 5$ . Then  $p \mid \Delta^2 = 5$ . Hence, from (4.21), we get  $P(5^m, f_n^{(k)}) = 4 \cdot 5^{m+1}$  for  $k = 2, 3, 4$ , or  $4 \cdot 5^m$  for  $k = 1$  and, from (4.22), we get  $P(5^m, \ell_n^{(k)}) = 4 \cdot 5^{m+1}$  for  $k = 3, 4$  or  $4 \cdot 5^m$  for  $k = 1, 2$ .

**Example 3:** We show that  $f_n^{(2)} \equiv 0 \pmod{10}$  iff  $n \equiv 0, \pm 1, \pm 2 \pmod{25}$ , and  $\ell_n^{(3)} \equiv 0 \pmod{30}$  iff  $n \equiv \pm 1, \pm 2 \pmod{25}$  or  $n \equiv 0 \pmod{5}$ .

**Proof of Example 3:** We have  $F_2(t) = (t^2 - t - 1)^3 = t^6 - 3t^5 + 5t^3 - 3t - 1 \equiv t^6 - 3t^5 - 3t - 1 \pmod{5}$  for  $x = 1$ . Let  $\{w_n\}$  be the principal sequence in  $\Omega(F_2(t))$ . Then  $w_{n+6} \equiv 3w_{n+5} + 3w_{n+1} + w_n \pmod{5}$ .

Calculate  $\{w_n \pmod{5}\}_0^\infty$  according to the last congruence:

$$0, 0, 0, 0, 0, 1, -2, -1, 2, 1, 1, -2, -1, 2, 1, 2, 1, -2, -1, 2, -2, -1, 2, 1, -2, 0, 0, 0, 0, 0, -2, \dots \pmod{5}.$$

This implies that  $s = P'(5, w_n) = 25$  and  $w_n \equiv 0 \pmod{5}$  iff  $n \equiv 0, 1, 2, 3, 4 \pmod{25}$ . Hence, the example is proved by Theorem 4.1 and Theorem 4.4.

### 5. EVALUATION OF SOME SERIES INVOLVING $u_n^{(k)}$ AND $v_n^{(k)}$

**Lemma 5.1:**

$$1^\circ. \sum_{i=0}^n u_i = (u_{n+1} + u_n - 1) / x \quad (x \neq 0). \tag{5.1}$$

$$2^\circ. \sum_{i=0}^n v_i = (v_{n+1} + v_n - 2) / x + 1 \quad (x \neq 0). \tag{5.2}$$

$$3^\circ. \sum_{i=0}^n \binom{n}{i} x^i h_{i+r} = h_{2n+r}, \text{ where } \{h_n\} \text{ is } \{u_n\} \text{ or } \{v_n\}. \tag{5.3}$$

$$4^\circ. \sum_{i=0}^n (-1)^i \binom{n}{i} h_{2i+r} = (-1)^n x^n h_{n+r}, \text{ where } \{h_n\} \text{ is } \{u_n\} \text{ or } \{v_n\}. \tag{5.4}$$

$$5^\circ. \sum_{i=0}^n \binom{n}{i} u_{2i+r} = (x^2 + 4)^{n/2} u_{n+r} \text{ for } 2 \mid n, \text{ or } (x^2 + 4)^{(n-1)/2} v_{n+r} \text{ for } 2 \nmid n. \tag{5.5}$$

$$6^\circ. \sum_{i=0}^n \binom{n}{i} v_{2i+r} = (x^2 + 4)^{n/2} v_{n+r} \text{ for } 2 \mid n, \text{ or } (x^2 + r)^{(n+1)/2} u_{n+r} \text{ for } 2 \nmid n. \tag{5.6}$$

**Proof:** We prove only 2° and 5°. The rest can be proved in the same way.

$$2^\circ. \sum_{i=0}^n v_i = \sum_{i=0}^n (\alpha^i + \beta^i) = (1 - \alpha^{n+1}) / (1 - \alpha) + (1 - \beta^{n+1}) / (1 - \beta) \\ = (1 - \alpha^{n+1} - \beta - \alpha^n + 1 - \beta^{n+1} - \alpha - \beta^n) / (-x) = (v_{n+1} + v_n - 2) / x + 1.$$

5°. We have

$$\sum_{i=0}^n \binom{n}{i} \alpha^{2i} = (1 + \alpha^2)^n = (-\alpha\beta + \alpha^2)^n = \Delta^n \alpha^n.$$

For the same reason

$$\sum_{i=0}^n \binom{n}{i} \beta^{2i} = (-1)^n \Delta^n \beta^n.$$

Hence,

$$\sum_{i=0}^n \binom{n}{i} u_{2i+r} = \sum_{i=0}^n \binom{n}{i} (\alpha^{2i+r} - \beta^{2i+r}) / \Delta = \Delta^n [\alpha^{n+r} - (-1)^n \beta^{n+r}] / \Delta \\ = \Delta^n [\alpha^{n+r} - \beta^{n+r}] / \Delta = (x^2 + 4)^{n/2} u_{n+r} \quad \text{for } 2|n, \\ \text{or} \quad = \Delta^{n-1} (\alpha^{n+r} + \beta^{n+r}) = (x^2 + 4)^{(n-1)/2} v_{n+r} \quad \text{for } 2 \nmid n. \quad \square$$

**Theorem 5.1:**

$$\sum_{i=0}^n u_i^{(k)} = \sum_{i=0}^k (-1)^i \binom{k}{i} [u_{n+1}^{(k-i)} + u_n^{(k-i)} - \delta_{k,i}] / x^{i+1} \quad (x \neq 0); \quad (5.7)$$

$$\sum_{i=0}^n v_i^{(k)} = \sum_{i=0}^k (-1)^i \binom{k}{i} [v_{n+1}^{(k-i)} + v_n^{(k-i)} - 2\delta_{k,i}] / x^{i+1} \quad (x \neq 0); \quad (5.8)$$

$$\sum_{i=0}^n \binom{n}{i} x^i h_{i+r}^{(k)} = \sum_{i=0}^k (-1)^i \binom{k}{i} (n)_i h_{2n-i+r}^{(k-i)}, \quad (5.9)$$

where  $\{h_n^{(i)}\}$  is  $\{u_n^{(i)}\}$  or  $\{v_n^{(i)}\}$  ( $i = 0, \dots, k$ );

$$\sum_{i=0}^n (-1)^i \binom{n}{i} h_{2i+r}^{(k)} = (-1)^n \sum_{i=0}^k \binom{k}{i} (n)_i x^{n-i} h_{n+r}^{(k-i)}, \quad (5.10)$$

where  $\{h_n^{(i)}\}$  is  $\{u_n^{(i)}\}$  or  $\{v_n^{(i)}\}$  ( $i = 0, \dots, k$ );

$$\sum_{i=0}^n \binom{n}{i} u_{2i+r}^{(k)} = \sum_{i=0}^k \binom{k}{i} u_{n+r}^{(k-i)} \frac{d^i}{dx^i} (x^2 + 4)^{n/2} \quad \text{for } 2|n, \quad (5.11)$$

$$\text{or} \quad = \sum_{i=0}^k \binom{k}{i} v_{n+r}^{(k-i)} \frac{d^i}{dx^i} (x^2 + 4)^{(n-1)/2} \quad \text{for } 2 \nmid n;$$

$$\sum_{i=0}^n \binom{n}{i} v_{2i+r}^{(k)} = \sum_{i=0}^k \binom{k}{i} v_{n+r}^{(k-i)} \frac{d^i}{dx^i} (x^2 + 4)^{n/2} \quad \text{for } 2|n, \quad (5.12)$$

$$\text{or} \quad = \sum_{i=0}^k \binom{k}{i} u_{n+r}^{(k-i)} \frac{d^i}{dx^i} (x^2 + r)^{(n+1)/2} \quad \text{for } 2 \nmid n.$$

**Proof:** Every one of (5.7), (5.8), (5.10)-(5.12) can be proved straightforwardly by differentiating the corresponding one of (5.1), (5.2), (5.4)-(5.6). The proof of (5.9) is as follows.

Let

$$g_{n,k,r} = g_{n,k,r}(x) = \sum_{i=0}^k \binom{n}{i} x^i h_{i+r}^{(k)}. \tag{5.13}$$

Then

$$g'_{n,k,r} = \sum_{i=0}^k \binom{n}{i} x^i h_{i+r}^{(k+1)} + \sum_{i=1}^n \binom{n}{i} i x^{i-1} h_{i+r}^{(k)} = g_{n,k+1,r} + n \cdot g_{n-1,k,r+1}.$$

So

$$g_{n,k+1,r} = g'_{n,k,r} - n \cdot g_{n-1,k,r+1}. \tag{5.14}$$

When  $k = 0$ , from (5.3), we can see that (5.9) holds. Assume that (5.9) holds for  $k$ ; then from (5.14), we have

$$g_{n,k+1,r} = \sum_{i=0}^k (-1)^i \binom{k}{i} (n)_i h_{2n-i+r}^{(k+1-i)} - n \sum_{i=0}^k (-1)^i \binom{k}{i} (n-1)_i h_{2n-1-i+r}^{(k-i)}.$$

The second summation in the right side of the last expression can be rewritten as

$$\begin{aligned} & -n \sum_{i=0}^{k-1} (-1)^i \binom{k}{i} (n-1)_i h_{2n-1-i+r}^{(k-i)} - n(-1)^k \cdot (n-1)_k h_{2n-1-k+r} \\ & = \sum_{i=1}^k (-1)^i \binom{k}{i-1} (n)_i h_{2n-i+r}^{(k+1-i)} + (-1)^{k+1} (n)_{k+1} h_{2n-(k+1)+r}. \end{aligned}$$

From this, it follows that

$$g_{n,k+1,r} = \sum_{i=0}^{k+1} (-1)^i \binom{k+1}{i} (n)_i h_{2n-i+r}^{(k+1-i)},$$

that is, (5.9) also holds for  $k + 1$ , and we are done.  $\square$

It is known that the generating function of  $\{u_n^{(k)}\}$  is expressed by (2.7). It is well known that the generating function of  $\{v_n\}$  is

$$V(t) = (2 - xt) / (1 - xt - t^2). \tag{5.15}$$

Differentiating (5.15), we can know that the generating function of  $\{v_n^{(k)}\}$  is

$$V_k(t) = k! t^k (1 + t^2) / (1 - xt - t^2)^{k+1} \quad (k \geq 1). \tag{5.16}$$

Obviously, the following identities hold:

$$\begin{aligned} U_k(t) \cdot U_r(t) &= \frac{k!r!}{(k+r+1)!} U_{k+r+1}(t); \\ V_k(t) \cdot V_r(t) &= \frac{k!r!}{(k+r+1)!} (t+t^{-1}) V_{k+r+1}(t) \quad (k, r \geq 1); \\ U_k(t) \cdot V_r(t) &= \frac{k!r!}{(k+r+1)!} V_{k+r+1}(t) \quad (r \geq 1); \\ U_k(t) \cdot V(t) &= \frac{1}{k+1} (2t^{-1} - x) U_{k+1}(t); \end{aligned}$$

$$V_k(t) \cdot V(t) = \frac{1}{k+1} (2t^{-1} - x) V_{k+1}(t) \quad (k \geq 1).$$

Equalizing the coefficients of  $t^n$  of the two sides in each of the above identities, we have

**Theorem 5.2:**

$$\sum_{i=0}^n u_i^{(k)} u_{n-i}^{(r)} = \frac{k!r!}{(k+r+1)!} u_n^{(k+r+1)}; \quad (5.17)$$

$$\sum_{i=0}^n v_i^{(k)} v_{n-i}^{(r)} = \frac{k!r!}{(k+r+1)!} (v_{n-i}^{(k+r+1)} + v_{n+1}^{(k+r+1)}) \quad (k, r \geq 1); \quad (5.18)$$

$$\sum_{i=0}^n u_i^{(k)} v_{n-i}^{(r)} = \frac{k!r!}{(k+r+1)!} v_n^{(k+r+1)} \quad (r \geq 1); \quad (5.19)$$

$$\sum_{i=0}^n u_i^{(k)} v_{n-i} = \frac{1}{k+1} (2u_{n+1}^{(k+1)} - xu_n^{(k+1)}); \quad (5.20)$$

$$\sum_{i=0}^n v_i^{(k)} v_{n-i} = \frac{1}{k+1} (2v_{n+1}^{(k+1)} - xv_n^{(k+1)}) \quad (k \geq 1). \quad (5.21)$$

#### REFERENCES

1. P. Filippini & A. F. Horadam. "Derivative Sequences of Fibonacci and Lucas Polynomials." In *Applications of Fibonacci Numbers* 4:99-108. Ed. G. E. Bergum, A. N. Philippou, & A. F. Horadam. Dordrecht: Kluwer, 1991.
2. P. Filippini & A. F. Horadam. "Second Derivative Sequences of Fibonacci and Lucas Polynomials." *The Fibonacci Quarterly* 31.3 (1993):194-204.
3. P. Filippini & A. F. Horadam. "Addendum to "Second Derivative Sequences of Fibonacci and Lucas Polynomials." *The Fibonacci Quarterly* 32.2 (1994):110.
4. K. S. Miller. *Advanced Complex Calculus*. New York, 1960.
5. E. Lucas. "Théorie des fonctions numériques simplement périodiques." *Amer. J. Math.* 1 (1878):184-240, 289-321.
6. Richard A. Brualdi. *Introductory Combinatorics*. The Hague: North-Holland, 1977.
7. Y. H. Harris Kwong. "Periodicities of a Class of Infinite Integer Sequences Modulo  $m$ ." *J. Number Theory* 31 (1989):64-79.
8. V. E. Hoggatt, Jr. *Fibonacci and Lucas Numbers*. Boston: Houghton Mifflin, 1969; rpt. Santa Clara, Calif.: The Fibonacci Association, 1979.
9. Chizhong Zhou. *Fibonacci-Lucas Sequences and Their Application* (in Chinese). China: Hunan Science and Technology Publishing House, 1993. [MR 95m:11027]

AMS Classification Numbers: 11B39, 26A24, 11B83

