

Research Article

On Genocchi Numbers and Polynomials

Seog-Hoon Rim, Kyoung Ho Park, and Eun Jung Moon

Department of Mathematics, Kyungpook National University, Taegu 702-701, South Korea

Correspondence should be addressed to Kyoung Ho Park, sagamath@yahoo.co.kr

Received 10 May 2008; Accepted 21 July 2008

Recommended by Lance Littlejohn

The main purpose of this paper is to study the distribution of Genocchi polynomials. Finally, we construct the Genocchi zeta function which interpolates Genocchi polynomials at negative integers.

Copyright © 2008 Seog-Hoon Rim et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Let p be a fixed odd prime number. Throughout this paper, \mathbb{Z}_p , \mathbb{Q}_p , \mathbb{C} , and \mathbb{C}_p will, respectively, denote the ring of p -adic rational integers, the field of p -adic rational numbers, the complex number field, and the completion of the algebraic closure of \mathbb{Q}_p . Let v_p be the normalized exponential valuation of \mathbb{C}_p with $|p|_p = p^{-v_p(p)} = 1/p$. When one talks about q -extension, q is variously considered as an indeterminate, a complex, $q \in \mathbb{C}$, or a p -adic number, $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$, one normally assumes $|q| < 1$. If $q \in \mathbb{C}_p$, then we assume $|q - 1|_p < 1$. The ordinary Genocchi polynomials are defined as the generating function:

$$F(t, x) = \frac{2t}{e^t + 1} e^{xt} = \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!}, \quad |t| < \pi. \quad (1.1)$$

For a fixed positive integer d with $(p, d) = 1$, set

$$\begin{aligned} X &= X_d = \varprojlim_{\mathbb{N}} \frac{\mathbb{Z}}{dP^N \mathbb{Z}}, \quad X_1 = \mathbb{Z}_p, \\ X^* &= \bigcup_{\substack{0 < a < dp, \\ (a, p) = 1}} (a + dp\mathbb{Z}_p), \\ a + dp^N \mathbb{Z}_p &= \{x \in X \mid x \equiv a \pmod{dp^N}\} \end{aligned} \quad (1.2)$$

(cf. [1–30]), where $a \in \mathbb{Z}$ satisfies the condition $0 \leq a < dp^N$. We say that f is uniformly differential function at $a \in \mathbb{Z}_p$ and write $f \in UD(\mathbb{Z}_p)$ if the difference quotients, $F_f(x, y) = (f(x) - f(y))/(x - y)$, have a limit $f'(a)$ as $(x, y) \rightarrow (a, a)$. Throughout this paper, we use the following notation:

$$[x]_{-q} = \frac{1 - (-q)^x}{1 + q}, \quad [x]_q = \frac{1 - q^x}{1 - q}. \quad (1.3)$$

For $f \in UD(\mathbb{Z}_p)$, the fermionic p -adic invariant q -integral on \mathbb{Z}_p is defined as

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_{-q}} \sum_{x=0}^{p^N-1} f(x)(-q)^x; \quad (1.4)$$

see [1–27]. Note that

$$I_{-1}(f) = \lim_{q \rightarrow -1} I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x). \quad (1.5)$$

In this paper, we investigate some interesting integral equations related to $I_{-1}(f)$. From these integral equations related to $I_{-1}(f)$, we can derive many interesting properties of Genocchi numbers and polynomials. The main purpose of this paper is to derive the distribution relations of the Genocchi polynomials, and to construct the Genocchi zeta function which interpolates the Genocchi polynomials at negative integers.

2. Genocchi numbers and polynomials

The Genocchi numbers are defined as

$$G_0 = 0, \quad (G + 1)^n + G_n = \begin{cases} 2 & \text{if } n = 1, \\ 0 & \text{if } n > 1, \end{cases} \quad (2.1)$$

where G^n is replaced by G_n , symbolically. The Genocchi polynomials are also defined as

$$G_n(x) = \sum_{k=0}^n \binom{n}{k} x^{n-k} G_k. \quad (2.2)$$

From (2.1), we note that $G_1 = 1$, $G_2 = -1$, $G_3 = 0$, $G_4 = 1, \dots, G_{2k+1} = 0$, and $G_{2k} \in \mathbb{Z}$ ($k = 1, 2, \dots$). The fermionic p -adic invariant integral on \mathbb{Z}_p is defined as

$$I_{-1}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_{-1}} \sum_{x=0}^{p^N-1} f(x)(-1)^x, \quad \text{see [1]}. \quad (2.3)$$

Let $f_1(x)$ be translation with $f_1(x) = f(x + 1)$. Then we have the following integral equation. Note that $I_{-1}(f_1) + I_{-1}(f) = 2f(0)$. From (2.3), we can derive

$$t \int_{\mathbb{Z}_p} e^{xt} d\mu_{-1}(x) = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!} = t \sum_{n=0}^{\infty} \frac{G_{n+1}}{n+1} \frac{t^n}{n!}, \quad (2.4)$$

$$t \int_{\mathbb{Z}_p} e^{(x+y)t} d\mu_{-1}(y) = \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!} = t \sum_{n=0}^{\infty} \frac{G_{n+1}(x)}{n+1} \frac{t^n}{n!}. \quad (2.5)$$

Thus, we obtain

$$\int_{\mathbb{Z}_p} x^n d\mu_{-1}(x) = \frac{G_{n+1}}{n+1}, \quad \int_{\mathbb{Z}_p} (x+y)^n d\mu_{-1}(y) = \frac{G_{n+1}(x)}{n+1}. \quad (2.6)$$

For $n \in \mathbb{N}$, we have

$$\int_{\mathbb{Z}_p} f(x+n) d\mu_{-1}(x) = (-1)^n \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) + 2 \sum_{\ell=0}^{n-1} (-1)^{n-1+\ell} f(\ell), \quad \text{see [1-27]}. \quad (2.7)$$

By (2.6) and (2.7), if we take $f(x) = x^k$ ($k \in \mathbb{Z}^+$), we easily see that

$$\int_{\mathbb{Z}_p} (x+n)^k d\mu_{-1}(x) - \int_{\mathbb{Z}_p} x^k d\mu_{-1}(x) = 2 \sum_{\ell=0}^{n-1} (-1)^{\ell-1} \ell^k \quad \text{if } n \equiv 0 \pmod{2}. \quad (2.8)$$

Thus, we have

$$\frac{G_{k+1}(n)}{k+1} - \frac{G_{k+1}}{k+1} = 2 \sum_{\ell=0}^{n-1} (-1)^{\ell-1} \ell^k \quad \text{if } n \equiv 0 \pmod{2}. \quad (2.9)$$

If $n \equiv 1 \pmod{2}$, then we know that

$$\int_{\mathbb{Z}_p} (x+n)^k d\mu_{-1}(x) + \int_{\mathbb{Z}_p} x^k d\mu_{-1}(x) = 2 \sum_{\ell=0}^{n-1} (-1)^{\ell} \ell^k \quad \text{if } n \equiv 1 \pmod{2}. \quad (2.10)$$

Thus, we get

$$\frac{G_{k+1}(n)}{k+1} + \frac{G_{k+1}}{k+1} = 2 \sum_{\ell=0}^{n-1} (-1)^{\ell} \ell^k, \quad \text{see [1-30]}. \quad (2.11)$$

Let χ be the Dirichlet character with conductor $d \in \mathbb{N}$, with $d \equiv 1 \pmod{2}$. Then, we consider the generalized Genocchi numbers attached to χ as follows:

$$\frac{G_{n+1, \chi}}{n+1} = \int_{\mathbb{X}} \chi(x) x^n d\mu_{-1}(x), \quad G_{0, \chi} = 0, \quad (2.12)$$

where $n \in \mathbb{Z}_+$. From (2.7) and (2.12), we note that

$$t \int_{\mathbb{X}} e^{xt} \chi(x) d\mu_{-1}(x) = \frac{2 \sum_{\ell=0}^{d-1} (-1)^{\ell} \chi(\ell) e^{\ell t}}{e^{dt} + 1} t = \sum_{n=0}^{\infty} \frac{G_{n, \chi}}{n!} t^n. \quad (2.13)$$

By (2.12) and (2.13), it is not difficult to show that

$$\begin{aligned} \frac{G_{n+1, \chi}}{n+1} &= \int_{\mathbb{X}} \chi(x) x^n d\mu_{-1}(x) \\ &= d^n \sum_{a=0}^{d-1} \chi(a) (-1)^a \int_{\mathbb{Z}_p} \left(\frac{a}{d} + x \right)^n d\mu_{-1}(x) \\ &= d^n \sum_{a=0}^{d-1} \chi(a) (-1)^a \frac{G_{n+1}(a/d)}{n+1}, \end{aligned} \quad (2.14)$$

$$\int_{\mathbb{X}} (x+y)^n d\mu_{-1}(y) = d^n \sum_{a=0}^{d-1} (-1)^a \int_{\mathbb{Z}_p} \left(\frac{x+a}{d} + y \right)^n d\mu_{-1}(y). \quad (2.15)$$

By (2.6) and (2.15), we obtain the following theorem.

Theorem 2.1. *Let $d \in \mathbb{N}$ with $d \equiv 1 \pmod{2}$, and let χ be the Dirichlet character with conductor d . Then, one has*

$$\frac{G_{n+1}(x)}{n+1} = d^n \sum_{a=0}^{d-1} (-1)^a \frac{G_{n+1}((x+a)/d)}{n+1} \quad (\text{distribution relation for Genocchi polynomials}), \quad (2.16)$$

$$\frac{G_{n+1,\chi}}{n+1} = d^n \sum_{a=0}^{d-1} \chi(a) (-1)^a \frac{G_{n+1}(a/d)}{n+1}. \quad (2.17)$$

3. Genocchi zeta function

Let $F(t, x)$ be the generating function of $G_k(x)$ in complex plane as follows:

$$\begin{aligned} F(t, x) &= \frac{2te^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!} \\ &= t \sum_{n=0}^{\infty} \frac{G_{n+1}(x)}{n+1} \frac{t^n}{n!}, \quad |t| < \pi. \end{aligned} \quad (3.1)$$

Then, we show that

$$F(t, x) = 2t \sum_{n=0}^{\infty} (-1)^n e^{(n+x)t}. \quad (3.2)$$

By (3.1) and (3.2), we easily see that

$$G_k(x) = \left. \frac{d^k F(t, x)}{dt^k} \right|_{t=0} = 2k \sum_{n=0}^{\infty} (-1)^n (n+x)^{k-1}. \quad (3.3)$$

Therefore, we obtain the following proposition.

Proposition 3.1. *For $k \in \mathbb{N}$, one has*

$$\frac{G_k(x)}{k} = 2 \sum_{n=0}^{\infty} (-1)^n (n+x)^{k-1}. \quad (3.4)$$

From Proposition 3.1, we can derive the Genocchi zeta function which interpolates Genocchi polynomials at negative integers.

For $s \in \mathbb{C}$, we define the Hurwitz-type Genocchi zeta function as follows.

Definition 3.2. For $s \in \mathbb{C}$,

$$\zeta_G(s, x) = 2 \sum_{n=0}^{\infty} \frac{(-1)^n}{(n+x)^s}, \quad \zeta_G(s) = 2 \sum_{n=1}^{\infty} \frac{(-1)^n}{n^s}. \quad (3.5)$$

By Proposition 3.1 and Definition 3.2, we obtain the following theorem.

Theorem 3.3. For $k \in \mathbb{N}$, one has

$$\zeta_G(1-k, x) = \frac{G_k(x)}{k}, \quad \zeta_G(1-k) = \frac{G_k}{k}. \quad (3.6)$$

Let χ be the Dirichlet character with conductor $d \in \mathbb{N}$, with $d \equiv 1 \pmod{2}$, and let $F_\chi(t)$ be the generating function in \mathbb{C} of $G_{n,\chi}$. Then, we have

$$F_\chi(t) = 2 \frac{\sum_{\ell=0}^{d-1} (-1)^\ell \chi(\ell) e^{\ell t}}{e^{dt} + 1} t = \sum_{n=0}^{\infty} G_{n,\chi} \frac{t^n}{n!}, \quad |t| < \frac{\pi}{d}. \quad (3.7)$$

From (3.7), we derive

$$\begin{aligned} F_\chi(t) &= \sum_{n=0}^{\infty} G_{n,\chi} \frac{t^n}{n!} = \sum_{n=1}^{\infty} G_{n,\chi} \frac{t^n}{n!} \\ &= t \sum_{n=0}^{\infty} \frac{G_{n+1,\chi}}{n+1} \frac{t^n}{n!} \\ &= t \sum_{n=0}^{\infty} \left(d^n \sum_{a=0}^{d-1} \chi(a) (-1)^a \frac{G_{n+1}(a/d)}{n+1} \right) \frac{t^n}{n!} \\ &= \sum_{a=0}^{d-1} \chi(a) (-1)^a \left(dt \sum_{n=0}^{\infty} \frac{G_{n+1}(a/d)}{n+1} \frac{d^n t^n}{n!} \right). \end{aligned} \quad (3.8)$$

By (3.1), (3.2), and (3.8), we easily see that

$$\begin{aligned} F_\chi(t) &= \sum_{a=0}^{d-1} \chi(a) (-1)^a \left(2t \sum_{k=0}^{\infty} (-1)^k e^{(k+a/d)t} \right) \\ &= 2t \sum_{k=0}^{\infty} \sum_{a=0}^{d-1} \chi(a+dk) (-1)^{a+dk} e^{(dk+a)t} \\ &= 2t \sum_{n=0}^{\infty} \chi(n) (-1)^n e^{nt} \\ &= 2t \sum_{n=1}^{\infty} \chi(n) (-1)^n e^{nt}. \end{aligned} \quad (3.9)$$

From (3.9), we can derive

$$G_{k,\chi} = \frac{d^k}{dt^k} F_\chi(t) \Big|_{t=0} = k \left(2 \sum_{n=1}^{\infty} \chi(n) (-1)^n n^{k-1} \right). \quad (3.10)$$

Thus, we have

$$\frac{G_{k,\chi}}{k} = 2 \sum_{n=1}^{\infty} \chi(n) (-1)^n n^{k-1}. \quad (3.11)$$

Now, we consider the Dirichlet-type Genocchi ℓ -function in complex plane as follows. For $s \in \mathbb{C}$, define

$$\ell_{G,\chi}(s) = 2 \sum_{n=1}^{\infty} \frac{(-1)^n \chi(n)}{n^s}. \quad (3.12)$$

By (3.11) and (3.12), we obtain the following theorem.

Theorem 3.4. *Let χ be the Dirichlet character with conductor $d \in \mathbb{N}$, with $d \equiv 1 \pmod{2}$, and let $k \in \mathbb{Z}^+$. Then, one has*

$$\ell_{G,\chi}(1-k) = \frac{G_{k,\chi}}{k}. \quad (3.13)$$



Remark 3.5. In [1], we can observe the value of Genocchi zeta function at positive integers as follows:

$$\zeta_G(2n) = \frac{(-1)^{n-1} \pi^{2n} (2-4^n)}{2(2n)!(1-4^n)} G_{2n}, \quad \text{cf. [1]}. \quad (3.14)$$

References

- [1] T. Kim, "Euler numbers and polynomials associated with zeta functions," *Abstract and Applied Analysis*, vol. 2008, Article ID 581582, 11 pages, 2008.
- [2] T. Kim, "The modified q -Euler numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 16, no. 2, pp. 161–170, 2008.
- [3] T. Kim, "A note on p -adic q -integral on \mathbb{Z}_p associated with q -Euler numbers," *Advanced Studies in Contemporary Mathematics*, vol. 15, no. 2, pp. 133–137, 2007.
- [4] T. Kim, " q -Volkenborn integration," *Russian Journal of Mathematical Physics*, vol. 9, no. 3, pp. 288–299, 2002.
- [5] T. Kim and Y. Simsek, "Analytic continuation of the multiple Daehee q - l -functions associated with Daehee numbers," *Russian Journal of Mathematical Physics*, vol. 15, no. 1, pp. 58–65, 2008.
- [6] T. Kim, " q -Bernoulli numbers and polynomials associated with Gaussian binomial coefficients," *Russian Journal of Mathematical Physics*, vol. 15, no. 1, pp. 51–57, 2008.
- [7] T. Kim, " q -extension of the Euler formula and trigonometric functions," *Russian Journal of Mathematical Physics*, vol. 14, no. 3, pp. 275–278, 2007.
- [8] T. Kim, J. Y. Choi, and J. Y. Sug, "Extended q -Euler numbers and polynomials associated with fermionic p -adic q -integral on \mathbb{Z}_p ," *Russian Journal of Mathematical Physics*, vol. 14, no. 2, pp. 160–163, 2007.
- [9] S.-H. Rim and T. Kim, "A note on p -adic Euler measure on \mathbb{Z}_p ," *Russian Journal of Mathematical Physics*, vol. 13, no. 3, pp. 358–361, 2006.
- [10] T. Kim, " q -generalized Euler numbers and polynomials," *Russian Journal of Mathematical Physics*, vol. 13, no. 3, pp. 293–298, 2006.
- [11] T. Kim, "Multiple p -adic L -function," *Russian Journal of Mathematical Physics*, vol. 13, no. 2, pp. 151–157, 2006.
- [12] H. M. Srivastava, T. Kim, and Y. Simsek, " q -Bernoulli numbers and polynomials associated with multiple q -zeta functions and basic L -series," *Russian Journal of Mathematical Physics*, vol. 12, no. 2, pp. 241–268, 2005.
- [13] T. Kim, "Power series and asymptotic series associated with the q -analog of the two-variable p -adic L -function," *Russian Journal of Mathematical Physics*, vol. 12, no. 2, pp. 186–196, 2005.
- [14] T. Kim, "Analytic continuation of multiple q -zeta functions and their values at negative integers," *Russian Journal of Mathematical Physics*, vol. 11, no. 1, pp. 71–76, 2004.
- [15] T. Kim, "Non-Archimedean q -integrals associated with multiple Changhee q -Bernoulli polynomials," *Russian Journal of Mathematical Physics*, vol. 10, no. 1, pp. 91–98, 2003.
- [16] T. Kim, "On Euler-Barnes multiple zeta functions," *Russian Journal of Mathematical Physics*, vol. 10, no. 3, pp. 261–267, 2003.

- [17] T. Kim, L.-C. Jang, and H. K. Pak, "A note on q -Euler and Genocchi numbers," *Proceedings of the Japan Academy. Series A*, vol. 77, no. 8, pp. 139–141, 2001.
- [18] T. Kim, "A note on q -Volkenborn integration," *Proceedings of the Jangjeon Mathematical Society*, vol. 8, no. 1, pp. 13–17, 2005.
- [19] T. Kim, "On the q -extension of Euler and Genocchi numbers," *Journal of Mathematical Analysis and Applications*, vol. 326, no. 2, pp. 1458–1465, 2007.
- [20] T. Kim, "A note on the q -Genocchi numbers and polynomials," *Journal of Inequalities and Applications*, vol. 2007, Article ID 71452, 8 pages, 2007.
- [21] T. Kim, et al., *Introduction to Non-Archimedean Analysis*, Seoul, Korea, Kyo Woo Sa, 2004.
- [22] L.-C. Jang and T. Kim, " q -Genocchi numbers and polynomials associated with fermionic p -adic invariant integrals on \mathbb{Z}_p ," *Abstract and Applied Analysis*, vol. 2008, Article ID 232187, 8 pages, 2008.
- [23] L.-C. Jang, T. Kim, D.-H. Lee, and D.-W. Park, "An application of polylogarithms in the analogs of Genocchi numbers," *Notes on Number Theory and Discrete Mathematics*, vol. 7, no. 3, pp. 65–69, 2001.
- [24] M. Cenkci, M. Can, and V. Kurt, " p -adic interpolation functions and Kummer-type congruences for q -twisted and q -generalized twisted Euler numbers," *Advanced Studies in Contemporary Mathematics*, vol. 9, no. 2, pp. 203–216, 2004.
- [25] H. Ozden, Y. Simsek, S.-H. Rim, and I. N. Cangul, "A note on p -adic q -Euler measure," *Advanced Studies in Contemporary Mathematics*, vol. 14, no. 2, pp. 233–239, 2007.
- [26] Y. Simsek, "On p -adic twisted q - L -functions related to generalized twisted Bernoulli numbers," *Russian Journal of Mathematical Physics*, vol. 13, no. 3, pp. 340–348, 2006.
- [27] Y. Simsek, I. N. Cangul, V. Kurt, and D. Kim, " q -Genocchi numbers and polynomials associated with q -Genocchi-type l -functions," *Advances in Difference Equations*, vol. 2008, Article ID 815750, 12 pages, 2008.
- [28] M. Schork, "Combinatorial aspects of normal ordering and its connection to q -calculus," *Advanced Studies in Contemporary Mathematics*, vol. 15, no. 1, pp. 49–57, 2007.
- [29] M. Schork, "Ward's "calculus of sequences", q -calculus and the limit $q \rightarrow -1$," *Advanced Studies in Contemporary Mathematics*, vol. 13, no. 2, pp. 131–141, 2006.
- [30] K. Shiratani and S. Yamamoto, "On a p -adic interpolation function for the Euler numbers and its derivatives," *Memoirs of the Faculty of Science. Kyushu University. Series A*, vol. 39, no. 1, pp. 113–125, 1985.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

