

RESEARCH

Open Access

Some identities on Bernoulli and Euler polynomials arising from the orthogonality of Laguerre polynomials

Taekyun Kim^{1*}, Seog-Hoon Rim², DV Dolgy³ and Sang-Hun Lee⁴

*Correspondence: tkkim@kw.ac.kr
¹Department of Mathematics,
Kwangwoon University, Seoul,
139-701, Republic of Korea
Full list of author information is
available at the end of the article

Abstract

In this paper, we derive some interesting identities on Bernoulli and Euler polynomials by using the orthogonal property of Laguerre polynomials.

1 Introduction

The generalized Laguerre polynomials are defined by

$$\frac{\exp(-\frac{xt}{1-t})}{(1-t)^{\alpha+1}} = \sum_{n=0}^{\infty} L_n^{\alpha}(x) t^n \quad (\alpha \in \mathbb{Q} \text{ with } \alpha > -1). \quad (1.1)$$

From (1.1), we note that

$$L_n^{\alpha}(x) = \sum_{r=0}^n \frac{(-1)^r \binom{n+\alpha}{n-r} x^r}{r!} \quad (\text{see [1-3]}). \quad (1.2)$$

By (1.2), we see that $L_n^{\alpha}(x)$ is a polynomial with degree n . It is well known that Rodrigues' formula for $L_n^{\alpha}(x)$ is given by

$$L_n^{\alpha}(x) = \frac{x^{-\alpha} e^x}{n!} \left(\frac{d^n}{dx^n} (e^{-x} x^{n+\alpha}) \right) \quad (\text{see [1-3]}). \quad (1.3)$$

From (1.3) and a part of integration, we note that

$$\int_0^{\infty} x^{\alpha} e^{-x} L_m^{\alpha}(x) L_n^{\alpha}(x) dx = \frac{\Gamma(\alpha + n + 1)}{n!} \delta_{m,n}, \quad (1.4)$$

where $\delta_{m,n}$ is a Kronecker symbol. As is well known, Bernoulli polynomials are defined by the generating function to be

$$\frac{t}{e^t - 1} e^{xt} = e^{B(x)t} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!} \quad (\text{see [1-29]}), \quad (1.5)$$

with the usual convention about replacing $B^n(x)$ by $B_n(x)$.

In the special case, $x = 0$, $B_n(0) = B_n$ are called the n th Bernoulli numbers. By (1.5), we get

$$B_n(x) = \sum_{l=0}^n \binom{n}{l} B_{n-l} x^l \quad (\text{see [1–29]}). \quad (1.6)$$

The Euler numbers are defined by

$$E_0 = 1, \quad (E + 1)^n + E_n = 2\delta_{0,n} \quad (\text{see [27, 28]}), \quad (1.7)$$

with the usual convention about replacing E^n by E_n .

In the viewpoint of (1.6), the Euler polynomials are also defined by

$$E_n(x) = (E + x)^n = \sum_{l=0}^n \binom{n}{l} E_{n-l} x^l \quad (\text{see [11–24]}). \quad (1.8)$$

From (1.7) and (1.8), we note that the generating function of the Euler polynomial is given by

$$\frac{2}{e^t + 1} e^{xt} = e^{E(x)t} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!} \quad (\text{see [15–29]}). \quad (1.9)$$

By (1.5) and (1.9), we get

$$\frac{2}{e^t + 1} e^{xt} = \frac{1}{t} \left(2 - 2 \frac{2}{e^t + 1} \right) \left(\frac{te^{xt}}{e^t - 1} \right) = -2 \sum_{n=0}^{\infty} \left(\sum_{l=0}^n \frac{E_{l+1}}{l+1} \binom{n}{l} B_{n-l}(x) \right) \frac{t^n}{n!}. \quad (1.10)$$

Thus, by (1.10), we see that

$$E_n(x) = -2 \sum_{l=0}^n \binom{n}{l} \frac{E_{l+1}}{l+1} B_{n-l}(x). \quad (1.11)$$

By (1.7) and (1.8), we easily get

$$\frac{t}{e^t - 1} e^{xt} = \frac{t}{2} \left(\frac{2e^{xt}}{e^t + 1} \right) + \left(\frac{t}{e^t - 1} \right) \left(\frac{2e^{xt}}{e^t + 1} \right). \quad (1.12)$$

Thus, by (1.12), we see that

$$B_n(x) = \sum_{k=0, k \neq 1}^n \binom{n}{k} B_k E_{n-k}(x). \quad (1.13)$$

Throughout this paper, we assume that $\alpha \in \mathbb{Q}$ with $\alpha > -1$. Let $\mathbb{P}_n = \{p(x) \in \mathbb{Q}[x] \mid \deg p(x) \leq n\}$ be the inner product space with the inner product

$$\langle p(x), q(x) \rangle = \int_0^\infty x^\alpha e^{-x} p(x) q(x) dx,$$

where $p(x), q(x) \in \mathbb{P}_n$. From (1.4), we note that $\{L_0^\alpha(x), L_1^\alpha(x), \dots, L_n^\alpha(x)\}$ is an orthogonal basis for \mathbb{P}_n .

In this paper, we give some interesting identities on Bernoulli and Euler polynomials which can be derived by an orthogonal basis $\{L_0^\alpha(x), L_1^\alpha(x), \dots, L_n^\alpha(x)\}$ for \mathbb{P}_n .

2 Some identities on Bernoulli and Euler polynomials

Let $p(x) \in \mathbb{P}_n$. Then $p(x)$ can be generated by $\{L_0^\alpha(x), L_1^\alpha(x), \dots, L_n^\alpha(x)\}$ in \mathbb{P}_n to be

$$p(x) = \sum_{k=0}^n C_k L_k^\alpha(x), \quad (2.1)$$

where

$$\begin{aligned} \langle p(x), L_k^\alpha(x) \rangle &= C_k \langle L_k^\alpha(x), L_k^\alpha(x) \rangle \\ &= C_k \int_0^\infty x^\alpha e^{-x} L_k^\alpha(x) L_k^\alpha(x) dx \\ &= C_k \frac{\Gamma(\alpha + k + 1)}{k!}. \end{aligned} \quad (2.2)$$

From (2.2), we note that

$$\begin{aligned} C_k &= \frac{k!}{\Gamma(\alpha + k + 1)} \langle p(x), L_k^\alpha(x) \rangle \\ &= \frac{k!}{\Gamma(\alpha + k + 1)} \frac{1}{k!} \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) p(x) dx \\ &= \frac{1}{\Gamma(\alpha + k + 1)} \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) p(x) dx. \end{aligned} \quad (2.3)$$

Let us take $p(x) = \sum_{m=0, m \neq 1}^n \binom{n}{m} B_m E_n - m(x) \in \mathbb{P}_n$. Then, from (2.3), we have

$$\begin{aligned} C_k &= \frac{1}{\Gamma(\alpha + k + 1)} \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) \sum_{m=0, m \neq 1}^n \binom{n}{m} B_m E_{n-m}(x) dx \\ &= \frac{(-1)^k}{\Gamma(\alpha + k + 1)} \sum_{m=0, m \neq 1}^{n-k} \sum_{l=k}^{n-m} \binom{n}{m} \binom{n-m}{l} B_m E_{n-m-l} \frac{l!}{(l-k)!} \int_0^\infty x^{l+\alpha} e^{-x} dx \\ &= \frac{(-1)^k}{\Gamma(\alpha + k + 1)} \sum_{m=0, m \neq 1}^{n-k} \sum_{l=k}^{n-m} \binom{n}{m} \binom{n-m}{l} B_m E_{n-m-l} \frac{l!}{(l-k)!} \Gamma(l + \alpha + 1) \\ &= (-1)^k \sum_{m=0, m \neq 1}^{n-k} \sum_{l=k}^{n-m} \binom{n}{m} \binom{n-m}{l} B_m E_{n-m-l} \frac{l!}{(l-k)!} \frac{(l+\alpha)(l+\alpha-1)\cdots\alpha}{(\alpha+k)(\alpha+k-1)\cdots\alpha} \\ &= (-1)^k n! \sum_{m=0, m \neq 1}^{n-k} \sum_{l=k}^{n-m} \frac{B_m}{m!} \frac{E_{n-m-l}}{(n-m-l)!} \binom{l+\alpha}{l-k}. \end{aligned} \quad (2.4)$$

Therefore, by (2.1) and (2.4), we obtain the following theorem.

Theorem 2.1 For $n \in \mathbb{Z}_+$, we have

$$\begin{aligned} & \sum_{m=0, m \neq 1}^n \binom{n}{m} B_m E_{n-m}(x) \\ &= n! \sum_{k=0}^{n-k} (-1)^k \left(\sum_{m=0, m \neq 1}^n \sum_{l=k}^{n-m} \frac{B_m}{m!} \frac{E_n - m - l}{(n-m-l)!} \binom{l+\alpha}{l-k} \right) L_k^\alpha(x). \end{aligned}$$

From (1.13), we can derive the following corollary.

Corollary 2.2 For $n \in \mathbb{Z}_+$, we have

$$B_n(x) = n! \sum_{k=0}^n (-1)^k \left(\sum_{m=0, m \neq 1}^{n-k} \sum_{l=k}^{n-m} \frac{B_m}{m!} \frac{E_n - m - l}{(n-m-l)!} \binom{l+\alpha}{l-k} \right) L_k^\alpha(x).$$

Let us take $p(x) = \sum_{l=0}^n \binom{n}{l} \frac{E_{l+1}}{l+1} B_{n-l}(x)$. By the same method, we get

$$\begin{aligned} C_k &= \frac{1}{\Gamma(\alpha + k + 1)} \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) \sum_{l=0}^n \binom{n}{l} \frac{E_{l+1}}{l+1} B_{n-l}(x) dx \\ &= \frac{1}{\Gamma(\alpha + k + 1)} \sum_{l=0}^{n-k} \sum_{m=0}^{n-l} \binom{n}{l} \binom{n-l}{m} \frac{E_{l+1}}{l+1} B_{n-l-m} \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) x^m dx \\ &= \frac{(-1)^k}{\Gamma(\alpha + k + 1)} \sum_{l=0}^{n-k} \sum_{m=k}^{n-l} \binom{n}{l} \binom{n-l}{m} \frac{E_{l+1}}{l+1} B_{n-l-m} \frac{m!}{(m-k)!} \Gamma(m + \alpha + 1) \\ &= (-1)^k \sum_{l=0}^{n-k} \sum_{m=k}^{n-l} \binom{n}{l} \binom{n-l}{m} \frac{m!}{(m-k)!} \frac{E_{l+1}}{(l+1)!} B_{n-l-m} \frac{(\alpha+m)(\alpha+m-1)\cdots\alpha}{(\alpha+k)(\alpha+k-1)\cdots\alpha} \\ &= (-1)^k n! \sum_{l=0}^{n-k} \sum_{m=k}^{n-l} \binom{\alpha+m}{m-k} \frac{E_{l+1}}{(l+1)!} \frac{B_{n-l-m}}{(n-l-m)!}. \end{aligned} \tag{2.5}$$

Therefore, by (1.11), (2.1), and (2.5), we obtain the following theorem.

Theorem 2.3 For $n \in \mathbb{Z}_+$, we have

$$\frac{E_n(x)}{2} = n! \sum_{k=0}^n (-1)^k \left(\sum_{l=0}^{n-k} \sum_{m=k}^{n-l} \binom{\alpha+m}{m-k} \frac{E_{m+1}}{(m+1)!} \frac{B_{n-m-l}}{(n-m-l)!} \right) L_k^\alpha(x).$$

For $n \in \mathbb{N}$ with $n \geq 2$ and $m \in \mathbb{Z}_+$ with $n - m \geq 0$, we have

$$\begin{aligned} B_{n-m}(x) B_m(x) &= \sum_r \left\{ \binom{n-m}{2r} m + \binom{m}{2r} (n-m) \right\} \frac{B_{2r} B_{n-2r}(x)}{n-2r} \\ &\quad + (-1)^{m+1} \frac{(n-m)! m!}{n!} B_n \in \mathbb{P}_n \quad (\text{see [8]}). \end{aligned} \tag{2.6}$$

Let us take $p(x) = B_{n-m}(x)B_m(x) \in \mathbb{P}_n$. Then $p(x)$ can be generated by an orthogonal basis $\{L_0^\alpha(x), L_1^\alpha(x), \dots, L_n^\alpha(x)\}$ in \mathbb{P}_n to be

$$p(x) = \sum_{k=0}^n C_k L_k^\alpha(x). \quad (2.7)$$

From (2.3), (2.6), and (2.7), we note that

$$\begin{aligned} C_k &= \frac{1}{\Gamma(\alpha + k + 1)} \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) p(x) dx \\ &= \frac{1}{\Gamma(\alpha + k + 1)} \sum_{r=0}^{\lfloor \frac{n}{2} \rfloor} \left\{ \binom{n-m}{2r} m + \binom{m}{2r} (n-m) \right\} \\ &\quad \times \frac{B_{2r}}{n-2r} \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) B_{n-2r}(x) dx \\ &= \frac{1}{\Gamma(\alpha + k + 1)} \sum_{r=0}^{\lfloor \frac{n}{2} \rfloor} \left\{ \binom{n-m}{2r} m + \binom{m}{2r} (n-m) \right\} \frac{B_{2r}}{n-2r} \\ &\quad \times \sum_{l=0}^{n-2r} \binom{n-2r}{l} B_{n-2r-l} \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) x^l dx \\ &= \frac{1}{\Gamma(\alpha + k + 1)} \sum_{r=0}^{\lfloor \frac{n}{2} \rfloor} \sum_{l=0}^{n-2r} \left\{ \binom{n-m}{2r} m + \binom{m}{2r} (n-m) \right\} \binom{n-2r}{l} \\ &\quad \times \frac{B_{2r} B_{n-2r-l}}{n-2r} \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) x^l dx \\ &= \frac{(-1)^k}{\Gamma(\alpha + k + 1)} \sum_{r=0}^{\lfloor \frac{n}{2} \rfloor} \sum_{l=k}^{n-2r} \left\{ \binom{n-m}{2r} m + \binom{m}{2r} (n-m) \right\} \binom{n-2r}{l} \\ &\quad \times \frac{B_{2r} B_{n-2r-l} l!}{(n-2r)(l-k)!} \Gamma(\alpha + l + 1). \end{aligned} \quad (2.8)$$

It is easy to show that

$$\begin{aligned} \frac{\Gamma(\alpha + l + 1)}{\Gamma(\alpha + k + 1)(l-k)!} &= \frac{(\alpha + l)(\alpha + l - 1) \cdots \alpha \Gamma(\alpha)}{(\alpha + k)(\alpha + k - 1) \cdots \alpha \Gamma(\alpha)(l-k)!} \\ &= \frac{(\alpha + l)(\alpha + l - 1) \cdots (\alpha + k + 1)}{(\alpha - k)!} = \binom{\alpha + l}{l - k}. \end{aligned} \quad (2.9)$$

By (2.8) and (2.9), we get

$$\begin{aligned} C_k &= (-1)^k \sum_{r=0}^{\lfloor \frac{n}{2} \rfloor} \sum_{l=k}^{n-2r} \left\{ \binom{n-m}{2r} m + \binom{m}{2r} (n-m) \right\} \\ &\quad \times \binom{n-2r}{l} \binom{\alpha + l}{l - k} \frac{l! B_{2r} B_{n-2r-l}}{(n-2r)}. \end{aligned} \quad (2.10)$$

Therefore, by (2.7) and (2.10), we obtain the following theorem.

Theorem 2.4 For $n \in \mathbb{N}$ with $n \geq 2$ and $m \in \mathbb{Z}_+$ with $n - m \geq 0$, we have

$$B_{n-m}(x)B_m(x) = \sum_{k=0}^n (-1)^k \left\{ \sum_{r=0}^{\lfloor \frac{n-k}{2} \rfloor} \sum_{l=k}^{n-2r} \left(\binom{n-m}{2r} m + \binom{m}{2r} (n-m) \right) \right. \\ \times \left. \binom{n-2r}{l} \binom{\alpha+l}{l-k} \frac{l! B_{2r} B_{n-2r-l}}{(n-2r)!} \right\} L_k^\alpha(x).$$

It is easy to show that

$$\frac{t^2 e^{t(x+y)}}{(e^t - 1)^2} = (x+y-1) \frac{t^2 e^{t(x+y-1)}}{e^t - 1} - \frac{t^2 d}{dt} \left(\frac{e^{t(x+y-1)}}{e^t - 1} \right). \quad (2.11)$$

From (2.11), we have

$$\sum_{k=0}^n \binom{n}{k} B_k(x) B_{n-k}(y) = (1-n) B_n(x+y) + (x+y-1) n B_{n-1}(x+y) \quad (\text{see [11]}). \quad (2.12)$$

Let $x = y$. Then by (2.12), we get

$$\sum_{k=0}^n \binom{n}{k} B_k(x) B_{n-k}(x) = (1-n) B_n(2x) + (2x-1) B_{n-1}(2x). \quad (2.13)$$

Let us take $p(x) = \sum_{k=0}^n \binom{n}{k} B_k(x) B_{n-k}(x) \in \mathbb{P}_n$. Then $p(x)$ can be generated by an orthogonal basis $\{L_0^\alpha(x), L_1^\alpha(x), \dots, L_n^\alpha(x)\}$ in \mathbb{P}_n to be

$$p(x) = \sum_{k=0}^n \binom{n}{k} B_k(x) B_{n-k}(x) = \sum_{k=0}^n C_k L_k^\alpha(x). \quad (2.14)$$

From (2.3), (2.13), and (2.14), we can determine the coefficients C_k 's to be

$$C_k = \frac{1}{\Gamma(\alpha+k+1)} \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) p(x) dx \\ = \frac{1}{\Gamma(\alpha+k+1)} \left\{ (1-n) \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) B_n(2x) dx \right. \\ \left. + n \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) (2x-1) B_{n-1}(2x) dx \right\}. \quad (2.15)$$

By simple calculation, we get

$$\frac{1}{\Gamma(\alpha+k+1)} \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) (2x-1) B_{n-1}(2x) dx \\ = 2(-1)^k \sum_{l=k-1}^{n-1} \binom{n-1}{l} 2^l B_{n-1-l} \binom{\alpha+l+1}{l-k+1} (l+1)! \\ + (-1)^{k+1} \sum_{l=k}^{n-1} \binom{n-1}{l} 2^l B_{n-1-l} \binom{\alpha+l}{l-k} l! \quad (2.16)$$

and

$$\begin{aligned} & \frac{1}{\Gamma(\alpha + k + 1)} \int_0^\infty \left(\frac{d^k}{dx^k} x^{k+\alpha} e^{-x} \right) B_n(2x) dx \\ &= (-1)^k \sum_{l=k}^n \binom{n}{l} 2^l B_{n-l} l! \binom{l+\alpha}{l-k}. \end{aligned} \quad (2.17)$$

Therefore, by (2.13), (2.14), (2.15), (2.16), and (2.17), we obtain the following theorem.

Theorem 2.5 For $n \in \mathbb{Z}_+$, we get

$$\begin{aligned} & \sum_{k=0}^n \binom{n}{k} B_k(x) B_{n-k}(x) \\ &= (1-n) \sum_{k=0}^n \left\{ (-1)^k \sum_{l=k}^n \binom{n}{l} 2^l B_{n-l} l! \binom{l+\alpha}{l-k} \right\} L_k^\alpha(x) \\ &+ n \sum_{k=0}^n (-1)^k \left\{ \sum_{l=k-1}^{n-1} \binom{n-1}{l} 2^{l+1} B_{n-1-l} (l+1)! \binom{\alpha+l+1}{l-k+1} \right. \\ &\quad \left. - \sum_{l=k}^{n-1} \binom{n-1}{l} 2^l B_{n-l-1} l! \binom{\alpha+l}{l-k} \right\} L_k^\alpha(x). \end{aligned}$$

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to the manuscript and typed, read, and approved the final manuscript.

Author details

¹Department of Mathematics, Kwangwoon University, Seoul, 139-701, Republic of Korea. ²Department of Mathematics Education, Kyungpook National University, Taegu, 702-701, Republic of Korea. ³Hanrimwon, Kwangwoon University, Seoul, 139-701, Republic of Korea. ⁴Division of General Education, Kwangwoon University, Seoul, 139-701, Republic of Korea.

Acknowledgements

The authors would like to express their deep gratitude to the referees for their valuable suggestions and comments.

Received: 8 August 2012 Accepted: 6 November 2012 Published: 22 November 2012

References

1. Carlitz, L: An integral for the product of two Laguerre polynomials. *Boll. Unione Mat. Ital.* **17**(17), 25-28 (1962)
2. Carlitz, L: On the product of two Laguerre polynomials. *J. Lond. Math. Soc.* **36**, 399-402 (1961)
3. Cangul, N, Kurt, V, Ozden, H, Simsek, Y: On the higher-order w - q -Genocchi numbers. *Adv. Stud. Contemp. Math.* **19**(1), 39-57 (2009)
4. Akemann, G, Kieburg, M, Phillips, MJ: Skew-orthogonal Laguerre polynomials for chiral real asymmetric random matrices. *J. Phys. A* **43**(37), Art. ID 375207 (2010)
5. Bayad, A, Kim, T: Identities for the Bernoulli, the Euler and the Genocchi numbers and polynomials. *Adv. Stud. Contemp. Math.* **20**(2), 247-253 (2010)
6. Bayad, A: Modular properties of elliptic Bernoulli and Euler functions. *Adv. Stud. Contemp. Math.* **20**(3), 389-401 (2010)
7. Carlitz, L: Note on the integral of the product of several Bernoulli polynomials. *J. Lond. Math. Soc.* **34**, 361-363 (1959)
8. Carlitz, L: Some generating functions for Laguerre polynomials. *Duke Math. J.* **35**, 825-827 (1968)
9. Costin, RD: Orthogonality of Jacobi and Laguerre polynomials for general parameters via the Hadamard finite part. *J. Approx. Theory* **162**(1), 141-152 (2010)
10. Ding, D, Yang, J: Some identities related to the Apostol-Euler and Apostol-Bernoulli polynomials. *Adv. Stud. Contemp. Math.* **20**(1), 7-21 (2010)
11. Hansen, ER: A Table of Series and Products. Prentice Hall, Englewood Cliffs (1975)

12. Kudo, A: A congruence of generalized Bernoulli number for the character of the first kind. *Adv. Stud. Contemp. Math.* **2**, 1-8 (2000)
13. Kim, T, Choi, J, Kim, YH, Ryoo, CS: On q -Bernstein and q -Hermite polynomials. *Proc. Jangjeon Math. Soc.* **14**(2), 215-221 (2011)
14. Kim, T: A note on q -Bernstein polynomials. *Russ. J. Math. Phys.* **18**(1), 73-82 (2011)
15. Kim, T: Some identities on the q -Euler polynomials of higher order and q -Stirling numbers by the fermionic p -adic integral on \mathbb{Z}_p . *Russ. J. Math. Phys.* **16**(4), 484-491 (2009)
16. Kim, T: Symmetry of power sum polynomials and multivariate fermionic p -adic invariant integral on \mathbb{Z}_p . *Russ. J. Math. Phys.* **16**(1), 93-96 (2009)
17. Ozden, H, Cangul, IN, Simsek, Y: Multivariate interpolation functions of higher-order q -Euler numbers and their applications. *Abstr. Appl. Anal.* **2008**, Art. ID 390857 (2008)
18. Choi, J, Kim, DS, Kim, T, Kim, YH: Some arithmetic identities on Bernoulli and Euler numbers arising from the p -adic integrals on \mathbb{Z}_p . *Adv. Stud. Contemp. Math.* **22**(2), 239-247 (2012)
19. Ozden, H, Cangul, IN, Simsek, Y: Remarks on q -Bernoulli numbers associated with Daehee numbers. *Adv. Stud. Contemp. Math.* **18**(1), 41-48 (2009)
20. Rim, SH, Lee, SJ: Some identities on the twisted (h, q) -Genocchi numbers and polynomials associated with q -Bernstein polynomials. *Int. J. Math. Math. Sci.* **2011**, Art. ID 482840 (2011)
21. Rim, SH, Bayad, A, Moon, EJ, Jin, JH, Lee, SJ: A new construction on the q -Bernoulli polynomials. *Adv. Differ. Equ.* **2011**, 34 (2011)
22. Rim, SH, Jin, JH, Moon, EJ, Lee, SJ: Some identities on the q -Genocchi polynomials of higher-order and q -Stirling numbers by the fermionic p -adic integral on \mathbb{Z}_p . *Int. J. Math. Math. Sci.* **2010**, Art. ID 860280 (2010)
23. Ryoo, CS: On the generalized Barnes type multiple q -Euler polynomials twisted by ramified roots of unity. *Proc. Jangjeon Math. Soc.* **13**(2), 255-263 (2010)
24. Ryoo, CS: A note on the Frobenius-Euler polynomials. *Proc. Jangjeon Math. Soc.* **14**(4), 495-501 (2011)
25. Ryoo, CS: Some relations between twisted q -Euler numbers and Bernstein polynomials. *Adv. Stud. Contemp. Math.* **21**(2), 217-223 (2011)
26. Ryoo, CS: Some identities of the twisted q -Euler numbers and polynomials associated with q -Bernstein polynomials. *Proc. Jangjeon Math. Soc.* **14**(2), 239-248 (2011)
27. Simsek, Y: Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions. *Adv. Stud. Contemp. Math. (Kyungshang)* **16**(2), 251-278 (2008)
28. Simsek, Y: Special functions related to Dedekind-type DC-sums and their applications. *Russ. J. Math. Phys.* **17**(4), 495-508 (2010)
29. Simsek, Y: On p -adic twisted q - L -functions related to generalized twisted Bernoulli numbers. *Russ. J. Math. Phys.* **13**(3), 340-348 (2006)

doi:10.1186/1687-1847-2012-201

Cite this article as: Kim et al.: Some identities on Bernoulli and Euler polynomials arising from the orthogonality of Laguerre polynomials. *Advances in Difference Equations* 2012 2012:201.

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com