



Generalized Bivariate Lucas p -Polynomials and Hessenberg Matrices

Kenan Kaygisiz and Adem Şahin

Department of Mathematics

Faculty of Arts and Sciences

Gaziosmanpaşa University

60250 Tokat

Turkey

kenan.kaygisiz@gop.edu.tr

adem.sahin@gop.edu.tr

Abstract

In this paper, we give some determinantal and permanental representations of generalized bivariate Lucas p -polynomials by using various Hessenberg matrices. The results that we obtained are important since generalized bivariate Lucas p -polynomials are general forms of, for example, bivariate Jacobsthal-Lucas, bivariate Pell-Lucas p -polynomials, Chebyshev polynomials of the first kind, Jacobsthal-Lucas numbers etc.

1 Introduction

The generalized Lucas p -numbers [15] are defined by

$$L_p(n) = L_p(n-1) + L_p(n-p-1) \quad (1)$$

for $n > p+1$, with boundary conditions $L_p(0) = (p+1)$, $L_p(1) = \dots = L_p(p) = 1$.

The Lucas [8], Pell-Lucas [2] and Chebyshev polynomials of the first kind [17] are defined as follows:

$$\begin{aligned} l_{n+1}(x) &= xl_n(x) + l_{n-1}(x), \quad n \geq 2 \text{ with } l_0(x) = 2, \quad l_1(x) = x \\ Q_{n+1}(x) &= 2xQ_n(x) + Q_{n-1}(x), \quad n \geq 2 \text{ with } Q_0(x) = 2, \quad Q_1(x) = 2x \\ T_{n+1}(x) &= 2xT_n(x) - T_{n-1}(x), \quad n \geq 2 \text{ with } T_0(x) = 1, \quad T_1(x) = x \end{aligned}$$

respectively.

The generalized bivariate Lucas p -polynomials [16] are defined as follows:

$$L_{p,n}(x, y) = xL_{p,n-1}(x, y) + yL_{p,n-p-1}(x, y)$$

for $n > p$, with boundary conditions $L_{p,0}(x, y) = (p + 1)$, $L_{p,n}(x, y) = x^n$, $n = 1, 2, \dots, p$.

A few terms of $L_{p,n}(x, y)$ for $p = 4$ and $p = 5$ are

$5, x, x^2, x^3, x^4, 5y + x^5, 6xy + x^6, x^7 + 7x^2y, x^8 + 8x^3y, x^9 + 9x^4y, 5y^2 + x^{10} + 10x^5y, \dots$ and $6, x, x^2, x^3, x^4, x^5, 5y + x^6, 6xy + x^7, x^8 + 7x^2y, x^9 + 8x^3y, \dots$ respectively.

MacHenry [9] defined generalized Lucas polynomials ($L_{k,n}(t)$) where t_i ($1 \leq i \leq k$) are constant coefficients of the core polynomial

$$P(x; t_1, t_2, \dots, t_k) = x^k - t_1x^{k-1} - \dots - t_k,$$

which is denoted by the vector $t = (t_1, t_2, \dots, t_k)$.

$G_{k,n}(t_1, t_2, \dots, t_k)$ is defined by

$$\begin{aligned} G_{k,n}(t) &= 0, \quad n < 0 \\ G_{k,0}(t) &= k \\ G_{k,1}(t) &= t_1 \\ G_{k,n+1}(t) &= t_1G_{k,n}(t) + \dots + t_kG_{k,n-k+1}(t). \end{aligned}$$

MacHenry obtained very useful properties of these polynomials in [10, 11].

Remark 1. [16]Cognate polynomial sequence are as follows

\mathbf{x}	\mathbf{y}	\mathbf{p}	$L_{p,n}(x, y)$
x	y	1	bivariate Lucas polynomials $L_n(x, y)$
x	1	p	Lucas p -polynomials $L_{p,n}(x)$
x	1	1	Lucas polynomials $l_n(x)$
1	1	p	Lucas p -numbers $L_p(n)$
1	1	1	Lucas numbers L_n
$2x$	y	p	bivariate Pell-Lucas p -polynomials $L_{p,n}(2x, y)$
$2x$	y	1	bivariate Pell-Lucas polynomials $L_n(2x, y)$
$2x$	1	p	Pell-Lucas p -polynomials $Q_{p,n}(x)$
$2x$	1	1	Pell-Lucas polynomials $Q_n(x)$
2	1	1	Pell-Lucas numbers Q_n
$2x$	-1	1	Chebyshev polynomials of the first kind $T_n(x)$
x	$2y$	p	bivariate Jacobsthal-Lucas p -polynomials $L_{p,n}(x, 2y)$
x	$2y$	1	Bivariate Jacobsthal-Lucas polynomials $L_n(x, 2y)$
1	$2y$	1	Jacobsthal-Lucas polynomials $j_n(y)$
1	2	1	Jacobsthal-Lucas numbers j_n

Remark 1 shows that $L_{p,n}(x, y)$ is a general form of all sequences and polynomials mentioned in that remark. Therefore, any result obtained from $L_{p,n}(x, y)$ is valid for all sequences and polynomials mentioned there.

Many researchers have studied determinantal and permanental representations of k sequences of generalized order- k Fibonacci and Lucas numbers. For example, Minc [12] defined

an $n \times n$ $(0,1)$ -matrix $F(n, k)$, and showed that the permanents of $F(n, k)$ are equal to the generalized order- k Fibonacci numbers. Nalli and Haukkanen [13] defined $h(x)$ -Fibonacci and Lucas polynomials and gave determinantal representations of these polynomials. The authors ([6, 7]) defined two $(0, 1)$ -matrices and showed that the permanents of these matrices are the generalized Fibonacci and Lucas numbers. Öcal et al. [14] gave some determinantal and permanental representations of k -generalized Fibonacci and Lucas numbers, and obtained Binet's formula for these sequences. Kılıc and Stakhov [4] gave permanent representation of Fibonacci and Lucas p -numbers. Kılıc and Tasci [5] studied permanents and determinants of Hessenberg matrices. Janjic [3] considers a particular case of upper Hessenberg matrices and gave a determinant representation of a generalized Fibonacci numbers.

In this paper, we give some determinantal and permanental representations of $L_{p,n}(x, y)$ by using various Hessenberg matrices. These results are a general form of determinantal and permanental representations of polynomials and sequences mentioned in Remark 1.

2 The determinantal representations

In this section, we give some determinantal representations of $L_{p,n}(x, y)$ using Hessenberg matrices.

Definition 2. An $n \times n$ matrix $A_n = (a_{ij})$ is called lower Hessenberg matrix if $a_{ij} = 0$ when $j - i > 1$ i.e.,

$$A_n = \begin{bmatrix} a_{11} & a_{12} & 0 & \cdots & 0 \\ a_{21} & a_{22} & a_{23} & \cdots & 0 \\ a_{31} & a_{32} & a_{33} & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ a_{n-1,1} & a_{n-1,2} & a_{n-1,3} & \cdots & a_{n-1,n} \\ a_{n,1} & a_{n,2} & a_{n,3} & \cdots & a_{n,n} \end{bmatrix}.$$

Theorem 3. [1] Let A_n be an $n \times n$ lower Hessenberg matrix for all $n \geq 1$ and $\det(A_0) = 1$. Then,

$$\det(A_1) = a_{11}$$

and for $n \geq 2$

$$\det(A_n) = a_{n,n} \det(A_{n-1}) + \sum_{r=1}^{n-1} \left[(-1)^{n-r} a_{n,r} \left(\prod_{j=r}^{n-1} a_{j,j+1} \right) \det(A_{r-1}) \right].$$

Theorem 4. Let $L_{p,n}(x, y)$ be the generalized bivariate Lucas p -polynomials and $W_{p,n} = (w_{ij})$ be an $n \times n$ Hessenberg matrix defined by

$$w_{ij} = \begin{cases} i, & \text{if } i = j - 1; \\ x, & \text{if } i = j; \\ i^p y, & \text{if } p = i - j \text{ and } j \neq 1; \\ (p + 1)i^p y, & \text{if } p = i - j \text{ and } j = 1; \\ 0, & \text{otherwise;} \end{cases}$$

that is,

$$W_{p,n} = \begin{bmatrix} x & i & 0 & \cdots & 0 \\ 0 & x & i & \ddots & \vdots \\ \vdots & 0 & x & & 0 \\ (p+1)i^p y & 0 & \vdots & \cdots & \\ 0 & i^p y & 0 & & 0 \\ \vdots & 0 & \ddots & x & i \\ 0 & 0 & \cdots & 0 & x \end{bmatrix}. \quad (2)$$

Then,

$$\det(W_{p,n}) = L_{p,n}(x, y) \quad (3)$$

where $n \geq 1$ and $i = \sqrt{-1}$.

Proof. To prove (3), we use mathematical induction on n . The result is true for $n = 1$ by hypothesis.

Assume that it is true for all positive integers less than or equal to n , namely $\det(W_{p,n}) = L_{p,n}(x, y)$. Then, we have

$$\begin{aligned} \det(W_{p,n+1}) &= q_{n+1,n+1} \det(W_{p,n}) + \sum_{r=1}^n \left[(-1)^{n+1-r} q_{n+1,r} \left(\prod_{j=r}^n q_{j,j+1} \right) \det(W_{p,r-1}) \right] \\ &= x \det(W_{p,n}) + \sum_{r=1}^{n-p} \left[(-1)^{n+1-r} q_{n+1,r} \left(\prod_{j=r}^n q_{j,j+1} \right) \det(W_{p,r-1}) \right] \\ &\quad + \sum_{r=n-p+1}^n \left[(-1)^{n+1-r} q_{n+1,r} \left(\prod_{j=r}^n q_{j,j+1} \right) \det(W_{p,r-1}) \right] \\ &= x \det(W_{p,n}) + \left[(-1)^p (i)^p y \prod_{j=n-p+1}^n i \det(W_{p,n-p}) \right] \\ &= x \det(W_{p,n}) + [(-1)^p y (i)^p \cdot (i)^p \det(W_{p,n-p})] \\ &= x \det(W_{p,n}) + y \det(W_{p,n-p}) \end{aligned}$$

by using Theorem 3. From the induction hypothesis and the definition of $L_{p,n}(x, y)$ we obtain

$$\det(W_{p,n+1}) = xL_{p,n}(x, y) + yL_{p,n-p}(x, y) = L_{p,n+1}(x, y).$$

Therefore, (3) holds for all positive integers n . \square

Example 5. We obtain the 5-th $L_{p,n}(x, y)$ for $p = 4$, by using Theorem 4

$$L_{4,5}(x, y) = \det \begin{bmatrix} x & i & 0 & 0 & 0 \\ 0 & x & i & 0 & 0 \\ 0 & 0 & x & i & 0 \\ 0 & 0 & 0 & x & i \\ 5i^4 y & 0 & 0 & 0 & x \end{bmatrix} = 5y + x^5.$$

Theorem 6. Let $p \geq 1$ be an integer, $L_{p,n}(x, y)$ be the generalized bivariate Lucas p -polynomials and $M_{p,n} = (m_{ij})$ be an $n \times n$ Hessenberg matrix defined by

$$m_{ij} = \begin{cases} -1, & \text{if } j = i + 1; \\ x, & \text{if } i = j; \\ y, & \text{if } p = i - j \text{ and } j \neq 1; \\ (p + 1)y, & \text{if } p = i - j \text{ and } j = 1; \\ 0, & \text{otherwise;} \end{cases}$$

that is,

$$M_{p,n} = \begin{bmatrix} x & -1 & 0 & \cdots & 0 \\ 0 & x & -1 & \cdots & 0 \\ 0 & 0 & x & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ (p+1)y & 0 & 0 & \cdots & 0 \\ 0 & y & 0 & \cdots & 0 \\ & \vdots & \vdots & \ddots & -1 \\ 0 & 0 & \cdots & 0 & x \end{bmatrix}. \quad (4)$$

Then,

$$\det(M_{p,n}) = L_{p,n}(x, y).$$

Proof. Since the proof is similar to the proof of Theorem 4, we omit the details. \square

3 The permanent representations

Let $A = (a_{i,j})$ be a square matrix of order n over a ring R . The permanent of A is defined by

$$\text{per}(A) = \sum_{\sigma \in S_n} \prod_{i=1}^n a_{i,\sigma(i)}$$

where S_n denotes the symmetric group on n letters.

Theorem 7. [14] Let A_n be an $n \times n$ lower Hessenberg matrix for all $n \geq 1$ and $\text{per}(A_0) = 1$. Then, $\text{per}(A_1) = a_{11}$ and for $n \geq 2$,

$$\text{per}(A_n) = a_{n,n} \text{per}(A_{n-1}) + \sum_{r=1}^{n-1} \left[a_{n,r} \left(\prod_{j=r}^{n-1} a_{j,j+1} \right) \text{per}(A_{r-1}) \right].$$

Theorem 8. Let $p \geq 1$ be an integer, $L_{p,n}(x, y)$ be the generalized bivariate Lucas p -polynomials and $H_{p,n} = (h_{rs})$ be an $n \times n$ lower Hessenberg matrix such that

$$h_{rs} = \begin{cases} -i, & \text{if } s - r = 1 ; \\ x, & \text{if } r = s ; \\ i^p y, & \text{if } p = r - s \text{ and } s \neq 1, ; \\ (p + 1)i^p y, & \text{if } p = r - s \text{ and } s = 1; \\ 0, & \text{otherwise;} \end{cases}$$

then

$$\text{per}(H_{p,n}) = L_{p,n}(x, y)$$

where $n \geq 1$ and $i = \sqrt{-1}$.

Proof. This is similar to the proof of Theorem 4 using Theorem 7. □

Example 9. We obtain the 6-th $L_{p,n}(x, y)$ for $p = 4$, by using Theorem 8

$$L_{4,6}(x, y) = \text{per} \begin{bmatrix} x & -i & 0 & 0 & 0 & 0 \\ 0 & x & -i & 0 & 0 & 0 \\ 0 & 0 & x & -i & 0 & 0 \\ 0 & 0 & 0 & x & -i & 0 \\ 5y & 0 & 0 & 0 & x & -i \\ 0 & y & 0 & 0 & 0 & x \end{bmatrix} = 6xy + x^6.$$

Theorem 10. Let $p \geq 1$ be an integer, $L_{p,n}(x, y)$ be the generalized bivariate Lucas p -polynomials and $K_{p,n} = (k_{ij})$ be an $n \times n$ lower Hessenberg matrix such that

$$k_{ij} = \begin{cases} 1, & \text{if } j = i + 1; \\ x, & \text{if } i = j; \\ y, & \text{if } i - j = p \text{ and } j \neq 1; \\ (p + 1)y, & \text{if } i - j = p \text{ and } j = 1; \\ 0, & \text{otherwise;} \end{cases}$$

then

$$\text{per}(K_{p,n}) = L_{p,n}(x, y).$$

Proof. This is similar to the proof of Theorem 4 by using Theorem 7. □

We note that the theorems given above are still valid for the sequences and polynomials mentioned in Remark 1

Corollary 11. *If we rewrite Theorem 4, Theorem 6, Theorem 8 and Theorem 10 for x, y, p , we obtain the following table.*

For	x	y	p	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{L}_{p,n+1}(\mathbf{x}, \mathbf{y}),$
for	x	y	1	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{L}_n(\mathbf{x}, \mathbf{y}),$
for	x	1	p	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{L}_{p,n}(\mathbf{x}),$
for	x	1	1	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = l_n(x),$
for	1	1	p	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{L}_p(\mathbf{n}),$
for	1	1	1	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{L}_n,$
for	$2x$	y	p	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{L}_{p,n}(2\mathbf{x}, \mathbf{y}),$
for	$2x$	y	1	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{L}_n(2\mathbf{x}, \mathbf{y}),$
for	$2x$	1	p	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{Q}_{p,n}(\mathbf{x}),$
for	$2x$	1	1	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{Q}_n(\mathbf{x}),$
for	2	1	1	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{Q}_n,$
for	$2x$	-1	1	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{T}_n(\mathbf{x}),$
for	x	$2y$	p	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{L}_{p,n}(\mathbf{x}, 2\mathbf{y}),$
for	x	$2y$	1	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{L}_n(\mathbf{x}, 2\mathbf{y}),$
for	1	$2y$	1	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{j}_n(\mathbf{y}),$
for	1	2	1	$\det(W_{p,n}) = \det(M_{p,n}) = \text{per}(H_{p,n}) = \text{per}(K_{p,n}) = \mathbf{j}_n.$

4 Conclusion

In this paper, we have given some determinantal and permanental representations of generalized bivariate Lucas p -polynomials. Our results allow us to derive determinantal and permanental representations of sequences and polynomials mentioned in Remark 1.

References

- [1] N. D. Cahill, J. R. D’Errico, D. A. Narayan, and J. Y. Narayan, Fibonacci determinants, *College Math. J.* **33** (2002), 221–225.
- [2] A. F. Horadam and J. M. Mahon, Pell and Pell-Lucas polynomials, *Fibonacci Quart.* **23** (1985), 17–20.
- [3] M. Janjic, Hessenberg matrices and integer sequences, *J. Integer Seq.* **13** (2010), Article 10.7.8.
- [4] E. Kılıç and A. P. Stakhov, On the Fibonacci and Lucas p -numbers, their sums, families of bipartite graphs and permanents of certain matrices, *Chaos Solitons Fractals* **40** (2009), 2210–2221.
- [5] E. Kılıç and D. Tasci, On the generalized Fibonacci and Pell sequences by Hessenberg matrices, *Ars Combin.* **94** (2010), 161–174.
- [6] G.-Y. Lee and S.-G. Lee, A note on generalized Fibonacci numbers, *Fibonacci Quart.* **33** (1995), 273–278.

- [7] G.-Y. Lee, k -Lucas numbers and associated bipartite graphs, *Linear Algebra Appl.* **320** (2000), 51–61.
- [8] A. Lupas, A guide of Fibonacci and Lucas polynomials, *Octagon Mathematics Magazine* **7** (1999), 2–12.
- [9] T. MacHenry, A subgroup of the group of units in the ring of arithmetic functions, *Rocky Mountain J. Math.* **29** (1999), 1055–1065.
- [10] T. MacHenry, Generalized Fibonacci and Lucas polynomials and multiplicative arithmetic functions, *Fibonacci Quart.* **38** (2000), 17–24.
- [11] T. MacHenry and K. Wong, Degree k linear recursions mod(p) and number fields. *Rocky Mountain J. Math.* **41** (2011), 1303–1327.
- [12] H. Minc, *Encyclopaedia of Mathematics and its Applications, Permanents, Vol.6*, Addison-Wesley Publishing Company, 1978.
- [13] A. Nalli and P. Haukkanen, On generalized Fibonacci and Lucas polynomials. *Chaos Solitons Fractals* **42** (2009), 3179–3186.
- [14] A. A. Öcal, N. Tuglu, and E. Altinisik, On the representation of k -generalized Fibonacci and Lucas numbers. *Appl. Math. Comput.* **170** (2005) 584–596.
- [15] A. P. Stakhov, *Introduction to Algorithmic Measurement Theory*, Publishing House “Soviet Radio”, Moscow, 1977. In Russian.
- [16] N. Tuglu, E. G. Kocer, and A. Stakhov, Bivariate Fibonacci like p -polynomials. *Appl. Math. Comput.* **217** (2011), 10239–10246.
- [17] G. Udrea, Chebshev polynomials and some methods of approximation, *Port. Math.*, Fasc. 3, **55** (1998), 261–269.

2010 *Mathematics Subject Classification*: Primary 11B37, Secondary 15A15, 15A51.

Keywords: Generalized bivariate Lucas p -polynomials, determinant, permanent and Hessenberg matrix.

Received November 30 2011; revised version received February 21 2012. Published in *Journal of Integer Sequences*, March 11 2012.

Return to [Journal of Integer Sequences home page](#).