

CARLITZ GENERALIZATIONS OF LUCAS AND LEHMER SEQUENCES

A. G. Shannon and R. S. Melham

University of Technology, Sydney, NSW, 2007, Australia
(Submitted May 1991)

1. INTRODUCTION

A Lucas fundamental sequence $\{u_n\}_{n=0}^{\infty}$ is a nondegenerate binary recurrence sequence with initial conditions $u_0 = 1$, $u_1 = P$ which satisfies the homogeneous second-order linear recurrence relation

$$(1.1) \quad u_n = Pu_{n-1} - Qu_{n-2}, \quad n \geq 2,$$

where P and Q are integers [12].

If the associated auxiliary equation

$$(1.2) \quad x^2 - Px + Q = 0$$

has roots α , β , then

$$(1.3) \quad u_n = (\alpha^{n+1} - \beta^{n+1}) / (\alpha - \beta).$$

The Fibonacci, Mersenne, and Fermat numbers are all types of Lucas numbers. Their properties were studied extensively by Carmichael [5].

Many authors have generalized aspects of them by various alterations to the characteristic equations. Some of these may be found in Dickinson [6], Feinberg [7], Harris & Styles [8], Horadam [10], Miles [14], Raab [15], Williams [19], and Zeitlin [20]. Atanassov et al. [1] have coupled the recurrence relations in their generalizations.

Lehmer [11] generalized the results of Lucas on the divisibility properties of Lucas numbers to numbers

$$(1.4) \quad \ell_n = \begin{cases} (\alpha^n - \beta^n) / (\alpha - \beta), & \text{for } n \text{ odd} \\ (\alpha^n - \beta^n) / (\alpha^2 - \beta^2), & \text{for } n \text{ even.} \end{cases}$$

It is a generalization of these numbers that we wish to consider in this paper. It is of interest to note in passing that McDaniel has also recently studied analogies between the Lucas and Lehmer sequences [13].

2. DEFINITIONS

Following Carlitz [4], we define

$$(2.1) \quad f_n^{(r)} = (\alpha^{nk+k} - \beta^{nk+k}) / (\alpha^k - \beta^k)$$

and

$$(2.2) \quad g_n^{(r)} = (\alpha^{n+k} - \beta^{n+k}) / (\alpha^k - \beta^k)$$

which are not necessarily integers, where $k = r - 1$, and α and β are the roots of (1.2) as before. For example,

$$f_n^{(2)} = (\alpha^{n+1} - \beta^{n+1}) / (\alpha - \beta) = g_n^{(2)} = u_n$$

so that these numbers are generalizations of the Lucas numbers. They are also generalizations of the Lehmer numbers if we let

$$(2.3) \quad \ell_n = \begin{cases} f_{n-1}^{(2)}, & \text{for } n \text{ odd,} \\ g_{n-2}^{(3)}, & \text{for } n \text{ even,} \end{cases}$$

Carlitz [4] first defined the $f_n^{(r)}$ in another context and proved that

$$f_n^{(r)} = \text{tr}(A_{n+1}^r)$$

where

$$A_{n+1} = \left[\begin{pmatrix} t & \\ & n-s \end{pmatrix} \right] \quad (t, s = 0, 1, \dots, n)$$

is a matrix of order $n + 1$.

Examples of $f_n^{(r)}$ and $g_n^{(r)}$ now follow. When $P = -Q = 1$ and $r = 2, 3, 4$, in turn, we have:

n	0	1	2	3	4	5	6	...
$f_n^{(2)}$	1	1	2	3	5	8	13	...
$f_n^{(3)}$	1	3	8	21	55	144	377	...
$f_n^{(4)}$	1	4	17	72	305	1292	5473	...
$g_n^{(2)}$	1	1	2	3	5	8	13	...
$g_n^{(3)}$	1	2	3	5	8	13	21	...
$g_n^{(4)}$	1	$\frac{3}{2}$	$\frac{5}{2}$	4	$\frac{13}{2}$	$\frac{21}{2}$	17	...

It can be seen from this table and the recurrence relations that other properties for these sequences could be developed by treating them as cases of Horadam's $Py_{k-1} + P^2y_{k-3} = v_k \cdot \{w_n\}$ [10].

3. RECURRENCE RELATIONS

We also need the Lucas primordial sequence $\{v_n\}_{n=0}^\infty$ defined by the recurrence relation (1.1) with initial terms $v_0 = 2$ and $v_1 = P$, so that the general term is given by

$$(3.1) \quad v_n = \alpha^n + \beta^n.$$

We can show that

$$(3.2) \quad f_{n+1}^{(r)} = v_{r-1} f_n^{(r)} - Q^{r-1} f_{n-1}^{(r)}$$

and

$$(3.3) \quad g_{n+1}^{(r)} = v_1 g_n^{(r)} - Q g_{n-1}^{(r)}.$$

The latter is the same as (1.1) when $v = 2$ since $v_1 = P$.

Proof of (3.2):

$$\begin{aligned} v_{r-1}f_n^{(r)} - Q^{r-1}f_{n-1}^{(r)} &= \left((\alpha^k + \beta^k)(\alpha^{nk+k} - \beta^{nk+k}) - (\alpha\beta)^k(\alpha^{nk} - \beta^{nk}) \right) / (\alpha^k - \beta^k) \\ &= \left(\alpha^{nk+2k} - \beta^{nk+2k} + (\alpha\beta)^k(\alpha^{nk} - \beta^{nk}) - (\alpha\beta)^k(\alpha^{nk} - \beta^{nk}) \right) / (\alpha^k - \beta^k) \\ &= (\alpha^{nk+2k} - \beta^{nk+2k}) / (\alpha^k - \beta^k) \\ &= f_{n+1}^{(r)}, \text{ as required.} \end{aligned}$$

Proof of (3.3):

$$\begin{aligned} v_1g_n^{(r)} - Qg_{n-1}^{(r)} &= \left((\alpha + \beta)(\alpha^{n+k} - \beta^{n+k}) - (\alpha\beta)(\alpha^{n+k-1} - \beta^{n+k-1}) \right) / (\alpha^k - \beta^k) \\ &= \left(\alpha^{n+k+1} - \beta^{n+k+1} + (\alpha\beta)(\alpha^{n+k-1} - \beta^{n+k-1}) - (\alpha\beta)(\alpha^{n+k-1} - \beta^{n+k-1}) \right) / (\alpha^k - \beta^k) \\ &= (\alpha^{n+k+1} - \beta^{n+k+1}) / (\alpha^k - \beta^k) \\ &= g_{n+1}^{(r)}, \text{ as required.} \end{aligned}$$

Thus, the ordinary generating functions will be given (formally) by

$$(3.4) \quad \sum_{n=0}^{\infty} f_n^{(r)} x^n = 1 / (1 - v_{r-1}x + Q^{r-1}x^2),$$

and

$$(3.5) \quad \sum_{n=0}^{\infty} g_n^{(r)} x^n = \left(1 - Q \left(\frac{u_{r-3}}{u_{r-2}} \right) x \right) / (1 - v_1x + Qx^2).$$

Proof of (3.4):

$$\begin{aligned} (1 - v_{r-1}x + Q^{r-1}x^2) \sum_{n=0}^{\infty} f_n^{(r)} x^n &= f_0^{(r)} + (f_1^{(r)} - f_0^{(r)}v_{r-1})x \quad [\text{by (3.2)}] \\ &= 1 + \left(\frac{\alpha^{2k} - \beta^{2k}}{\alpha^k - \beta^k} - (\alpha^k + \beta^k) \right) x \quad [\text{by (2.1)}] \\ &= 1. \end{aligned}$$

Proof of (3.5):

$$\begin{aligned} (1 - v_1x + Qx^2) \sum_{n=0}^{\infty} g_n^{(r)} x^n &= g_0^{(r)} + (g_1^{(r)} - g_0^{(r)}v_1)x \\ &= 1 + \left(\frac{\alpha^{k+1} - \beta^{k+1}}{\alpha^k - \beta^k} - (\alpha + \beta) \right) x \\ &= 1 - Q \left(\frac{u_{k-2}}{u_{k-1}} \right) x, \text{ as required.} \end{aligned}$$

The $g_n^{(r)}$ are related to the $f_n^{(r)}$ by

$$(3.6) \quad g_{kn}^{(r)} = f_n^{(r)},$$

and to the Lucas primordial numbers v_n by

$$(3.7) \quad v_n = P g_{n-1}^{(r)} - Q v_{r-2} g_{n-k-1}^{(r)}$$

and

$$(3.8) \quad v_n = g_{2n-k}^{(r)} / g_{n-k}^{(r)}.$$

Proof of (3.7):

$$\begin{aligned} P g_n^{(r)} - Q v_{r-2} g_{n-k}^{(r)} &= ((\alpha + \beta)(\alpha^{n+k} - \beta^{n+k}) - (\alpha\beta)(\alpha^{k-1} + \beta^{k-1})(\alpha^n - \beta^n)) / (\alpha^k - \beta^k) \\ &= (\alpha^{n+k+1} - \beta^{n+k+1} + \alpha^k \beta^{n+1} - \alpha^{n+1} \beta^k) / (\alpha^k - \beta^k) \\ &= (\alpha^{n+1}(\alpha^k - \beta^k) + \beta^{n+1}(\alpha^k - \beta^k)) / (\alpha^k - \beta^k) \\ &= v_{n+1}, \text{ as required.} \end{aligned}$$

Proof of (3.8):

$$\begin{aligned} g_{n-k}^{(r)} v_n &= (\alpha^n - \beta^n)(\alpha^n + \beta^n) / (\alpha^k - \beta^k) \\ &= (\alpha^{2n} - \beta^{2n}) / (\alpha^k - \beta^k) \\ &= g_{2n-k}^{(r)}, \text{ as required.} \end{aligned}$$

4. GENERALIZATIONS OF BARAKAT'S RESULTS

As an analog of Simson's relation, we have

$$(g_n^{(r)})^2 - g_{n-k}^{(r)} g_{n+k}^{(r)} = Q^n.$$

Proof: The numerator of the left-hand side reduces to

$$\begin{aligned} (\alpha\beta)^n \alpha^{2k} + (\alpha\beta)^n \beta^{2k} - 2(\alpha\beta)^n (\alpha\beta)^k &= (\alpha\beta)^n (\alpha^k - \beta^k)^2 \\ &= Q^n (\alpha^k - \beta^k)^2 \end{aligned}$$

which is Q^n times the denominator of the left-hand side.

When $P = -Q$, we are able to relate the $f_n^{(r)}$ to the ordinary Lucas fundamental numbers, u_n , by means of a generalization of a result of Barakat [2] for the ordinary Lucas fundamental numbers. Barakat proved

$$\begin{aligned} u_n &= \sum_{0 \leq m \leq n} \binom{n-m}{m} P^{n-2m} (-Q)^m \\ &= \sum_{0 \leq m \leq n} \binom{n-m}{m} P^{n-m} \text{ when } P = -Q. \end{aligned}$$

We define

$$x_n = u_n / P = (\alpha^{n+1} - \beta^{n+1}) / (\alpha^2 - \beta^2),$$

and, for notational convenience, set $y_n = x_{n+1}$. Thus, from Simson's relation, which can be expressed as $u_k^2 - u_{k-1} u_{k+1} = (-P)^k$, we have

$$(4.2) \quad x_k y_{k-1} - x_{k-1} y_k = (-P)^{k-2}.$$

Proof: If we divide the left-hand side of Simson's relation by P^2 , we get

$$x_k y_{k-1} - x_{k-1} y_k = (-P)^{k-2}$$

from which the result follows.

A variation of equation (3.7) is then

$$(4.3) \quad P y_{k-1} + P^2 y_{k-3} = v_k.$$

Proof: The numerator of the left-hand side is

$$\begin{aligned} & (\alpha + \beta)(\alpha^{k+1} - \beta^{k+1}) + (-\alpha\beta)(\alpha + \beta)(\alpha^{k-1} - \beta^{k-1}) \\ &= \alpha^{k+2} - \beta^{k+2} - \alpha\beta^{k+1} + \alpha^{k+1}\beta - \alpha^{k-1}\beta + \alpha^2\beta^k - \alpha^k\beta^2 + \alpha\beta^{k+1} \\ &= (\alpha^k + \beta^k)(\alpha^2 - \beta^2) \\ &= v_k \cdot (\alpha^2 - \beta^2) \quad \text{which gives the result (4.3).} \end{aligned}$$

We are now in a position to assert a property which relates these generalized Fibonacci numbers to ordinary Fibonacci numbers and at the same time gives an iterative formula for the general term. This formula generalizes Barakat [2] and Shannon [16].

$$f_n^{(r)} = \sum_{0 \leq m+s \leq n} \binom{m}{s} \binom{n-m}{s} u_{k-2}^{m-s} u_{k-1}^{2s} u_k^{n-m-s} P^m.$$

Proof:

$$\begin{aligned} \sum_{n=0}^{\infty} f_n^{(r)} z^n &= (1 - v_k z + (-P)^k z^2)^{-1} && \text{[from (3.4)]} \\ &= (1 - (P^2 y_{k-3} + P y_{k-1})z + (-P)^k z^2)^{-1} && \text{[from (4.3)]} \\ &= (1 - (P^2 x_{k-2} + P y_{k-1})z + P^3 (x_{k-2} y_{k-1} - x_{k-1} y_{k-2})z^2)^{-1} && \text{[from (4.2)]} \\ &= ((1 - P^2 x_{k-2} z)(1 - P y_{k-1} z) - P^3 x_{k-1} y_{k-2} z^2)^{-1} \\ &= \sum_{s=0}^{\infty} (1 - P^2 x_{k-2} z)^{-s-1} (1 - P y_{k-1} z)^{-s-1} x_{k-1}^s y_{k-2}^s P^{3s} z^{2s} \\ &= \sum_{s=0}^{\infty} \sum_{m=0}^{\infty} \binom{m+s}{s} (1 - P y_{k-1} z)^{-s-1} x_{k-1}^s x_{k-2}^m y_{k-2}^s P^{3s+2m} z^{2s+m} \\ &= \sum_{m=0}^{\infty} \sum_{s=0}^m \binom{m}{s} (1 - P y_{k-1} z)^{-s-1} x_{k-1}^s x_{k-2}^{m-s} y_{k-2}^s P^{s+2m} z^{m+s} \\ &= \sum_{m=0}^{\infty} \sum_{s=0}^m \sum_{t=0}^{\infty} \binom{m}{s} \binom{t+s}{s} x_{k-1}^s x_{k-2}^{m-s} y_{k-1}^t y_{k-2}^s P^{s+2m+t} z^{m+s+t} \\ &= \sum_{n=0}^{\infty} \sum_{0 \leq m+s \leq n} \binom{m}{s} \binom{n-m}{s} x_{k-1}^s x_{k-2}^{m-s} y_{k-1}^{n-m-s} y_{k-2}^s P^{n+m} z^n. \end{aligned}$$

So by equating coefficients of z^n we find

$$f_n^{(r)} = \sum_{0 \leq m+s \leq n} \binom{m}{s} \binom{n-m}{s} \frac{u_{k-1}^{2s} u_{k-2}^{m-s} u_k^{n-m-s} P^{n+m}}{P^{s+m-s+n-m-s+s}}$$

$$= \sum_{0 \leq m+s \leq n} \binom{m}{s} \binom{n-m}{s} u_{k-2}^{m-s} u_{k-1}^{2s} u_k^{n-m-s} P^m, \text{ as required.}$$

For example, when $r = 2$ (and so $k = 1$), since $f_{-1}^{(2)} = 0$,

$$f_n^{(2)} = \sum_{0 \leq 2m \leq n} \binom{n-m}{m} P^{n-2m} P^m$$

$$= \sum_{0 \leq 2m \leq n} \binom{n-m}{m} P^{n-m},$$

which agrees with the result due to Barakat above. Note that Bruckman [3] has provided a neater proof for (4.3) in the case when $P = 1$.

5. CONCLUDING COMMENTS

Other properties can be readily developed to relate the $f_n^{(r)}$ and $g_n^{(r)}$ to other parts of the recurrence relation theory. For instance, we can prove that

$$(5.1) \quad \left\{ \begin{matrix} n \\ j \end{matrix} \right\} \prod_{i=2}^{j+1} g_{n-2i+4}^{(i)}$$

where $\left\{ \begin{matrix} n \\ j \end{matrix} \right\}$ is the analogue of the binomial coefficient used extensively in recurrence relation work (for example, Horadam [9]), and defined by

$$\left\{ \begin{matrix} n \\ j \end{matrix} \right\} = \frac{u_n u_{n-1} \dots u_{n-j-1}}{u_0 u_1 \dots u_{j-1}}$$

Proof:

$$\left\{ \begin{matrix} n \\ j \end{matrix} \right\} = \frac{(\alpha^{n+1} - \beta^{n+1})(\alpha^n - \beta^n) \dots (\alpha^{n-j+2} - \beta^{n-j+2})}{(\alpha - \beta)(\alpha^2 - \beta^2) \dots (\alpha^j - \beta^j)}$$

$$= g_n^{(2)} g_{n-2}^{(3)} \dots g_{n-2j+2}^{(j+1)} \quad [\text{by (2.2)}]$$

$$= \prod_{i=2}^{j+1} g_{n-2i+4}^{(i)}.$$

As another instance, consider

$$(5.2) \quad f_n^{(r)} = \beta^{kn} \left(\frac{\alpha}{\beta} \right)^k \Big|_{n+1}$$

where \underline{x}_n represents the n^{th} reduced Fermatian of index x as mentioned by Whitney [18] and utilized by Shannon [17]. It is defined formally by $\underline{x}_n = 1 + x + x^2 + \dots + x^{n-1}$.

Proof of (5.2):

$$f_n^{(r)} = \frac{(\alpha^k)^{n+1} - (\beta^k)^{n+1}}{(\alpha^k) - (\beta^k)} = \beta^{kn} \left(1 + \left(\frac{\alpha}{\beta} \right)^k + \dots + \left(\frac{\alpha}{\beta} \right)^{kn} \right) = \beta^{kn} (\alpha / \beta)^{kn}.$$

REFERENCES

1. K. T. Atanassov, L. C. Atanassov, & D. D. Sasselov. "A New Perspective to the Generalization of the Fibonacci Sequence." *Fibonacci Quarterly* **23.1** (1985):21-28.
2. R. Barakat. "The Matrix Operator e^x and the Lucas Polynomials." *J. Math. and Physics* **43.4** (1964):332-35.
3. P. Bruckman. Solution to Problem H-233. *Fibonacci Quarterly* **14.1** (1976):90-92.
4. L. Carlitz. "The Characteristic Polynomial of a Certain Matrix of Binomial Coefficients." *Fibonacci Quarterly* **3.2** (1965):81-89.
5. R. D. Carmichael. "On the Numerical Factors of the Arithmetic Forms $\alpha^n \pm \beta^n$." *Annals of Math. (2)* **15.1** (1913):30-70.
6. D. Dickinson. "On Sums Involving Binomial Coefficients." *Amer. Math. Monthly* **57.1** (1950):82-86.
7. M. Feinberg. "New Slants." *Fibonacci Quarterly* **2.3** (1964):223-27.
8. V. C. Harris & C. C. Styles. "A Generalization of Fibonacci Numbers." *Fibonacci Quarterly* **2.4** (1964):277-89.
9. A. F. Horadam. "Generating Functions for Powers of a Certain Generalized Sequence of Numbers." *Duke Math. J.* **32.3** (1965):437-46.
10. A. F. Horadam. "Basic Properties of a Certain Generalized Sequence of Numbers." *Fibonacci Quarterly* **3.3** (1965):161-76.
11. D. H. Lehmer. "An Extended Theory of Lucas' Functions." *Annals of Math. (2)* **31.4** (1930):419-48.
12. E. Lucas. *The Theory of Simply Periodic Numerical Functions*. Ed. D. A. Lind, tr. S. Kravitz. Santa Clara, Calif.: The Fibonacci Association, 1969.
13. W. L. McDaniel. "The G.C.D. in Lucas Sequences and Lehmer Number Sequences." *Fibonacci Quarterly* **29.1** (1991):24-29.
14. E. P. Miles, Jr. "Generalized Fibonacci Numbers and Associated Matrices." *Amer. Math. Monthly* **67.8** (1960):745-52.
15. J. A. Raab. "A Generalization of the Connection between the Fibonacci Sequence and Pascal's Triangle." *Fibonacci Quarterly* **1.3** (1963):21-31.
16. A. G. Shannon. "Iterative Formulas Associated with Generalized Third-Order Recurrence Relations." *SIAM J Appl. Math.* **23.3** (1972):364-68.
17. A. G. Shannon. "Ramanujan's q -Series as Generalized Binomial Coefficients." Ramanujan's Birth Centenary Year International Symposium on Analysis, Dec.26-28, 1987, Pune, India.
18. R. E. Whitney. "On a Class of Difference Equations." *Fibonacci Quarterly* **8.5** (1970):470-75.
19. H. C. Williams. "On a Generalization of the Lucas Functions." *Acta Arithmetica* **20.1** (1972):33-51.
20. D. Zeitlin. "On Solutions of Homogeneous Linear Difference Equations with Constant Coefficients." *Amer. Math. Monthly* **68.2** (1961):134-37.

