

TOPICS ON MEIXNER FAMILIES

MAREK BOŻEJKO

*Institut Matematyczny, Uniwersytet Wrocławski,
Pl. Grunwaldzki 2/4,
50-384 Wrocław, Poland.*

NIZAR DEMNI

*Faculty of Mathematics,
Bielefeld University, Germany.*

Abstract. We shed the light on the inter-connections between different characterizations leading to the classical Meixner family. This allow to set free analogs of both the Sheffer's and the Al-Salam and Chihara's characterizations in the classical case by the use of the free derivative operator. The paper is closed with a discussion of the q -deformed case, $|q| < 1$.

1. Introduction. A remarkable and exciting feature of Mathematics is the fact that several independent and seemingly unrelated works may lead to a common result. It is often the lack of communication among various schools that prevents the full understanding of the existing inter-connections, thereby arises the problem of unifying the different approaches, which stands for an important, useful and also tricky task. In this spirit, the so-called *Meixner family* of probability distributions, referred to as *Meixner distributions*, admits the following characterizations:

- When it first appeared in [23], the Meixner family was defined as the set of probability measures μ with finite exponential moments in a neighborhood of zero such that:

$$\psi(z, x) := \sum_{n \geq 0} P_n(x) z^n = \frac{e^{xH(z)}}{\mathbb{E}(e^{XH(z)})}, \quad (1)$$

where H is analytic around $z = 0$ such that $H(0) = 0, H'(0) = 1$, X is a random

2000 *Mathematics Subject Classification:*

Key words and phrases: Meixner families; orthogonal polynomials; free and quantum probability

Nizar Demni was supported by CRC 701.

The paper is in final form and no version of it will be published elsewhere.

variable in some probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with law $\mu = \mathbb{P} \circ X^{-1}$ and $(P_n)_{n \geq 0}$ is the set of orthogonal polynomials (OP) with respect to μ . ψ is an OP generating function of exponential-type. Up to translations and dilations, the Meixner family consists of Gaussian, Poisson, Gamma, negative binomial, Meixner and binomial distributions. The latter being non-infinitely-divisible in the classical sense. Another proof of this characterization was given in [19].

- Sheffer characterized $(P_n)_n$ as the set of OP of type zero ([26]).
- Meixner laws are the solutions of a quadratic regression problem ([21]).
- Morris showed that the variance of the natural exponential family parametrized by the mean associated with a probability measure μ , is an at most quadratic polynomial in the mean if and only if μ belongs to the Meixner family ([24]).
- OP with respect to Meixner laws form a distinguished subclass of the so-called Al-Salam and Chihara set of OP ([1]).

After the birth of free probability theory, for which miscellaneous analogs of known results in classical probability theory were derived, the *free Meixner family* was defined in [4] and covers six probability distributions called analogously to their classical counterparts free Gaussian or Wigner, free Poisson or Marchenko-Pastur, free Gamma, free negative binomial, free Meixner and free binomial known also as the free stationary Jacobi law ([13],[15],[16]). Their study revealed free versions of some of the above characterizations: the exponential-type OP generating functions are replaced by generating functions of Cauchy-Stieltjes-type, ([4], [10],[20]), *free Meixner distributions* solve a quadratic regression problem in a free setting ([9]) and to them are associated Cauchy-Stieltjes-type families with linear means and quadratic variances ([11]). To our best knowledge, no free analogs of both the Sheffer's and the Al-Salam and Chihara's characterizations exist in literature.

In the present paper, we highlight some intrinsic connections between the different characterizations in both the classical and the free settings. More precisely, we start by explaining how the Sheffer's characterization leads to Al-Salam and Chihara results and vice-versa. Then we do the same for both the Kubo's and the Morris characterizations. In the free setting, the connection is set between Bryc and Ismail's characterization using Cauchy-Stieltjes families of quadratic variances and the one given by the authors via Cauchy-Stieltjes-type generating functions of OP ([10]).

The study of those connections shows a parallel between the classical and free settings manifested by classical and free cumulants generating functions on the one side and by classical and free derivatives on the other side. As a matter of fact, it reveals what free Sheffer's and free Al-Salam and Chihara characterizations should be. Moreover, we observed that, to a given OP generating function of a probability measure satisfying suitable integrability conditions, corresponds a family of probability measures parametrized by the mean such that the variance is an at most quadratic polynomial in the mean (we will say for short 'an at most quadratic variance'). This fact is an immediate consequence of the orthogonality or equivalently the three-terms recurrence relation satisfied by the OP. Besides, the coefficients of the variance are easily expressed in terms of the Jacobi-Szegő parameters. This general observation gives a machinery that we used to solve the pro-

blem of characterizing probability distributions μ_λ (of finite all order moments) having *ultraspherical-type* generating functions ([17]). The paper is closed with pointing out the Difficulties arising in the q -deformed case, $|q| < 1$.

REMARK 1.1. *The Meixner family will be referred to as the classical Meixner family. Besides, we shall often use the word ‘law’ instead of ‘probability distributions’ and also the acronymes ‘Meixner, Al-Salam Chihara, Sheffer, Kubo etc...’ to refer to the corresponding characterizations cited above. We hope there will be no confusion while reading the paper.*

2. On Sheffer and Al-Salam and Chihara characterizations. Few years after the Meixner’s classification was given, Sheffer ([26]) provided an easy proof of it while studying the following general situation: let $(Q_n)_n$ be a set of polynomials, that is Q_n is of degree n for each $n \geq 0$, which are not necessarily orthogonal. Let D denote the derivative operator and take $Q_0(x) = 1$, then there exists a (unique) differential operator $J := \sum_{n \geq 1} c_n D^n$ with constant coefficients $(c_n)_n, c_1 \neq 0$ such that $JQ_n = Q_{n-1}, Q_{-1} = 0$. $(Q_n)_n$ is then said to be a set of *type zero* with respect to J or belongs to J . Sheffer showed that being a set of type zero is equivalent to

$$\sum_{n \geq 0} Q_n(x) z^n = A(z) e^{xH(z)} \quad (2)$$

where A is an entire function around zero (called the determining function) and H is the inverse function of J when the latter is viewed as an entire function (of course, the series defining J is supposed to be a convergent). It is then obvious that one immediately recovers the Meixner family under the additional orthogonality assumption on $(Q_n)_n$. In the middle 70’s, Al-Salam and Chihara ([1]) addressed the following problem: characterize the set of OP $(r_n)_n, (q_n)_n$ such that their convolution

$$s_n(x, y) = \sum_{k=0}^n r_k(x) q_{n-k}(y) \quad (3)$$

defines a set of OP in the variable x for all y . $(r_n)_n$ and $(q_n)_n$ belong then to the so-called Al-Salam and Chihara set of OP. If one further requires that $s_n(x, y) := p_n(x + y)$ for some polynomial p_n , then $r_n = q_n$ for all n and $(r_n)_n$ are OP with respect to a probability measure from the classical Meixner family.

REMARK 2.1. *This last characterization was firstly noticed by Lancaster in [22] and we would like to thank Prof. Gérard Letac for helping us with this reference.*

2.1. From Sheffer to Al-Salam and Chihara. Let us now see how one carries the Al-Salam and Chihara problem into the setting of Sheffer. Let $(r_n)_n, (q_n)_n, (p_n)_n$ three sets of OP related by the convolution below

$$p_n(x + y) = \sum_{k=0}^n r_k(x) q_{n-k}(y) \quad (4)$$

and let us prove that $(r_n), (q_n)$ are of type zero with respect to the same operator. Denote J and \tilde{J} the (unique) operators ([26]) such that $J(r_n) = r_{n-1}$ and $\tilde{J}(q_n) = q_{n-1}$ and

write

$$J = \sum_{k \geq 1} c_k(x) D^k, \quad \tilde{J} = \sum_{k \geq 1} \tilde{c}_k(x) D^k$$

where c_k, \tilde{c}_k are polynomials of degree at most $k - 1$ and $c_1 = \tilde{c}_1 = 1$. Note first that if (4) holds, then $c_n(x) = \tilde{c}_n(x)$ for all n and this follows from the symmetry of $(x, y) \mapsto p_n(x + y)$ in both arguments. More precisely,

$$\begin{aligned} J(p_n(\cdot + y))(x) &= \sum_{k=1}^n r_{k-1}(x) q_{n-k}(y) = p_{n-1}(x + y) \\ &= \sum_{k=0}^{n-1} r_k(y) q_{n-k-1}(x) = \tilde{J}(p_n(\cdot + y))(x) \end{aligned}$$

which proves our claim by induction on the degree n . Again, the symmetry gives

$$p_{n-1}(x + y) = J(p_n(\cdot + y))(x) = J(p_n(x + \cdot))(y)$$

which yields $c_n(x) = c_n(y) = c_n$ for all n . As a result, $(r_n)_n, (q_n)_n$ are of type zero with respect to the same operator J (hence the same H) and Sheffer's result applies.

REMARK 2.2. *Since the expectation of $\psi(z, X)$ equals 1 when $(P_n)_n$ is a set of OP, then $A(z) = 1/\mathbb{E}(e^{H(z)x})$ and is then entirely determined by the knowledge of H . Thus, $r_n = q_n$ for all n whenever both sets of OP belong to the same lowering operator J . This claim is no more true for sets of not necessarily OP.*

2.2. From Al-Salam and Chihara to Sheffer. Consider a set $(r_n)_n$ of OP with respect to a probability distribution μ having a finite exponential moment in a neighborhood of zero and assume that it is of type zero with respect to J :

$$J = \sum_{n \geq 1} c_n D^n, \quad c_1 = 1.$$

Form the convolution

$$s_n(x, y) = \sum_{k=0}^n r_k(x) r_{n-k}(y),$$

it then follows that $J s_n(\cdot, y) = J(s_n(x, \cdot))$ which implies (by induction) that $s_n(x, y) = p_n(x + y)$ for a certain polynomial p_n . The orthogonality of $(p_n(\cdot + y))_n$ follows from

$$\begin{aligned} \int_{\mathbb{R}} p_n(x + t) p_m(x + t) \mu(dx) \mu(dt) &= \sum_{k=0}^n \sum_{l=0}^m \int_{\mathbb{R}} r_k(x) r_l(x) \mu(dx) \int_{\mathbb{R}} r_{n-k}(t) r_{m-l}(t) \mu(dt) \\ &= \sum_{k=0}^n \sum_{l=0}^m \|r_k\|^2 \|r_{n-k}\|^2 \delta_{kl} \delta_{n-k, m-l} = \sum_{k=0}^n \|r_k\|^2 \|r_{n-k}\|^2 \delta_{n, m}, \end{aligned}$$

where μ is the orthogonality probability distribution of $(r_n)_n$.

REMARK 2.3. *Since*

$$p_n(x) = \sum_{k=0}^n r_k(x - y) q_{n-k}(y),$$

for all y , then $(p_n)_n$ belongs to J too and by orthogonality, $p_n = r_n$ for all n .

3. On Kubo and Morris characterizations. In the early eighties, the classical Meixner family appeared in [24] as the set of probability measures which generate natural exponential families parametrized by the means, such that the variance of a given exponential family is at most quadratic of the mean. More precisely, given a (non-degenerate) probability measure μ with finite exponential moments in some neighborhood of zero, its natural exponential family is the one-parameter family of probability measures defined by

$$\mathbb{P}_\theta(dx) := \frac{e^{\theta x}}{\mathbb{E}(e^{\theta X})} \mu(dx).$$

Let z and V denote respectively the mean and the variance of \mathbb{P}_θ for a given θ . Then $z = \tau'(\theta)$ and $V = \tau''(\theta)$ where

$$\tau(\theta) := \log \mathbb{E}(e^{\theta X})$$

is the cumulants generating function of μ . Since $V = V(\theta) > 0$, then τ' is invertible and $\theta = (\tau')^{-1}(z) := S(z)$ makes sense for small z . We also have $S(0) = 0, S'(z) = 1/V(z)$ and $S'(0) \neq 0$ which one may suppose to equal 1. This gives rise to a new exponential family parametrized by the mean z

$$\mathbb{P}_z(dx) := \frac{e^{S(z)x}}{\mathbb{E}(e^{S(z)X})} \mu(dx)$$

of mean z and variance $V(z) = \tau''(S(z))$. It was shown in [24] that V is at most quadratic in z if and only if μ belongs to the classical Meixner family.

Few years ago, another proof of the Meixner's classification was given in ([19]) via OP generating functions using Asai-Kuo-Kubo's criterion ([2]). Keeping the notations in (2), the author proved that (2) generates orthogonal polynomials if and only if

$$bH^{-1}(z) + a = \tau'(z), \quad (5)$$

where $a \in \mathbb{R}, b > 0$ are respectively the mean and the variance of the orthogonality measure and

$$H'(z) = \frac{1}{\beta z^2 + \gamma z + 1}, \quad (\beta \geq 0) \text{ or } (\beta < 0, b/\beta \in \mathbb{R} \setminus \mathbb{Z}). \quad (6)$$

The values $(\beta < 0, b/\beta \in \mathbb{Z})$ correspond to a signed measure. Note that, for $a = 0, b = 1$. (5) together with (6) give a Ricatti-type equation for τ' :

$$\tau''(z) = (H^{-1})'(z) = \frac{1}{H'(\tau'(z))} = \beta(\tau')^2(z) + \gamma\tau'(z) + 1.$$

3.1. From Kubo to Morris. It is known that the analyticity assumptions on S (since $S^{-1} = \tau'$ and τ' is analytic near $z = 0$) allow to expand

$$\frac{e^{S(z)x}}{\mathbb{E}(e^{S(z)x})} = \sum_{n \geq 0} P_n(x) z^n, \quad (7)$$

where P_n is a polynomial of exact degree n and $(P_n)_{n \geq 0}$ need not to be orthogonal (see [25] p.45). Assume that μ has zero mean and unit variance (standard), then one uses

Kubo's results above after identifying S with H so that $S'(z) = H'(z) = 1/V(z)$ fits (6) (one further needs to check the sign of the parameters involved in the variance).

3.2. From Morris to Kubo. Let $(V_n)_n$ be the set of monic OP with respect to the standard μ and consider the exponential generating function

$$\frac{e^{H(z)x}}{\mathbb{E}(e^{H(z)X})} = \sum_{n \geq 0} a_n V_n(x) z^n = \sum_{n \geq 0} \frac{1}{n!} V_n(x) z^n \quad (8)$$

where the last equality follows from substituting x with x/z then letting $z \rightarrow 0$ and finally using $H'(0) = 1$. It is known that

$$xV_n(x) = V_{n+1}(x) + \alpha_n V_n(x) + \omega_n V_{n-1}(x), \quad V_{-1} := 0, \omega_0 := 1$$

where $(\alpha_n)_n, (\omega_n)_n$ are the Jacobi-Szegő parameters ([14]). Then, intertwining the order of integration and using the orthogonality of $(V_n)_n$, it easily follows that

$$m(z) := \sum_{n \geq 0} \int_{\mathbb{R}} x V_n(x) \mu(dx) \frac{z^n}{n!} = \alpha_0 + \omega_1 z = z$$

since μ is standard. Similarly, one gets

$$V(z) = (\omega_2 - 2) \frac{z^2}{2} + \alpha_1 z + 1$$

which fits into the work of Morris and identifies the coefficients β, γ in (6).

What is quite interesting is that every OP generating function of a given measure satisfying suitable integrability conditions defines a one-parameter family of probability measures parametrized by the mean such that, for a fixed value of the parameter (the mean), the variance is at most quadratic in the mean. When the OP generating function is handable (which is not the case for instance for the q -Meixner family due to the infinite product, [5]), one gets a machinery that can be used to characterize a family of probability measures via their OP generating functions. This is, as we will see, the case of the free Meixner family with Cauchy-Stieltjes-type OP generating functions and of a natural generalization of ultraspherical polynomials ([17]).

4. Free Meixner family. A parallel definition of Lévy-Meixner processes in free probability theory appeared in [4] after the pioneering work of Biane [7] on free processes with free increments, where the author derived free analogs of exponential martingales involving free cumulants generating functions. Then, the free Meixner family was defined in [9] by proving a free Laha-Lukacs characterization and it consists of six laws which we think of as the free analogs of the six laws forming the classical Meixner family (see the introduction). Only the free binomial distribution is not infinitely divisible with respect to the free additive convolution. Similarly to the classical case, one can set the connection between Bryc and Ismail characterization via Cauchy-Stieltjes families with at most quadratic variances and the one recently given by the authors via Cauchy-Stieltjes-type OP generating functions ([9]).

On the one hand, a combination of [4] and [9] allowed us to characterize the free Meixner distributions and Asai-Kuo-Kubo criterion ([2]) was used to derive analogous identities to (5) and (6) involving the free cumulants generating function ([10]). More

precisely, Let μ be a standard probability distribution with finite all order moments. Then, μ belongs to the free Meixner family if and only if

$$\psi(z, x) := \sum_{n \geq 0} P_n(x) z^n = \frac{1}{u(z)[f(z) - x]} \quad (9)$$

where

$$f(z) = K(u(z)) = z + \frac{1}{u(z)} \quad (10)$$

$$\frac{u(z)}{z} = \frac{1}{1 + az + bz^2} \quad (11)$$

for $a \in \mathbb{R}, b \geq -1$ (parameters of μ and they do not have to be confused with a, b used previously), where K is the right inverse in a neighborhood of zero of the Cauchy-Stieltjes transform of μ :

$$G(z) := \int \frac{1}{z - x} \mu(dx), \quad z \in \mathbb{C} \setminus \text{supp}(\mu).$$

The second equality in (10) is analogous to (5) if it is written as $R(u(z)) = z$ where $R(z) = K(z) - 1/z$ is the free cumulants generating function, while (11) is analogous to (6) where the classical derivative is replaced by the free derivative defined by

$$\begin{cases} D^0(f)(z) = \frac{f(z) - f(0)}{z}, & z \neq 0 \\ D^0(f)(0) = f'(0). \end{cases} \quad (12)$$

On the other hand, to a compactly-supported probability measure is associated a Cauchy-Stieltjes family defined by ([12]):

$$\left\{ \mathbb{P}_\theta(dx) := \frac{1}{M(\theta)} \frac{1}{1 - \theta x} \mu(dx) \right\},$$

for small complex θ , where M is the moment generating function defined by

$$M(\theta) := \int_{\mathbb{R}} \frac{1}{1 - \theta x} \mu(dx) = \frac{1}{\theta} G\left(\frac{1}{\theta}\right).$$

If μ is not a Dirac mass, then the mean of \mathbb{P}_θ

$$z := \int_{\mathbb{R}} x \mathbb{P}_\theta(dx) = \frac{M(\theta) - 1}{\theta M(\theta)} := k(\theta)$$

is invertible as a function of small enough θ , thus a new parametrization by the mean is performed on \mathbb{P}_θ giving rise to

$$\mathbb{P}_z(dx) := \frac{1}{M(v(z))} \frac{1}{1 - v(z)x} \mu(dx) \quad \theta := k^{-1}(z) = v(z).$$

Let $V = V(z)$ denote the variance of P_z for z small enough. Then:

$$\mathbb{P}_z(dx) := \frac{V(z)}{V(z) + z(x - z)} \mu(dx) \quad (13)$$

when μ has mean zero. It was shown in [12] that V is quadratic if and only if μ belongs to the free Meixner family. To deduce this result using OP generating functions, expand

the ratio in (13) as

$$\frac{V(z)}{V(z) + z(x-z)} = \sum_{n \geq 0} P_n(x) z^n$$

with non-necessarily OP. Set $u(z) := z/V(z)$ with $V(z) = 1 + az + bz^2$, then equation (2.1) in [12] fits the second equality in (10). It follows that the density in (13) is given by (9) which gives the result.

Conversely, suppose we are given a standard probability measure with finite all order moments and the OP generating function (9) with $u(0) = 0, u'(0) = 1, uf(0) = 1$. Then, one defines a Cauchy-Stieltjes family $(\mathbb{P}_z)_z$ parametrized by the mean with an at most quadratic variance. Since $z \mapsto R(z)$ is defined and analytic in a neighborhood of zero, then μ is compactly-supported ([6]). Moreover, setting $V(z) = z/u(z)$, \mathbb{P}_z takes the form of (13) so that Theorem 1.2 in [12] applies.

5. Free-type zero polynomials and free Sheffer's characterization. Let $(U_n)_n$ be the set of monic Tchebycheff polynomials of the second kind ([14]) defined by

$$\sum_{n \geq 0} U_n(x) z^n = \frac{1}{1 - zx + z^2}.$$

It is easy to see from the above OP generating function that $(U_n)_n$ can not be a set of finite A, B, C -types described in [26] (the A -type includes the type zero). This was already noticed for the Legendre polynomials (see p.621-622 in [26]) and the same reasoning applies for $(U_n)_n$. Easily speaking, this is due to the fact that the above generating function is not of an exponential-type (A -type) and its derivative with respect to either x or z will raise it to the square so that, for each n , the polynomials coefficients in the expansion of U'_n (B -type) and nU_n (C -type) in terms of (U_{n-1}, \dots, U_0) have unbounded degrees in n . Nevertheless, they will be of type zero in the free sense, that is, when substituting the classical derivative D by its free analog D^0 defined by (12) and the exponential function by $x \mapsto (1-x)^{-1}$. Following the lines of Sheffer, we first define 'free differential operators' by

$$J^0 := \sum_{n \geq 0} c_n(x) (D^0)^n$$

where c_n are polynomials. We will say that a set of polynomials $(P_n)_n$ (that is P_n has degree n) belongs to J if $J^0 P_n = P_{n-1}, P_0 \neq 0, P_{-1} := 0$. Since $(D^0)^k x^n = x^{n-k}$ for $1 \leq k \leq n$ and $D^0 1 = 0$, then the sum defining J^0 terminates when acting on polynomials. In addition, J^0 maps x^n to a polynomial of degree $\leq n-1$ if and only if $c_0 = 0$ and the degree of $c_k, 1 \leq k \leq n$ does not exceed $k-1$. Writing in that case

$$c_k(x) = c_{k,0} + c_{k,1}x + \dots + c_{k,k-1}x^{k-1}, \quad 1 \leq k \leq n,$$

then J^0 maps x^n to a polynomial of exact degree $n-1$ if and only if there exist at least k such that $c_{k,k-1} \neq 0$.

Second, to a given set of polynomials $(P_n)_n$ corresponds a unique free differential operator J^0 such that $J^0 P_n = P_{n-1}$ and $c_1 \neq 0$, for otherwise P_1 will be constant which disagrees the fact that $(P_n)_n$ is a set of polynomials. Recall also ([26]) that $(P_n)_n$ and $(Q_n)_n$ belong

to the same operator J^0 if and only if there exists a sequence $(a_n)_{n \geq 0}$ such that for each n

$$P_n(x) = a_0 Q_n(x) + \dots + a_{n-1} Q_1(x) + a_n Q_0(x), \quad a_0 \neq 0. \quad (14)$$

Finally, $(P_n)_n$ is said to be of *free-type* k , $k \in \mathbb{N}$ with respect to J^0 if it belongs to J^0 and, for all $n \geq 1$, the degree of c_n is at most k . Thus, a set of *free-type zero* if and only if $c_n(x) = c_n \in \mathbb{R}$ for all $n \geq 1$.

Let us suppose that $c_1 = 1$ (which holds for instance if $(P_n)_n$ are monic) such that J^0 , viewed as an entire series, is invertible near 0 with inverse ρ with $\rho'(0) = 1$. Since D^0 acts on $x \mapsto (1 - zx)^{-1}$ by multiplication by z similarly as D does on $x \mapsto e^{zx}$, then we claim that a set of polynomials $(P_n)_n$ is of free type zero if and only if

$$\sum_{n \geq 0} P_n(x) z^n = A(z)(1 - \rho(z)x)^{-1}, \quad A(0) = \sum_{n \geq 0} a_n z^n. \quad (15)$$

As a result, the free Sheffer characterization is stated as: $(P_n)_n$ is a set of monic OP with respect to a probability measure μ with finite all order moments and of free-type zero with respect to a given J^0 if and only if μ belongs to the free Meixner family with $\rho(J(z)) = z$. Recall from (10) that:

$$\rho(z) = \frac{1}{f(z)} = \frac{1}{K(u(z))}, \quad A(z) = \frac{1}{f(z)u(z)} = \frac{f(z) - z}{f(z)} = 1 - z\rho(z)$$

when μ is a standard free Meixner distribution of parameters $a \in \mathbb{R}$, $b \geq -1$. This shows that the knowledge of J is enough to characterize $(P_n)_n$ and implies that $J(z) = R(G(1/z))$ for small enough z (recall that $R(u(z)) = z$). Using the relation

$$R(G(z)) + \frac{1}{G(z)} = z$$

it follows that

$$J(z) = \frac{1}{z} - \frac{1}{G(1/z)} = \frac{1}{z} - F\left(\frac{1}{z}\right).$$

REMARK 5.1. *The lowering operator J was used in [4] when characterizing OP of Cauchy-Stieltjes type generating functions. However, no explicit expression as a free differential operator with constant coefficients was given.*

6. Free Al-Salam and Chihara's characterization. Let μ be a standard free Meixner distribution with parameters $a \in \mathbb{R}$, $b \geq -1$ and $(P_n)_n$ be the set of monic OP with respect to μ . Recall that these polynomials satisfy the three-terms recurrence relation ([4]):

$$xP_n(x) = P_{n+1}(x) + aP_n(x) + (1+b)P_{n-1}(x), \quad n \geq 2$$

with $P_0(x) = 1$, $P_1(x) = x - a$, and that

$$P_n(x) = (1+b)^{n/2} U_n(x-a, \sqrt{1+b}) + a U_{n-1}(x-a, \sqrt{1+b}) + b U_{n-2}(x-a, \sqrt{1+b}) \quad (16)$$

where

$$U_n(x-a, \sqrt{1+b}) := (1+b)^{n/2} U_n\left(\frac{x-a}{\sqrt{1+b}}\right)$$

is the monic shifted Tchebycheff polynomial of the second kind. Using the elementary relation

$$\frac{x-y}{(z-x)(z-y)} = \frac{1}{z-x} - \frac{1}{z-y}, \quad x \neq y$$

then

$$\sum_{n \geq 0} \frac{P_n(x) - P_n(y)}{x-y} z^n = u(z)\psi(z,x)\psi(z,y)$$

where ψ is given by (9). Since and $P_0(x) = P_0(y) = 1$ and $z/u(z) = 1 + az + bz^2$, then

$$(1 + az + bz^2) \sum_{n \geq 0} \frac{P_{n+1}(x) - P_{n+1}(y)}{x-y} z^n = \psi(z,x)\psi(z,y). \quad (17)$$

Set

$$R_n(x, y) := \frac{P_{n+1}(x) - P_{n+1}(y)}{x-y},$$

then $R_0(x) = 1$ and R_n is a linear combination of

$$\frac{x^{k+1} - y^{k+1}}{x-y} = x^k + x^{k-1}y + \cdots + xy^{k-1} + y^k = \sum_{r=0}^k y^r (D_x^0)^r (x^k) = \sum_{r=0}^k x^r (D_y^0)^r (y^k)$$

as one easily checks. Thus, (17) is equivalent to

$$\sum_{k=0}^n P_k(x)P_{n-k}(y) = R_n(x, y) + aR_{n-1}(x, y) + bR_{n-2}(x, y), \quad R_{-1} = R_{-2} := 0. \quad (18)$$

For $a = b = 0$, one has

$$\sum_{k=0}^n U_k(x)U_{n-k}(y) = \frac{U_{n+1}(x) - U_{n+1}(y)}{x-y}$$

which was already considered in [5] when studying free Appell sets. One also easily checks that if a polynomial s_n in two variables (x, y) of total degree n satisfies the Cauchy equation $D_x^0 s_n = D_y^0 s_n$, then it is a linear combination of $x^k + x^{k-1}y + xy^{k-1} + y^k$, $0 \leq k \leq n$. Therefore, the free Al-Salam and Chihara's characterization is stated as

PROPOSITION 6.1. *Let $(r_n)_n, (q_n)_n$ be two sets of monic OP and let*

$$s_n(x, y) = \sum_{k=0}^n r_n(x)q_{n-k}(y)$$

be their convolution. Then, s_n has the expansion

$$s_n(x, y) = \sum_{k=0}^n b_k \frac{x^{k+1} - y^{k+1}}{x-y}$$

if and only if $r_n = q_n$ for all n and $(r_n)_n$ belongs to the free Meixner family.

REMARKS. 1/A remarkable difference with the classical setting is the fact that $(s_n(\cdot, y))_n$ no longer defines a set of OP. Let us for instance consider the case of monic Tchebycheff polynomials $(U_n)_n$, then one easily derives for all $n \geq 1$,

$$xR_n(x, y) = R_{n+1}(x, y) + R_{n-1}(x, y) - U_{n+1}(y).$$

This also can be noticed from Al-Salam and Chihara paper [1], where it is shown that, in order to get the desired orthogonality, it is not allowed, except for the classical Meixner family, to take $r_n = q_n, n \geq 1$.

2/Integrating (18) with respect to μ , one obtains:

$$Q_n(x) + aQ_{n-1}(x) + bQ_{n-2}(x) = P_n(x), \quad Q_n(x) := \int_{\mathbb{R}} \frac{P_{n+1}(x) - P_{n+1}(y)}{x - y} \mu(dy).$$

Q_{n-1} is the n -th numerator polynomial of P_n ([14]) and Q_n is known to be equal to $U_n(x - a, \sqrt{1 + b})$ ([14] p. 89). Thus, the above equality coincides with (16).

Proof: the sufficiency was already proved and the necessity can be proved using the free Sheffer's characterization. Indeed, let

$$J = \sum_{k \geq 1} c_k(x)(D^0)^k, \quad \tilde{J} = \sum_{k \geq 1} \tilde{c}_k(x)(D^0)^k,$$

denote the unique operators satisfying $J(r_n) = r_{n-1}, \tilde{J}(q_n) = q_{n-1}$ ($c_1 = \tilde{c}_1 = 1$). Then, by symmetry of $(x, y) \mapsto s_n(x, y)$ in both variables, we first get that $J = \tilde{J}$. Using,

$$\begin{aligned} D_x^0(x^k + x^{k-1}y + \dots + xy^{k-1} + y^k) &= x^{k-1} + x^{k-2}y + \dots + xy^{k-2} + y^{k-1} \\ &= D_y^0(x^k + x^{k-1}y + \dots + xy^{k-1} + y^k) \end{aligned}$$

for $k \geq 1$, one has $J^0 s_n(\cdot, y) = J^0 s_n(x, \cdot)$. Thus, $c_n(x) = c_n(y) = c_n, n \geq 1$ by induction on n , for some constants $(c_n)_n$ so that $(r_n)_n$ is a family of OP of type zero. Moreover, we have already seen that a set of OP is entirely determined by the knowledge of J thereby $r_n = q_n$. ■

6.1. Wick product. We have already seen that for the classical Meixner family, the convolution polynomial $s_n(x, y) = p_n(x + y)$ for some polynomial p_n in one variable. In the free setting, we have a similar relation but with non commutative variables and even more, it is related to the so-called Wick product ([8]).

Let $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ be a complex Hilbert space and let $\mathcal{H}^{\otimes n}$ be the n -fold tensor product of H with $\mathcal{H}^{\otimes 0} = \mathbb{C}\Omega$ for a distinguished unit-norm vector $\Omega \in \mathcal{H}$ called the vacuum. Let $\Gamma_0(\mathcal{H})$ denote the full Fock space associated with H defined by

$$\Gamma_0(\mathcal{H}) := \bigoplus_{n \geq 0} \mathcal{H}^{\otimes n}$$

and for $f \in \mathcal{H}$, consider the creation and the annihilation operators on $\Gamma_0(\mathcal{H})$ defined respectively by

$$\begin{aligned} a_f^* \Omega &= f, \\ a_f^*(f_1 \otimes f_2 \cdots \otimes f_n) &= \langle f, f_1 \rangle f_2 \otimes \cdots \otimes f_n \end{aligned}$$

and

$$\begin{aligned} a_f \Omega &= 0, \\ a_f(f_1 \otimes f_2 \cdots \otimes f_n) &= f \otimes f_1 \otimes f_2 \otimes \cdots \otimes f_n. \end{aligned}$$

These operators satisfy the *free commutation* relation

$$a_f a_f^* = \langle f, f \rangle \mathbf{1},$$

where $\mathbf{1}$ is the identity operator ([8]). Take $f \in \mathcal{H}$ such that $\langle f, f \rangle = 1$, then it was shown in ([8]) that for all n

$$a_f^n + a_f^{n-1}a_f^* + \cdots + a_f(a_f^*)^{n-1} + (a_f^*)^n = U_n(a_f + a_f^*)$$

and it is known as the free Wick product. As a result, with the same notations of the above Proposition,

$$s_n(a_f, a_f^*) = \sum_{k=0}^n b_k U_k(a_f + a_f^*).$$

7. Concluding remarks: the q -deformed Meixner family, $|q| < 1$. Though both the classical and free settings provide different characterizations of their corresponding Meixner families, the q -deformed setting is rather complicated and we record below the main difficulties arising in this case. Let us first recall that the q -deformed Meixner family was already defined in [5] and its elements are (up to a rescaling) the orthogonality measures of the Al-Salam and Chihara OP characterized by the following property: $(r_n)_n$ and $(q_n)_n$ belong to the Al-Salam and Chihara family of OP if and only if their convolution

$$s_n(x, y) = \sum_{k=0}^n r_n(x)q_{n-k}(y)$$

defines a set of OP in x for infinitely many y ([1]). Up to a rescaling, their generating function is given by ([11], p.14)

$$\psi_q(z, x) = \prod_{k=0}^{\infty} \frac{1 + aq^k z + bq^{2k} z^2}{1 + (a - (1 - q)x)q^k z + (b + 1 - q)q^{2k} z^2},$$

for $a \in \mathbb{R}, b \geq -1 + \max(q, 0)$, which may be written as

$$\psi_q(z, x) = A_q(z) \prod_{k=0}^{\infty} \frac{1}{1 - (1 - q)q^k H_k(q, z)x} \quad (19)$$

for some analytic function $H(q, \cdot)$ around zero with $H(q, 0) = 0, H'(q, 0) \neq 0$. Thus, ψ_q can not in general be expressed through the q -exponential function e_q defined by

$$e_q(x) := \sum_{k \geq 0} \frac{x^k}{[k]_q!} = \prod_{k=0}^{\infty} \frac{1}{1 - (1 - q)q^k x}$$

whenever it makes sense. The difficulties arising in this case sum up into the infinite products or the infinite sums involved in the expressions, which reduce to only one term when $q = 0$:

1. The infinite product form of ψ_q prevents to derive q -analogs of relations (5), (6), (10), (11) by performing direct computations of the first and the second moment.
2. The following characterization via q -exponential families was given in [11]: a q -exponential family associated with a compactly-supported measure and parametrized by the mean has an at most quadratic variance if and only if the measure belongs to the q -Meixner family. Thus, regarding relations (6) and (11), one consi-

ders a quadratic variance function

$$V(z) = bz^2 + az + 1, \quad b \geq -1 + \max(q, 0), \quad a \in \mathbb{R}$$

of a q -Meixner distribution and defines f by:

$$D^q f(q, z) = \frac{1}{V(z)}, \quad f(q, 0) = 0, \quad z \in V(0),$$

where D^q is the q -derivative defined by

$$D^q f(q, z) := \frac{f(q, z) - f(q, qz)}{(1-q)z}.$$

It is easy to see that

$$f(q, z) = (1-q) \sum_{k \geq 0} \frac{q^k z}{1 + aq^k z + bq^{2k} z^2}.$$

Then, f is invertible and we think of its inverse as a q -analog of the free cumulants generating function for $|q| < 1$. Unfortunately, the inversion procedure stands for a hard task (use Lagrange's formula for instance).

3. Though e_q is uniquely determined by the requirement

$$D_x^q(e_q(zx)) = ze_q(zx),$$

it is not immediate that ψ_q is given by (19) since the function H may depend on both k and q .

4. Finally, the dependence of H on q shows that

$$D_x^q[\psi_q(z, x)\psi_q(z, y)] \neq D_q^x[\psi_q(z, x)\psi_q(z, y)]$$

in general (except in the case $a = b = 0$ corresponding to the q -Hermite polynomials, [8]). Nevertheless, a polynomial s_n in two variables x, y of degree n satisfies the cauchy equation $D_x^q s_n = D_y^q s_n$ if and only if s_n is a linear combination of the homogeneous polynomials

$$h_k(x, y) := \sum_{k=0}^n \binom{n}{k}_q x^k y^{n-k} \quad 1 \leq k \leq n.$$

where $\binom{n}{k}_q$ is the q -binomial coefficient ([8]) and the Wick product representation holds, that is

$$h_k(a_f, a_f^*) = H_k^q(a_f + a_f^*)$$

where H_k^q is the k -th q -Hermite polynomial and a_f, a_f^* are the annihilation and creation operators satisfying the q -commutation relation $a_f a_f^* - q a_f^* a_f = \mathbf{1}$ for a unit norm function f in an infinite Hilbert space ([8]).

Acknowledgments: both authors attribute special thanks to Prof. Friedrich Goetze. The first author would like to thank the organizing committee of the workshop for their hospitality.

Literatura

- [1] *W. Al-Salam, T. S. Chihara.* Convolutions of orthogonal polynomials. *SIAM J. Funct. Anal.* **7**, 1976, 16-28.
- [2] *N. Asai, I. Kubo, H. H. Kuo.* Multiplicative renormalization and generating function I. *Taiwan. J. Math.* **7**, **1**, 2003, 89-101.
- [3] *N. Asai, I. Kubo, H. H. Kuo.* Generating functions of orthogonal polynomials and Szegő-Jacobi parameters. *Probab. Math. Stat.* **23**, 2003, 273-291.
- [4] *M. Anshelevich.* Free martingale polynomials. *J. Funct. Anal.* **201**, 2003, 228-261.
- [5] *M. Anshelevich.* Appell polynomials and their relatives. *Int. Math. Res. Not.* **65**, 2004, 3469–3531.
- [6] *F. Benaych-Georges.* Taylor expansions of R -transforms: application to supports and moments. *Indiana Univ. Math. J.* **55**, no. 2, 2006, 465-481.
- [7] *P. Biane.* Processes with free increments. *Math. Z.* **227**, No 1, 1998, 143-174.
- [8] *M. Bożejko, B. Kümmerner, R. Speicher.* q -Gaussian processes: non-commutative and classical aspects. *Comm. Math. Phys.* **185**, no. 1, 1997, 129-154.
- [9] *M. Bożejko, W. Bryc.* On a class of free Lévy laws related to a regression problem. *J. Funct. Anal.* **236**, no. 1, 2006, 59-77.
- [10] *M. Bożejko, N. Demni.* Generating functions of Cauchy-Stieltjes type for orthogonal polynomials. *Submitted in Infinite. Dimen. Anal. Quantum Probab. Rel. Top.*
- [11] *W. Bryc, M. E. H. Ismail.* Approximation operators, q -exponential and free exponential families. *Available on arXiv.*
- [12] *W. Bryc, M. E. H. Ismail.* Cauchy-Stieltjes kernel families. *Available on arXiv.*
- [13] *M. Capitaine, M. Casalis.* Asymptotic freeness by generalized moments for Gaussian and Wishart matrices. Application to beta random matrices. *Indiana Univ. Math. J.* **53**, no. 2, 2004, 397-431.
- [14] *T. S. Chihara.* An Introduction to Orthogonal polynomials. *Gordon and Breach Science Publishers*, 1978.
- [15] *B. Collins.* Product of random projections, Jacobi ensembles and universality problems arising from free probability. *P. T. R. F.*, **133**, 2005, no. 3, 315-344.
- [16] *N. Demni.* Free Jacobi process. *J. Theor. Probab.* **21**, 2008, 118-143.
- [17] *N. Demni.* Ultraspherical type generating functions for orthogonal polynomials. *Probab. Math. Stat.* **29**, no. 2, 2009, 281-296.
- [18] *M. E. H. Ismail.* Classical and Quantum orthogonal polynomials in one variable. *Cambridge University Press*, 2005.
- [19] *I. Kubo.* Generating functions of exponential-type for orthogonal polynomials. *Inf. Dimens. Anal. Quantum Probab. Related Topics.* **7**, no 1, 2004, 155-159.
- [20] *I. Kubo, H. H. Kuo, S. Namli.* The characterization of a class of probability measures by multiplicative renormalization. *Communications on Stochastic Analysis.* **1**, no. 3, 2007, 455-472.
- [21] *R. G. Laha, E. Lukacs.* On a problem connected with quadratic regression. *Biometrika.* **47**, 1960, 335-343.
- [22] *H. O. Lancaster.* Joint probability distributions in the Meixner classes. *J. Roy. Stat. Soc. Series B.* **37**, no. 3, 1975, 434-443.
- [23] *J. Meixner.* Orthogonale polynomsysteme mit einer besonderern der erzeugenden funktion. *J. London. Math. Soc.* **9**, 1934, 6-13.
- [24] *C. N. Morris.* Natural exponential families with quadratic variance function. *Ann. Stat.* **10**, no. 1, 1982, 65-80.
- [25] *W. Schoutens.* Stochastic Processes and Orthogonal Polynomials. Lecture notes in Statistics. Springer, 2000.
- [26] *I. M. Sheffer.* Some properties of polynomial set of type zero. *Duke Math. J.* **5**, 1939, 590-622.