

Editorial Comment

Mark Feinberg is a fourteen-year-old student in the ninth grade of the Susquehanna Township Junior High School and recently became the Pennsylvania State Grand Champion in the Junior Academy of Science. This paper is based on his winning project and is editorially uncut. Mark Feinberg, in this editor's opinion, will go far in his chosen field of endeavor. Congratulations from the editorial staff of the Fibonacci Quarterly Journal, Mark!




Figure 1




Figure 3

## FIBONACCI-TRIBONACCI\*

MARK FEINBERG

For this Junior High School Science Fair project two variations of the Fibonacci series were worked out.

### "TRIBONACCI"

Just as in the Fibonacci series where each number is the sum of the preceding two, or  $p_{n+1} = p_n + p_{n-1}$ , the first variation is a series in which each number is the sum of the preceding three, or  $q_{n+1} = q_n + q_{n-1} + q_{n-2}$ ; hence the series is called "Tribonacci." Its first few numbers are

1, 1, 2, 4, 7, 13, 24, 44, 81, 149, 274, 504...

Like the Fibonacci series, the Tribonacci series is convergent. Where the Fibonacci fractions  $p_n/p_{n+1}$  and  $p_{n+1}/p_n$  converge on .6180339... and 1.6180339... , the Tribonacci fraction of any number of the series divided by the preceding one ( $q_n/q_{n+1}$ ) approaches .54368901... . While the Fibonacci convergents are termed "Phi" ( $\phi$ ), the Tribonacci convergents might be called "Tri-Phi" ( $\phi_3$ ).

Series-repeating characteristics are shown in the famed Fibonacci Golden Rectangle.\* A rectangle can be made of the Tribonacci series which also has series-repeating characteristics but since they are less obvious this rectangle might be called the "Silver Rectangle." Its length ( $q_{n+1}$ ) and its width ( $q_n$ ) make it proportionately longer than the Golden Rectangle.

By removing the squares  $q_n$  by  $q_n$  and  $q_{n-1}$  by  $q_{n-1}$ , two new rectangles in the proportion of the original appear (shaded areas). One is  $q_{n-1}$  by  $q_{n-2}$ ; but the other is composed of numbers not found in the Tribonacci series. This rectangle is  $(q_{n+1} - q_n)$  by  $(q_n - q_{n-1})$  and is formed of numbers from an intermediate series obtained by subtracting each Tribonacci number from the one after it.

---

\*See editorial remarks, page 70. Figure 1 appears on page 70.

By carrying the rectangle out farther new numbers found in neither the original Tribonacci series nor the intermediate series appear. These are of a second intermediate series and are obtained by subtracting each number of the first intermediate series from the succeeding one. New numbers of new intermediate series also appear by further carrying out the rectangle. These other series are formed by triangulating in the same way as the first two intermediate series. All these intermediate series are convergent upon the "Tri-Phi" values and each number in each of these series is the sum of the preceding three. Figure 2 shows the first two intermediate series.

$$\begin{array}{cccccccccccc} 1, & 1, & 1, & 3, & 5, & 9, & 17, & 31 & \dots \\ 0, & 1, & 2, & 3, & 6, & 11, & 20, & 37, & 68 & \dots \\ 1, & 1, & 2, & 4, & 7, & 13, & 24, & 44, & 81, & 149 & \dots \end{array}$$

Figure 2

The two Fibonacci convergents fit the quadratic equation  $x = 1 + 1/x$ . The Tribonacci convergent of any number in the series divided by the preceding one ( $q_{n+1}/q_n$ ) fits the cubic equation  $y = 1 + 1/y + 1/y^2$ . It is derived thus:

The formula giving any number in the series is

$$q_{n+1} = q_n + q_{n-1} + q_{n-2} .$$

Dividing by  $q_{n-1}$ :

$$\frac{q_{n+1}}{q_{n-1}} = \frac{q_n}{q_{n-1}} + 1 + \frac{q_{n-2}}{q_{n-1}} .$$

Let

$$\frac{q_{n+1}}{q_n} = t_n ; \quad \frac{q_n}{q_{n-1}} = t_{n-1} ; \quad \frac{q_{n-1}}{q_{n-2}} = t_{n-2}$$

Then since

$$\frac{q_{n+1}}{q_{n-1}} = \frac{q_{n+1}}{q_{n-1}} \cdot \frac{q_n}{q_n} = \frac{q_{n+1}}{q_n} \cdot \frac{q_n}{q_{n-1}} = t_n \cdot t_{n-1}$$

Therefore

$$t_n \cdot t_{n-1} = \frac{q_n}{q_{n-1}} + 1 + \frac{q_{n-2}}{q_{n-1}}$$

Substituting for the rest of the formula:

$$t_n \cdot t_{n-1} = t_{n-1} + 1 + \frac{1}{t_{n-2}}$$

Dividing by  $t_{n-1}$ :

$$t_n = 1 + \frac{1}{t_{n-1}} + \frac{1}{t_{n-1} \cdot t_{n-2}}$$

All the  $t_n$  terms converge upon one value ( $y$ ). Therefore " $y$ " can be substituted for all  $t_n$  terms. So

$$y = 1 + \frac{1}{y} + \frac{1}{y^2}$$

The convergent approached by any number of the series divided by the succeeding one ( $q_n/q_{n+1}$ ) fits the cubic equation  $1/y = 1 + y + y^2$  and is derived through a similar process.

Charting the Fibonacci convergent .6180339... on polar coordinate paper is known to produce the famed spiral found all over nature. By charting the Tribonacci convergent .54368901... a slightly tighter spiral is produced.

It is not known whether the Tribonacci series has any natural applications. A well-known Fibonacci application is of a hypothetical tree. If each limb were to sprout another limb one year and rest the next, the number of limbs per year would total 1, 2, 3, 5, 8... in Fibonacci sequence. However if each limb on the tree were to sprout for two years and rest for a year, the number of limbs per year would total 1, 2, 4, 7, 13... in Tribonacci sequence. See Figure 3, page 70.

Could such a tree as that on the right be called a "Tree-bonacci?"

#### "TETRANACCI"

The second variation of the Fibonacci sequence is a series in which each number is the sum of the preceding four numbers or  $r_{n+1} = r_n + r_{n-1} + r_{n-2}$

+  $r_{n-3}$ . Therefore this series is called "Tetranacci." Its first few numbers are:

1, 1, 2, 4, 8, 15, 29, 56, 108, 208, 401, 773...

Like the Fibonacci and Tribonacci series, the Tetranacci series is convergent. The fraction  $r_{n+1}/r_n$  converges upon 1.9275619... and fits the fourth power equation  $z = 1 + 1/z + 1/z^2 + 1/z^3$ .

The fraction  $r_n/r_{n+1}$  converges upon .51879006... which fits the equation  $1/z = 1 + z + z^2 + z^3$ .

The derivation of these formulas follows the same algebraic process as that given above and will be gladly furnished upon request.

XXXXXXXXXXXXXXXXXXXXXXXXXXXX

The Fibonacci Association invites Educational Institutions to apply for Academic Membership in the Association. The minimum subscription fee is \$25 annually. (Academic Members will receive two copies of each issue and will have their names listed in the Journal.)

#### REQUEST

The Fibonacci Bibliographical Research Center desires that any reader finding a Fibonacci reference send a card giving the reference and a brief description of the contents. Please forward all such information to:

Fibonacci Bibliographical Research Center,  
Mathematics Department,  
San Jose State College,  
San Jose, California

#### NOTICE TO ALL SUBSCRIBERS !!!

Please notify the Managing Editor AT ONCE of any address change. The Post Office Department, rather than forwarding magazines mailed third class, sends them directly to the dead-letter office. Unless the addressee specifically requests the Fibonacci Quarterly be forwarded at first class rates to the new address, he will not receive it. (This will usually cost about 30 cents for first-class postage.) If possible, please notify us AT LEAST THREE WEEKS PRIOR to publication dates: February 15, April 15, October 15, and December 15.