

Research Article

A Note on Some Identities of Frobenius-Euler Numbers and Polynomials

J. Choi,¹ D. S. Kim,² T. Kim,³ and Y. H. Kim¹

¹ Division of General Education-Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea

² Department of Mathematics, Sogang University, Seoul 121-742, Republic of Korea

³ Department of Mathematics, Kwangwoon University, Seoul 139-701, Republic of Korea

Correspondence should be addressed to D. S. Kim, dskim@sogang.ac.kr
and T. Kim, tkkim@kw.ac.kr

Received 28 November 2011; Accepted 9 January 2012

Academic Editor: Feng Qi

Copyright © 2012 J. Choi et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The purpose of this paper is to give some identities on the Frobenius-Euler numbers and polynomials by using the fermionic p -adic q -integral equation on \mathbb{Z}_p .

1. Introduction

Let p be a fixed odd prime number. Throughout this paper, \mathbb{Z}_p , \mathbb{Q}_p , and \mathbb{C}_p will denote the ring of p -adic rational integers, the field of p -adic rational numbers, and the completion of algebraic closure of \mathbb{Q}_p , respectively. Let \mathbb{N} be the set of natural numbers and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. The p -adic absolute value on \mathbb{C}_p is normalized so that $|p|_p = 1/p$. Assume that $q \in \mathbb{C}_p$ with $|1 - q|_p < 1$.

Let f be a continuous function on \mathbb{Z}_p . Then the fermionic p -adic q -integral on \mathbb{Z}_p for f is defined by Kim as follows:

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N \rightarrow \infty} \frac{1+q}{1+q^{p^N}} \sum_{x=0}^{p^N-1} f(x) (-q)^x, \quad (\text{see [1]}). \quad (1.1)$$

From (1.1), we note that

$$q^n I_{-q}(f_n) = (-1)^n I_{-q}(f) + (1+q) \sum_{l=0}^{n-1} (-1)^{n-1-l} f(l) q^l, \quad (1.2)$$

where $n \in \mathbb{N}$ and $f_n(x) = f(x+n)$ (see [1]). The ordinary Euler polynomials $E_n(x)$ are defined by

$$\frac{2}{e^t + 1} e^{xt} = e^{E(x)t} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}, \quad (1.3)$$

with the usual convention about replacing $E^n(x)$ by $E_n(x)$ (see [1–10]). In the special case, $x = 0$, $E_n(0) = E_n$ is called the n th Euler number.

As the extension of (1.3), the Frobenius-Euler polynomials are defined by

$$\frac{1-q}{e^t - q} e^{xt} = \sum_{n=0}^{\infty} H_n(q, x) \frac{t^n}{n!}, \quad (\text{see [2]}). \quad (1.4)$$

In the special case, $x = 0$, $H_n(q, 0) = H_n(q)$ is called the n th Frobenius-Euler number.

By (1.3) and (1.4), we easily get $H_n(-1, x) = E_n(x)$.

From (1.4), we note that

$$H_n(q, x) = \sum_{l=0}^n \binom{n}{l} H_l(q) x^{n-l} = (H(q) + x)^n, \quad (\text{see [2]}), \quad (1.5)$$

with the usual convention about replacing $H(q)^n$ by $H_n(q)$.

In this paper, we consider the fermionic p -adic q -integral on \mathbb{Z}_p for the Frobenius-Euler numbers and polynomials. From these p -adic q -integrals on \mathbb{Z}_p , we derive some new and interesting identities on the Frobenius-Euler numbers and polynomials.

2. Identities on the Frobenius-Euler Numbers

From (1.2) and (1.4), we can derive the following:

$$\int_{\mathbb{Z}_p} e^{(x+y)t} d\mu_{-q}(y) = \frac{1+q^{-1}}{e^t + q^{-1}} e^{xt} = \sum_{n=0}^{\infty} H_n(-q^{-1}, x) \frac{t^n}{n!}. \quad (2.1)$$

Thus, by (2.1), we get Witt's formula for $H_n(-q^{-1}, x)$ as follows:

$$\int_{\mathbb{Z}_p} (x+y)^n d\mu_{-q}(y) = H_n(-q^{-1}, x), \quad n \in \mathbb{Z}_+. \quad (2.2)$$

By (1.5) and (2.1), we get

$$(H(-q^{-1}) + 1)^n + q^{-1} H_n(-q^{-1}) = \begin{cases} 1 + q^{-1}, & \text{if } n = 0, \\ 0, & \text{if } n > 0, \end{cases} \quad (2.3)$$

with the usual convention about replacing $H(-q^{-1})^n$ by $H_n(-q^{-1})$.

From (1.5) and (2.3), we note that

$$H_0(-q^{-1}) = 1, \quad H_n(-q^{-1}, 1) + q^{-1}H_n(-q^{-1}) = 0, \quad \text{if } n > 0. \quad (2.4)$$

By (2.1) and (2.2), we get

$$\int_{\mathbb{Z}_p} (\mathbf{y} + 1 - x)^n d\mu_{-q}(\mathbf{y}) = (-1)^n \int_{\mathbb{Z}_p} (\mathbf{y} + x)^n d\mu_{-q^{-1}}(\mathbf{y}). \quad (2.5)$$

Therefore, by (2.5), we obtain the following lemma.

Lemma 2.1. For $n \in \mathbb{Z}_+$, one has

$$H_n(-q^{-1}, 1 - x) = (-1)^n H_n(-q, x). \quad (2.6)$$

From (2.3), we can derive the following:

$$\begin{aligned} q^2 H_n(-q^{-1}, 2) - q^2 - q &= q^2 \sum_{l=0}^n \binom{n}{l} (H(-q^{-1}) + 1)^l - q^2 - q \\ &= q \sum_{l=1}^n \binom{n}{l} q (H(-q^{-1}) + 1)^l - q \\ &= -q \sum_{l=0}^n \binom{n}{l} H_l(-q^{-1}) \\ &= -(1 + q)\delta_{0,n} + H_n(-q^{-1}), \end{aligned} \quad (2.7)$$

where $\delta_{k,n}$ is the Kronecker symbol.

Therefore, by (2.7), we obtain the following theorem.

Theorem 2.2. For $n \in \mathbb{Z}_+$, one has

$$H_n(-q^{-1}, 2) = 1 + q^{-1} - q^{-2}(1 + q)\delta_{0,n} + q^{-2}H_n(-q^{-1}). \quad (2.8)$$

First we consider the fermionic p -adic q -integral on \mathbb{Z}_p for the n th Frobenius-Euler polynomials as follows:

$$\begin{aligned}
 I_1 &= \int_{\mathbb{Z}_p} H_n(-q^{-1}, x) d\mu_{-q}(x) \\
 &= \sum_{l=0}^n \binom{n}{l} H_l(-q^{-1}) \int_{\mathbb{Z}_p} x^{n-l} d\mu_{-q}(x) \\
 &= \sum_{l=0}^n \binom{n}{l} H_l(-q^{-1}) H_{n-l}(-q^{-1}), \quad \text{where } n \in \mathbb{Z}_+.
 \end{aligned} \tag{2.9}$$

On the other hand, by (2.5) and Lemma 2.1, we get

$$\begin{aligned}
 I_1 &= \int_{\mathbb{Z}_p} H_n(-q^{-1}, x) d\mu_{-q}(x) = (-1)^n \int_{\mathbb{Z}_p} H_n(-q, 1-x) d\mu_{-q}(x) \\
 &= (-1)^n \sum_{l=0}^n \binom{n}{l} H_{n-l}(-q) \int_{\mathbb{Z}_p} (1-x)^l d\mu_{-q}(x) \\
 &= \sum_{l=0}^n \binom{n}{l} (-1)^{n-l} H_{n-l}(-q) \int_{\mathbb{Z}_p} (x-1)^l d\mu_{-q}(x) \\
 &= \sum_{l=0}^n \binom{n}{l} (-1)^{n-l} H_{n-l}(-q) H_l(-q^{-1}, -1).
 \end{aligned} \tag{2.10}$$

From Lemma 2.1, Theorem 2.2, and (2.10), we note that

$$\begin{aligned}
 I_1 &= \sum_{l=0}^n \binom{n}{l} (-1)^{n-l} H_{n-l}(-q) H_l(-q^{-1}, -1) \\
 &= \sum_{l=0}^n \binom{n}{l} (-1)^n H_{n-l}(-q) H_l(-q, 2) \\
 &= \sum_{l=0}^n \binom{n}{l} (-1)^n H_{n-l}(-q) \{1 + q - q^2(1 + q^{-1})\delta_{0,l} + q^2 H_l(-q)\} \\
 &= (-1)^n (1+q) ((1+q)\delta_{0,n} - qH_n(-q)) - H_n(-q)(q+q^2)(-1)^n \\
 &\quad + (-1)^n q^2 \sum_{l=0}^n \binom{n}{l} H_{n-l}(-q) H_l(-q).
 \end{aligned} \tag{2.11}$$

Therefore, by (2.10) and (2.11), we obtain the following theorem.

Theorem 2.3. For $n \in \mathbb{Z}_+$, one has

$$\begin{aligned} \sum_{l=0}^n \binom{n}{l} H_l(-q^{-1}) H_{n-l}(-q^{-1}) &= (-1)^n (1+q) ((1+q)\delta_{0,n} - 2qH_n(-q)) \\ &+ (-1)^n q^2 \sum_{l=0}^n \binom{n}{l} H_{n-l}(-q) H_l(-q). \end{aligned} \tag{2.12}$$

In particular, for $n \in \mathbb{N}$, one has

$$\begin{aligned} \sum_{l=0}^n \binom{n}{l} H_l(-q^{-1}) H_{n-l}(-q^{-1}) &= 2(-1)^{n+1} q(1+q) H_n(-q) \\ &+ (-1)^n q^2 \sum_{l=0}^n \binom{n}{l} H_{n-l}(-q) H_l(-q). \end{aligned} \tag{2.13}$$

Let us consider the following fermionic p -adic q -integral on \mathbb{Z}_p for the product of Bernoulli and Frobenius-Euler polynomials as follows:

$$\begin{aligned} I_2 &= \int_{\mathbb{Z}_p} B_m(x) H_n(-q^{-1}, x) d\mu_{-q}(x) \\ &= \sum_{k=0}^m \sum_{l=0}^n \binom{m}{k} \binom{n}{l} B_{m-k} H_{n-l}(-q^{-1}) \int_{\mathbb{Z}_p} x^{k+l} d\mu_{-q}(x) \\ &= \sum_{k=0}^m \sum_{l=0}^n \binom{m}{k} \binom{n}{l} B_{m-k} H_{n-l}(-q^{-1}) H_{k+l}(-q^{-1}). \end{aligned} \tag{2.14}$$

It is known that $B_n(x) = (-1)^n B_n(1-x)$.

On the other hand, by Lemma 2.1, we get

$$\begin{aligned} I_2 &= (-1)^{m+n} \int_{\mathbb{Z}_p} B_m(1-x) H_n(-q, 1-x) d\mu_{-q}(x) \\ &= (-1)^{m+n} \sum_{k=0}^m \sum_{l=0}^n \binom{m}{k} \binom{n}{l} B_{m-k} H_{n-l}(-q) \int_{\mathbb{Z}_p} (1-x)^{k+l} d\mu_{-q}(x) \\ &= (-1)^{m+n} \sum_{k=0}^m \sum_{l=0}^n \binom{m}{k} \binom{n}{l} B_{m-k} H_{n-l}(-q) \{ (1+q) - q^2(1+q^{-1})\delta_{0,k+l} + q^2 H_{k+l}(-q) \} \\ &= (-1)^{m+n} (1+q) B_m(1) H_n(-q, 1) - (q^2+q) (-1)^{m+n} B_m H_n(-q) \\ &\quad + q^2 (-1)^{m+n} \sum_{k=0}^m \sum_{l=0}^n \binom{m}{k} \binom{n}{l} B_{m-k} H_{n-l}(-q) H_{k+l}(-q). \end{aligned} \tag{2.15}$$

Therefore, by (2.14) and (2.15), we obtain the following theorem.

Theorem 2.4. For $m, n \in \mathbb{Z}_+$, one has

$$\begin{aligned} & \sum_{k=0}^m \sum_{l=0}^n \binom{m}{k} \binom{n}{l} B_{m-k} H_{n-l}(-q^{-1}) H_{k+l}(-q^{-1}) \\ &= (-1)^{m+n} (1+q) B_m(1) H_n(-q, 1) - (q^2 + q) (-1)^{m+n} B_m H_n(-q) \\ & \quad + q^2 (-1)^{m+n} \sum_{k=0}^m \sum_{l=0}^n \binom{m}{k} \binom{n}{l} B_{m-k} H_{n-l}(-q) H_{k+l}(-q). \end{aligned} \quad (2.16)$$

In particular, for $m \in \mathbb{N} - \{1\}$, $n \in \mathbb{N}$, one has

$$\begin{aligned} & \sum_{k=0}^m \sum_{l=0}^n \binom{m}{k} \binom{n}{l} B_{m-k} H_{n-l}(-q^{-1}) H_{k+l}(-q^{-1}) \\ &= 2(-1)^{m+n+1} (q^2 + q) B_m H_n(-q) \\ & \quad + q^2 (-1)^{m+n} \sum_{k=0}^m \sum_{l=0}^n \binom{m}{k} \binom{n}{l} B_{m-k} H_{n-l}(-q) H_{k+l}(-q). \end{aligned} \quad (2.17)$$

It is known that $x^n = (1/(n+1)) \sum_{l=0}^n \binom{n+1}{l} B_l(x)$. Let us consider the following fermionic p -adic q -integral on \mathbb{Z}_p :

$$\begin{aligned} \int_{\mathbb{Z}_p} x^n d\mu_{-q}(x) &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} \int_{\mathbb{Z}_p} B_l(x) d\mu_{-q}(x). \\ &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} \sum_{k=0}^l \binom{l}{k} B_{l-k} \int_{\mathbb{Z}_p} x^k d\mu_{-q}(x) \\ &= \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} \sum_{k=0}^l \binom{l}{k} B_{l-k} H_k(-q^{-1}). \end{aligned} \quad (2.18)$$

Therefore by (2.18), we obtain the following theorem.

Theorem 2.5. For $n \in \mathbb{Z}_+$, one has

$$H_n(-q^{-1}) = \frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} \sum_{k=0}^l \binom{l}{k} B_{l-k} H_k(-q^{-1}). \tag{2.19}$$

From (1.3), we can derive the following:

$$x^n = E_n(x) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} E_l(x). \tag{2.20}$$

Let us take the fermionic p -adic q -integral on \mathbb{Z}_p in (2.20) as follows:

$$\begin{aligned} \int_{\mathbb{Z}_p} x^n d\mu_{-q}(x) &= \int_{\mathbb{Z}_p} E_n(x) d\mu_{-q}(x) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} \int_{\mathbb{Z}_p} E_l(x) d\mu_{-q}(x) \\ &= \sum_{l=0}^n \binom{n}{l} E_{n-l} H_l(-q^{-1}) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} \sum_{k=0}^l \binom{l}{k} E_{l-k} H_k(-q^{-1}). \end{aligned} \tag{2.21}$$

Therefore, by (2.21), we obtain the following theorem.

Theorem 2.6. For $n \in \mathbb{N}$, one has

$$H_n(-q^{-1}) = \sum_{l=0}^n \binom{n}{l} E_{n-l} H_l(-q^{-1}) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} \sum_{k=0}^l \binom{l}{k} E_{l-k} H_k(-q^{-1}). \tag{2.22}$$

By Theorems 2.5 and 2.6, we obtain the following corollary.

Corollary 2.7. For $n \in \mathbb{N}$, one has

$$\begin{aligned} &\frac{1}{n+1} \sum_{l=0}^n \binom{n+1}{l} \sum_{k=0}^l \binom{l}{k} B_{l-k} H_k(-q^{-1}) \\ &= \sum_{l=0}^n \binom{n}{l} E_{n-l} H_l(-q^{-1}) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} \sum_{k=0}^l \binom{l}{k} E_{l-k} H_k(-q^{-1}). \end{aligned} \tag{2.23}$$

By (1.3), we easily get $E_n(x) = (-1)^n E_n(1-x)$.

Thus, we have

$$\begin{aligned}
 & \int_{\mathbb{Z}_p} x^n d\mu_{-q}(x) \\
 &= (-1)^n \int_{\mathbb{Z}_p} E_n(1-x) d\mu_{-q}(x) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} (-1)^l \int_{\mathbb{Z}_p} E_l(1-x) d\mu_{-q}(x) \\
 &= (-1)^n \sum_{l=0}^n \binom{n}{l} E_{n-l} \int_{\mathbb{Z}_p} (1-x)^l d\mu_{-q}(x) \\
 &\quad + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} (-1)^l \sum_{k=0}^l \binom{l}{k} E_{l-k} \int_{\mathbb{Z}_p} (1-x)^k d\mu_{-q}(x) \\
 &= \sum_{l=0}^n \binom{n}{l} E_{n-l} (-1)^{n-l} H_l(-q^{-1}, -1) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} \sum_{k=0}^l \binom{l}{k} E_{l-k} (-1)^{l-k} H_k(-q^{-1}, -1) \\
 &= \sum_{l=0}^n \binom{n}{l} E_{n-l} (-1)^n H_l(-q, 2) + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} \sum_{k=0}^l \binom{l}{k} E_{l-k} (-1)^l H_k(-q, 2).
 \end{aligned} \tag{2.24}$$

Therefore, by (2.24), we obtain the following theorem.

Theorem 2.8. For $n \in \mathbb{N}$, one has

$$\begin{aligned}
 H_n(-q^{-1}) &= \sum_{l=0}^n \binom{n}{l} E_{n-l} (-1)^n H_l(-q, 2) \\
 &\quad + \frac{1}{2} \sum_{l=0}^{n-1} \binom{n}{l} \sum_{k=0}^l \binom{l}{k} E_{l-k} (-1)^l H_k(-q, 2).
 \end{aligned} \tag{2.25}$$




Acknowledgment

The second author was supported by National Research Foundation of Korea Grant funded by the Korean Government 2009-0072514.

References

- [1] T. Kim, "Some identities on the q -Euler polynomials of higher order and q -Stirling numbers by the fermionic p -adic integral on \mathbb{Z}_p ," *Russian Journal of Mathematical Physics*, vol. 16, no. 4, pp. 484–491, 2009.
- [2] L. Carlitz, "The product of two eulerian polynomials," *Mathematics Magazine*, vol. 36, no. 1, pp. 37–41, 1963.
- [3] A. Bayad, "Modular properties of elliptic Bernoulli and Euler functions," *Advanced Studies in Contemporary Mathematics*, vol. 20, no. 3, pp. 389–401, 2010.
- [4] C.-P. Chen and L. Lin, "An inequality for the generalized-Euler-constant function," *Advanced Studies in Contemporary Mathematics*, vol. 17, no. 1, pp. 105–107, 2008.
- [5] T. Kim, J. Choi, and Y. H. Kim, "A note on Bernstein polynomials associated with q -Euler numbers and polynomials," *International Journal of Mathematics and Analysis*, vol. 5, no. 48, pp. 2413–2420, 2011.

- [6] H. Ozden, I. N. Cangul, and Y. Simsek, "Remarks on q -Bernoulli numbers associated with Daehee numbers," *Advanced Studies in Contemporary Mathematics*, vol. 18, no. 1, pp. 41–48, 2009.
- [7] S.-H. Rim, S. J. Lee, E. J. Moon, and J. H. Jin, "On the q -Genocchi numbers and polynomials associated with q -zeta function," *Proceedings of the Jangjeon Mathematical Society*, vol. 12, no. 3, pp. 261–267, 2009.
- [8] C. S. Ryoo, "A note on the weighted q -Euler numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 21, pp. 47–54, 2011.
- [9] C. S. Ryoo, "On the generalized Barnes type multiple q -Euler polynomials twisted by ramified roots of unity," *Proceedings of the Jangjeon Mathematical Society*, vol. 13, no. 2, pp. 255–263, 2010.
- [10] Y. Simsek, "Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions," *Advanced Studies in Contemporary Mathematics*, vol. 16, no. 2, pp. 251–278, 2008.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

