

Comparing two matrices of generalized moments defined by continued fraction expansions

Paul Barry
 School of Science
 Waterford Institute of Technology
 Ireland
 pbarry@wit.ie

Abstract

We study two matrices N and M defined by the parameters of equivalent S - and J -continued fraction expansions, and compare them by examining the product $N^{-1}M$. Using examples based on the Catalan numbers, the little Schröder numbers and powers of q , we indicate that this matrix product is an object worthy of study. In the case of the little Schröder numbers, we find that the matrix N has an interleaved structure based on two Riordan arrays.

1 Introduction

In this note, we study two matrices whose elements may be considered to be generalized moments. The matrices are defined using the coefficients of simple Jacobi and Stieltjes continued fractions.

In familiar cases, these matrices are well-known, though this study examines them from a fresh perspective. It will be assumed that the reader is familiar with the basics of orthogonal polynomials [4, 8, 17], Riordan arrays [14], production matrices [6, 7, 13], continued fractions [18] and the interplay between these areas [1, 2].

Our point of departure is a sequence a_n , with $a_1 = 1$, whose elements are either integers or polynomials with integer coefficients.

We will use these numbers to define two lower-triangular matrices, which we then compare.

In both cases, the elements of the first column will be the sequence μ_n generated by the continued fraction

$$\frac{1}{1 - \frac{a_1 x}{1 - \frac{a_2 x}{1 - \frac{a_3 x}{1 - \dots}}}}$$

We require that this sequence be Catalan-like, in the sense that we require all the Hankel determinants $|\mu_{i+j}|_{0 \leq i, j \leq n}$ to be non-zero.

An equivalence transformation ensures that the sequence μ_n is the same as that generated by

$$\frac{1}{1 - a_1x - \frac{a_1a_2x^2}{1 - (a_2 + a_3)x - \frac{a_3a_4x^2}{1 - (a_4 + a_5)x - \dots}}}$$

This exhibits μ_n as the moment sequence of the family of orthogonal polynomials $P_n(x)$ that satisfy

$$P_n(x) = (x - (a_{2n-2} + a_{2n-1}))P_{n-1}(x) - a_{2n-3}a_{2n-2}P_{n-2}(x),$$

with $P_0(x) = 1$, $P_1(x) = x - a_1$.

The first matrix M that we shall be interested in is the inverse of the matrix of coefficients of these polynomials. This matrix therefore has production matrix given by

$$\begin{pmatrix} a_1 & 1 & 0 & 0 & 0 & 0 & \dots \\ a_1a_2 & a_2 + a_3 & 1 & 0 & 0 & 0 & \dots \\ 0 & a_3a_4 & a_4 + a_5 & 1 & 0 & 0 & \dots \\ 0 & 0 & a_5a_6 & a_6 + a_7 & 1 & 0 & \dots \\ 0 & 0 & 0 & a_7a_8 & a_8 + a_9 & 1 & \dots \\ 0 & 0 & 0 & 0 & a_9a_{10} & a_{10} + a_{11} & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

The form of this production matrix ensures that the matrix generated by it will be lower-triangular with 1's on the diagonal. We obtain a matrix which begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ \alpha & 1 & 0 & 0 & 0 \\ \alpha(\alpha + \beta) & \alpha + \beta + \gamma & 1 & 0 & 0 \\ \alpha((\alpha + \beta)^2 + \beta\gamma) & (\alpha + \beta)^2 + \beta\gamma + \gamma(\alpha + \beta + \gamma + \delta) & \alpha + \beta + \gamma + \delta + \epsilon & 1 & 0 \\ \alpha((\alpha + \beta)^3 + \beta\gamma(2\alpha + 2\beta + \gamma + \delta)) & (\alpha + \beta + \gamma + \delta + \epsilon)^2 - \alpha(\gamma + \delta + \epsilon) - \beta(\delta + \epsilon) - \epsilon(\gamma - \phi) & \dots & \alpha + \beta + \gamma + \delta + \epsilon + \phi & 1 \\ \vdots & \vdots & \vdots & \vdots & \vdots \end{pmatrix}$$

where we have written $\alpha = a_1$, $\beta = a_2$, and so on.

In order to define the second matrix N , which again will have μ_n in the first column, we also use a production matrix. To construct this production matrix, we have two alternative routes. The first one proceeds as follows; we take the inverse of the matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ -a_1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & -a_2 & 1 & 0 & 0 & 0 & \dots \\ 0 & 0 & -a_3 & 1 & 0 & 0 & \dots \\ 0 & 0 & 0 & -a_4 & 1 & 0 & \dots \\ 0 & 0 & 0 & 0 & -a_5 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

to obtain the matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ a_1 & 1 & 0 & 0 & 0 & 0 & \dots \\ a_1 a_2 & a_2 & 1 & 0 & 0 & 0 & \dots \\ a_1 a_2 a_3 & a_2 a_3 & a_3 & 1 & 0 & 0 & \dots \\ a_1 a_2 a_3 a_4 & a_2 a_3 a_4 & a_3 a_4 & a_4 & 1 & 0 & \dots \\ a_1 a_2 a_3 a_4 a_5 & a_2 a_3 a_4 a_5 & a_3 a_4 a_5 & a_4 a_5 & a_5 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

We now behead this matrix (we remove the first row) to obtain the following production matrix.

$$\begin{pmatrix} a_1 & 1 & 0 & 0 & 0 & 0 & \dots \\ a_1 a_2 & a_2 & 1 & 0 & 0 & 0 & \dots \\ a_1 a_2 a_3 & a_2 a_3 & a_3 & 1 & 0 & 0 & \dots \\ a_1 a_2 a_3 a_4 & a_2 a_3 a_4 & a_3 a_4 & a_4 & 1 & 0 & \dots \\ a_1 a_2 a_3 a_4 a_5 & a_2 a_3 a_4 a_5 & a_3 a_4 a_5 & a_4 a_5 & a_5 & 1 & \dots \\ a_1 a_2 a_3 a_4 a_5 a_6 & a_2 a_3 a_4 a_5 a_6 & a_3 a_4 a_5 a_6 & a_4 a_5 a_6 & a_5 a_6 & a_6 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

The form of this production matrix ensures that the matrix N that it generates will be lower-triangular with 1's on the diagonal. The matrix N that we seek then begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & \dots \\ \alpha & 1 & 0 & 0 & 0 & \dots \\ \alpha(\alpha + \beta) & \alpha + \beta & 1 & 0 & 0 & \dots \\ \alpha((\alpha + \beta)^2 + \beta\gamma) & (\alpha + \beta)^2 + \beta\gamma & \alpha + \beta + \gamma & 1 & 0 & \dots \\ \alpha((\alpha + \beta)^3 + \beta\gamma(2\alpha + 2\beta + \gamma + \delta)) & (\alpha + \beta)^3 + \beta\gamma(2\alpha + 2\beta + \gamma + \delta) & (\alpha + \beta)^2 + \gamma(\alpha + \delta) + (\beta + \gamma)^2 & \alpha + \beta + \gamma + \delta & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

where we have used $\alpha = a_1$, $\beta = a_2$, and so on.

There is an alternative production matrix approach to the construction of N . Multiplying the (n, k) -th element of N by

$$\prod_{j=1}^k a_j$$

produces a lower triangular matrix whose first column is the same as that of N , and whose production matrix takes the simple form of

$$\begin{pmatrix} a_1 & a_1 & 0 & 0 & 0 & 0 & \dots \\ a_2 & a_2 & a_2 & 0 & 0 & 0 & \dots \\ a_3 & a_3 & a_3 & a_3 & 0 & 0 & \dots \\ a_4 & a_4 & a_4 & a_4 & a_4 & 0 & \dots \\ a_5 & a_5 & a_5 & a_5 & a_5 & a_5 & \dots \\ a_6 & a_6 & a_6 & a_6 & a_6 & a_6 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

We can clearly reverse this process, starting with the sequence a_n , to produce N .

In order to compare the two matrices M and N , it is natural to examine the product $N^{-1}M$.

Example 1. The Catalan matrices. We let $a_n = 1$. Thus we are interested in the sequence generated by the continued fraction

$$\frac{1}{1 - \frac{x}{1 - \frac{x}{1 - \frac{x}{\dots}}}}$$

This is the sequence of Catalan numbers $C_n = \frac{1}{n+1} \binom{2n}{n}$, [A000108](#). In this instance, the production matrix of N for both methods of generation is given by

$$\begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 1 & 0 & 0 & 0 & \dots \\ 1 & 1 & 1 & 1 & 0 & 0 & \dots \\ 1 & 1 & 1 & 1 & 1 & 0 & \dots \\ 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ 1 & 1 & 1 & 1 & 1 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

and N is the Riordan array

$$N = (c(x), xc(x)) = (1 - x, x(1 - x))^{-1} \quad \text{\a href="#">A033184.$$

The matrix M is given by the Riordan array

$$M = (c(x), xc(x)^2) = \left(\frac{1}{1+x}, \frac{x}{(1+x)^2} \right)^{-1} \quad \text{\a href="#">A039599,$$

and the associated orthogonal polynomials are the Chebyshev polynomials $U_n(\frac{x}{2})$. The production matrix of $(c(x), xc(x)^2)$ is given by

$$\begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 1 & 2 & 1 & 0 & 0 & 0 & \dots \\ 0 & 1 & 2 & 1 & 0 & 0 & \dots \\ 0 & 0 & 1 & 2 & 1 & 0 & \dots \\ 0 & 0 & 0 & 1 & 2 & 1 & \dots \\ 0 & 0 & 0 & 0 & 1 & 2 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

corresponding to the generating function

$$\frac{1}{1 - x - \frac{x^2}{1 - 2x - \frac{x^2}{1 - 2x - \dots}}}$$

of C_n .

A straight-forward Riordan array calculation now shows that in this case,

$$N^{-1} \cdot M = (c(x), xc(x))^{-1} \cdot (c(x), xc(x)^2) = \left(1, \frac{x}{1-x}\right),$$

which is the shifted binomial matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 1 & 0 & 0 & 0 & \dots \\ 0 & 1 & 2 & 1 & 0 & 0 & \dots \\ 0 & 1 & 3 & 3 & 1 & 0 & \dots \\ 0 & 1 & 4 & 6 & 4 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Example 2. The q -case. We take the example of

$$a_n = \frac{q^n}{q} - \frac{0^n}{q}.$$

Starting with the matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ -q & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & -q^2 & 1 & 0 & 0 & 0 & \dots \\ 0 & 0 & -q^3 & 1 & 0 & 0 & \dots \\ 0 & 0 & 0 & -q^4 & 1 & 0 & \dots \\ 0 & 0 & 0 & 0 & -q^5 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

we invert it and behold the resulting matrix to get the production matrix

$$\begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ q & q & 1 & 0 & 0 & 0 & \dots \\ q^3 & q^3 & q^2 & 1 & 0 & 0 & \dots \\ q^6 & q^6 & q^5 & q^3 & 1 & 0 & \dots \\ q^{10} & q^{10} & q^9 & q^7 & q^4 & 1 & \dots \\ q^{15} & q^{15} & q^{14} & q^{12} & q^9 & q^5 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

which we use to generate the matrix N:

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & 0 & \dots \\ q+1 & q+1 & 1 & 0 & 0 & \dots \\ q^3+q^2+2q+1 & q^3+q^2+2q+1 & q^2+q+1 & 1 & 0 & \dots \\ q^6+q^5+2q^4+3q^3+3q^2+3q+1 & q^6+q^5+2q^4+3q^3+3q^2+3q+1 & q^5+q^4+2q^3+2q^2+2q+1 & q^3+q^2+q+1 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

In the left column we recognize the q -Catalan sequence μ_n with generating function

$$\frac{1}{1 - \frac{x}{1 - \frac{qx}{1 - \frac{q^2x}{1 - \dots}}}}.$$

Alternatively we may begin with the production matrix

$$\begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ q & q & q & 0 & 0 & 0 & \dots \\ q^2 & q^2 & q^2 & q^2 & 0 & 0 & \dots \\ q^3 & q^3 & q^3 & q^3 & q^3 & 0 & \dots \\ q^4 & q^4 & q^4 & q^4 & q^4 & q^4 & \dots \\ q^5 & q^5 & q^5 & q^5 & q^5 & q^5 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Let \tilde{N} be the matrix generated by this production matrix. Dividing column k of the \tilde{M} by

$$\prod_{i=1}^k a_i = \prod_{i=1}^k q^i = q^{\binom{k}{2}},$$

we recover the matrix N .

The generating function

$$\frac{1}{1 - \frac{x}{1 - \frac{qx}{1 - \frac{q^2x}{1 - \dots}}}}.$$

is equivalent to

$$\frac{1}{1 - x - \frac{qx^2}{1 - (q + q^2)x - \frac{q^5x^2}{1 - (q^3 + q^4)x - \frac{q^9x^2}{1 - (q^5 + q^6)x - \dots}}}}.$$

This leads to the production matrix

$$\begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ q & q + q^2 & 1 & 0 & 0 & 0 & \dots \\ 0 & q^5 & q^3 + q^4 & 1 & 0 & 0 & \dots \\ 0 & 0 & q^9 & q^5 + q^6 & 1 & 0 & \dots \\ 0 & 0 & 0 & q^{13} & q^7 + q^8 & 1 & \dots \\ 0 & 0 & 0 & 0 & q^{17} & q^9 + q^{10} & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

which generates the matrix M , with first column equal to μ_n . The inverse of the matrix M is the coefficient array of the orthogonal polynomials defined by

$$P_n(x) = (x - q^{2n-3}(1+q))P_{n-1}(x) - q^{4n-7}P_{n-2}(x),$$

where $P_0(x) = 1$ and $P_1(x) = x - 1$.

For $N^{-1} \cdot M$, we obtain the matrix that begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & q^2 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & q^5 & q^3 + q^4 & 1 & 0 & 0 & 0 & \dots \\ 0 & q^9 & q^7 + q^8 + q^9 & q^4 + q^5 + q^6 & 1 & 0 & 0 & \dots \\ 0 & q^{14} & q^{12} + q^{13} + q^{14} + q^{15} & q^9 + q^{10} + q^{11} + q^{12} + q^{13} & q^5 + q^6 + q^7 + q^8 & 1 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Dividing each element $M_{n,k}$ of M by

$$q^{\binom{n-k+2}{2}-1},$$

we obtain the matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & q(1+q) & 1 & 0 & 0 & 0 & \dots \\ 0 & 1 & q^2(1+q+q^2) & q^2(1+q+q^2) & 1 & 0 & 0 & \dots \\ 0 & 1 & q^3(1+q+q^2+q^3) & q^4(1+q+2q^2+q^3+q^4) & q^3(1+q+q^2+q^3) & 1 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

which is the Hadamard product of the matrices

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 1+q & 1 & 0 & 0 & 0 & \dots \\ 0 & 1 & 1+q+q^2 & 1+q+q^2 & 1 & 0 & 0 & \dots \\ 0 & 1 & 1+q+q^2+q^3 & 1+q+2q^2+q^3+q^4 & 1+q+q^2+q^3 & 1 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

and

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 1 & 0 & 0 & 0 & \dots \\ 0 & 1 & q & 1 & 0 & 0 & \dots \\ 0 & 1 & q^2 & q^2 & 1 & 0 & \dots \\ 0 & 1 & q^3 & q^4 & q^3 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

where the first matrix is the q -Riordan array $\begin{bmatrix} n-1 \\ n-k \end{bmatrix}_q$ [3], and the second matrix is a shifted version of the matrix $q^{k(n-k)}$.

The production matrix of $N^{-1} \cdot M$ begins

$$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & q^2 & 1 & 0 & 0 & 0 & \dots \\ 0 & q^5(q-1) & q^2(q^2+q-1) & 1 & 0 & 0 & \dots \\ 0 & 0 & q^7(q^2-1) & q^3(q^3+q^2-1) & 1 & 0 & \dots \\ 0 & 0 & 0 & q^{10}(q^3-1) & q^4(q^4+q^3-1) & 1 & \dots \\ 0 & 0 & 0 & 0 & q^{13}(q^4-1) & q^5(q^5+q^4-1) & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

indicating that in this case, the inverse matrix $(N^{-1} \cdot M)^{-1} = M^{-1} \cdot N$ is the coefficient array of a family of orthogonal polynomials whose parameters are given by the production matrix above.

We look more closely at the case of $q = 2$. We find that

$$M = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 3 & 7 & 1 & 0 & 0 & 0 & \dots \\ 17 & 77 & 31 & 1 & 0 & 0 & \dots \\ 171 & 1471 & 1333 & 127 & 1 & 0 & \dots \\ 3113 & 51653 & 98487 & 21717 & 511 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

while

$$N = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 3 & 3 & 1 & 0 & 0 & 0 & \dots \\ 17 & 17 & 7 & 1 & 0 & 0 & \dots \\ 171 & 171 & 77 & 51 & 1 & 0 & \dots \\ 3113 & 3113 & 1471 & 325 & 31 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Then

$$N^{-1} \cdot M = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 4 & 1 & 0 & 0 & 0 & \dots \\ 0 & 32 & 24 & 1 & 0 & 0 & \dots \\ 0 & 512 & 896 & 112 & 1 & 0 & \dots \\ 0 & 16384 & 61440 & 17920 & 480 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Looking at the reduced matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 4 & 1 & 0 & 0 & 0 & 0 & \dots \\ 32 & 24 & 1 & 0 & 0 & 0 & \dots \\ 512 & 896 & 112 & 1 & 0 & 0 & \dots \\ 16384 & 61440 & 17920 & 480 & 1 & 0 & \dots \\ 1048576 & 8126464 & 5079040 & 317440 & 1984 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

we see that it is the moment array of the family of orthogonal polynomials whose parameters are given in the production matrix

$$\begin{pmatrix} 4 & 1 & 0 & 0 & 0 & 0 & \dots \\ 16 & 20 & 1 & 0 & 0 & 0 & \dots \\ 0 & 384 & 88 & 1 & 0 & 0 & \dots \\ 0 & 0 & 7168 & 368 & 1 & 0 & \dots \\ 0 & 0 & 0 & 122880 & 1504 & 1 & \dots \\ 0 & 0 & 0 & 0 & 2031616 & 6080 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

We deduce that the sequence $1, 4, 32, 512, 16384, \dots$ or $2^{n(n+3)/2}$ [A036442](#) has a generating function given by

$$\frac{1}{1 - 4x - \frac{16x^2}{1 - 20x - \frac{384x^2}{1 - 88x - \frac{7168x^2}{1 - 368x - \dots}}}},$$

or equivalently,

$$\frac{1}{1 - \frac{4x}{1 - \frac{4x}{1 - \frac{16x}{1 - \frac{24x}{1 - \frac{64x}{1 - \dots}}}}}}.$$

In this latter expression, the coefficients are given by the sequence

$$b(n) = 2^{n+2} - 2^{(n+1)/2}(1 - (-1)^n).$$

The Hankel transform of $2^{n(n+3)/2}$ is then given by [\[10, 11, 12\]](#)

$$h_n = \prod_{k=0}^{n-1} (b(2k+1)b(2k+2))^{n-k}.$$

A similar analysis can be carried out for $q^{n(n+3)/2}$.

Example 3. The little Schröder numbers. In this example, we take a base sequence a_n given by

$$1, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, 1, 2, \dots$$

The sequence with generating function

$$\frac{1}{1 - \frac{a_1 x}{1 - \frac{a_2 x}{1 - \frac{a_3 x}{1 - \dots}}}} = \frac{1}{1 - \frac{x}{1 - \frac{2x}{1 - \frac{x}{1 - \dots}}}}$$

is the sequence of little Schröder numbers [A001003](#)

$$1, 1, 3, 11, 45, 197, 903, \dots$$

These numbers are also generated by

$$\frac{1}{1 - x - \frac{2x^2}{1 - 3x - \frac{2x^2}{1 - 3x - \dots}}}$$

We have

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ -1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & -2 & 1 & 0 & 0 & 0 & \dots \\ 0 & 0 & -1 & 1 & 0 & 0 & \dots \\ 0 & 0 & 0 & -2 & 1 & 0 & \dots \\ & 0 & 0 & 0 & -1 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 2 & 2 & 1 & 0 & 0 & 0 & \dots \\ 2 & 2 & 1 & 1 & 0 & 0 & \dots \\ 4 & 4 & 2 & 2 & 1 & 0 & \dots \\ 4 & 4 & 2 & 2 & 1 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

so that the matrix N in this case begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 3 & 3 & 1 & 0 & 0 & 0 & \dots \\ 11 & 11 & 4 & 1 & 0 & 0 & \dots \\ 45 & 45 & 17 & 6 & 1 & 0 & \dots \\ 197 & 197 & 76 & 31 & 7 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

with production matrix

$$\begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 2 & 2 & 1 & 0 & 0 & 0 & \dots \\ 2 & 2 & 1 & 1 & 0 & 0 & \dots \\ 4 & 4 & 2 & 2 & 1 & 0 & \dots \\ 4 & 4 & 2 & 2 & 1 & 1 & \dots \\ 8 & 8 & 4 & 4 & 2 & 2 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

For instance, we have

$$17 = 1.11 + 1.4 + 2.1 + 2.0 + \dots,$$

and

$$45 = 1.11 + 2.11 + 2.4 + 4.1 + \dots.$$

The matrix M is given by

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 3 & 4 & 1 & 0 & 0 & 0 & \dots \\ 11 & 17 & 7 & 1 & 0 & 0 & \dots \\ 45 & 76 & 40 & 10 & 1 & 0 & \dots \\ 197 & 353 & 216 & 72 & 13 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

This is the Riordan array [A172094](#)

$$\left(\frac{1+x-\sqrt{1-6x+x^2}}{4x}, \frac{1-3x-\sqrt{1-4x+x^2}}{4x} \right) = \left(\frac{1}{1+x}, \frac{x}{1+3x+2x^2} \right)^{-1},$$

with production matrix

$$\begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 2 & 3 & 1 & 0 & 0 & 0 & \dots \\ 0 & 2 & 3 & 1 & 0 & 0 & \dots \\ 0 & 0 & 2 & 3 & 1 & 0 & \dots \\ 0 & 0 & 0 & 2 & 3 & 1 & \dots \\ 0 & 0 & 0 & 0 & 2 & 3 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

The matrix N is a “mixture” (in left to right interleaved fashion) [5] of this Riordan array and the related Riordan array

$$\left(1, \frac{1-3x-\sqrt{1-6x+x^2}}{4x} \right) = \left(1, \frac{x}{1+3x+2x^2} \right)^{-1},$$

which has production matrix

$$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 3 & 1 & 0 & 0 & 0 & \dots \\ 0 & 2 & 3 & 1 & 0 & 0 & \dots \\ 0 & 0 & 2 & 3 & 1 & 0 & \dots \\ 0 & 0 & 0 & 2 & 3 & 1 & \dots \\ 0 & 0 & 0 & 0 & 2 & 3 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

We have

$$N_{n,k} = \begin{cases} [x^n] \frac{1+x-\sqrt{1-6x+x^2}}{4x} \left(\frac{1-3x-\sqrt{1-6x+x^2}}{4} \right)^{k/2}, & \text{if } k \text{ is even} \\ [x^n] x^k \left(\frac{1-3x-\sqrt{1-6x+x^2}}{4x^2} \right)^{(k+1)/2}, & \text{if } k \text{ is odd.} \end{cases}$$

We note that in like fashion, the matrix N^{-1} , which begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ -1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & -3 & 1 & 0 & 0 & 0 & \dots \\ 0 & 1 & -4 & 1 & 0 & 0 & \dots \\ 0 & 0 & 7 & -6 & 1 & 0 & \dots \\ 0 & 0 & -1 & 11 & -7 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

is a ‘‘mixture’’ (in shifted alternate row fashion) of the two matrices

$$\left(\frac{1}{1+3x+2x^2}, \frac{x}{1+3x+2x^2} \right) \quad \text{and} \quad \left(\frac{1}{1+x}, \frac{x}{1+3x+2x^2} \right).$$

For instance, the array $\left(\frac{1}{1+3x+2x^2}, \frac{x}{1+3x+2x^2} \right)$ begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ -3 & 1 & 0 & 0 & 0 & 0 & \dots \\ 7 & -6 & 1 & 0 & 0 & 0 & \dots \\ -15 & 23 & -9 & 1 & 0 & 0 & \dots \\ 31 & -72 & 48 & -12 & 1 & 0 & \dots \\ -63 & 201 & -198 & 82 & -15 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix},$$

while the array $\left(\frac{1}{1+x}, \frac{x}{1+3x+2x^2} \right)$ begins

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ -1 & 1 & 0 & 0 & 0 & 0 & \dots \\ 1 & -4 & 1 & 0 & 0 & 0 & \dots \\ -1 & 11 & -7 & 1 & 0 & 0 & \dots \\ 1 & -26 & 30 & -10 & 1 & 0 & \dots \\ -1 & 57 & -102 & 58 & -13 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

We have

$$N^{-1} \cdot M = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 1 & 0 & 0 & 0 & \dots \\ 0 & 2 & 3 & 1 & 0 & 0 & \dots \\ 0 & 2 & 5 & 4 & 1 & 0 & \dots \\ 0 & 4 & 12 & 13 & 6 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

We find that the production matrix of the inverse of this matrix is given by

$$\begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & -1 & 1 & 0 & 0 & 0 & \dots \\ 0 & 0 & -2 & 1 & 0 & 0 & \dots \\ 0 & 0 & 0 & -1 & 1 & 0 & \dots \\ 0 & 0 & 0 & 0 & -2 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & -1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

This is the beheading of the inverse of the matrix

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 1 & 0 & 0 & 0 & \dots \\ 0 & 2 & 2 & 1 & 0 & 0 & \dots \\ 0 & 2 & 2 & 1 & 1 & 0 & \dots \\ 0 & 4 & 4 & 2 & 2 & 1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

The form of the production matrix of the inverse is reflected in the structure of $N^{-1} \cdot M$ as follows: the internal elements of each even row satisfy

$$t_{i,j} = 1 \cdot t_{1-1,j-1} + 1 \cdot t_{i-1,j},$$

while for odd rows we have

$$t_{i,j} = 1 \cdot t_{i-1,j-1} + 2 \cdot t_{i-1,j}.$$

We remark that it is clear that the interleaved structure of N , based on two Riordan arrays, will be replicated in the case of any sequence a_n of the form $1, 1, r, 1, r, 1, r, 1, \dots$

2 Conclusion

Using the parameters of equivalent Stieltjes and Jacobi continued fractions, we have defined two matrices N and M , and we have studied the product $N^{-1}M$ in three specific cases. In each case, some noteworthy results have emerged. We conclude that the matrix $N^{-1}M$ is worthy of further study.

References

- [1] P. Barry and A. Hennessy, Meixner-type results for Riordan arrays and associated integer sequences, *J. Integer Seq.*, **13** (2010), [Article 10.9.4](#)
- [2] P. Barry, Riordan arrays, orthogonal polynomials as moments, and Hankel transforms, *J. Integer Seq.*, **14** (2011), [Article 11.2.2](#)
- [3] G-S. Cheon, J-H. Jung, and Y. Lim, A q -analogue of the Riordan group, *Linear Algebra Appl.*, Available online 5 November 2013.
- [4] T. S. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, New York, 1978.
- [5] D. E. Davenport, L. W. Shapiro, and L. C. Woodson, The double Riordan group, *Electron. J. Combin.*, **18** (2012), #P33.
- [6] E. Deutsch, L. Ferrari, and S. Rinaldi, Production matrices, *Adv. in Appl. Math.*, **34** (2005), 101–122.
- [7] E. Deutsch, L. Ferrari, and S. Rinaldi, Production matrices and Riordan arrays, *Ann. Comb.*, **13** (2009), 65–85.
- [8] W. Gautschi, *Orthogonal Polynomials: Computation and Approximation*, Clarendon Press, Oxford.
- [9] I. Graham, D. E. Knuth, and O. Patashnik, *Concrete Mathematics*, Addison–Wesley, Reading, MA, 1994.
- [10] C. Krattenthaler, Advanced determinant calculus, *Séminaire Lotharingien Combin.* **42** (1999), Article B42q, available electronically at <http://www.emis.de/journals/SLC/wpapers/s42kratt.html>.
- [11] C. Krattenthaler, Advanced determinant calculus: A complement, *Linear Algebra Appl.*, **411** (2005), 68–166.
- [12] J. W. Layman, The Hankel transform and some of its properties, *J. Integer Seq.*, **4** (2001), [Article 01.1.5](#).

- [13] P. Peart and W.-J. Woan, Generating functions via Hankel and Stieltjes matrices, *J. Integer Seq.*, **3**(2000), [Article 00.2.1](#).
- [14] L. W. Shapiro, S. Getu, W.-J. Woan, and L.C. Woodson, The Riordan Group, *Discr. Appl. Math.* **34** (1991), 229–239.
- [15] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*. Published electronically at <http://oeis.org>, 2011.
- [16] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences, *Notices Amer. Math. Soc.*, **50** (2003), 912–915.
- [17] G. Szegö, *Orthogonal Polynomials*, 4th ed. Providence, RI, Amer. Math. Soc., 1975.
- [18] H. S. Wall, *Analytic Theory of Continued Fractions*, AMS Chelsea Publishing, 1967.

2010 *Mathematics Subject Classification*: Primary 15B36; Secondary 11B83, 15A09, 30B70, 42C05.

Keywords: Matrix, Stieltjes continued fraction, Jacobi continued fraction, orthogonal polynomials, production matrix, Riordan array, Hankel transform

Concerns sequences [A000108](#), [A001003](#), [A033184](#), [A036442](#), [A039599](#), [A172094](#).