

The Fibonacci Quart. 1987 (vol.25,1): 72-75

A COMPLETE CHARACTERIZATION OF B-POWER FRACTIONS THAT CAN BE REPRESENTED AS SERIES OF GENERAL n-BONACCI NUMBERS

JIN-ZAI LEE and JIA-SHENG LEE
Soochow University, Taipei, Taiwan, R.O.C.
 (Submitted April 1985)

1. INTRODUCTION AND MAIN RESULT

In 1953, Fenton Stancliff [5] noted that

$$\frac{1}{89} = .0112358 \quad = \sum_{k=0}^{\infty} 10^{-(k+1)} F_k,$$

13
21
...

where F_k denotes the k^{th} Fibonacci number. Until recently, this expansion was regarded as an anomalous numerical curiosity, possibly related to the fact that 89 is a Fibonacci number (see Remark in [5]), but not generalizing to other fractions in an obvious manner.

In 1980, C. F. Winans [6] showed that the sums $\sum 10^{-(k+1)} F_{\alpha k}$ approximate $1/71$, $2/59$, and $3/31$ for $\alpha = 2, 3$, and 4 , respectively. Moreover, he showed that the sums $\sum 10^{-2(k+1)} F_{\alpha k}$ approximate $1/9899$, $1/9701$, $2/9599$, and $3/9301$ for $\alpha = 1, 2, 3$, and 4 , respectively.

Since then, several authors proved general theorems on fractions that can be represented as series involving Fibonacci numbers and general n -Bonacci numbers [1, 2, 3, 4]. In the present paper we will prove a theorem which includes as special cases all the earlier results. We introduce some notation in order to state our theorem.

Let arbitrary complex numbers $A_0, A_1, \dots, A_m, W_0, W_1, \dots, W_m$, and B be given. Construct the sequence W_k by the recursion

$$W_{n+m+1} = \sum_{r=0}^m A_r W_{n+m-r}$$

for $n \geq 0$ or, equivalently, by the formula

$$W_n = \sum_{r=0}^m \lambda_r \omega_r^n$$

for any integer n where ω_r ($r = 0, 1, \dots, m$) are the zeros of the polynomial

$$q(z) = z^{m+1} - \sum_{r=0}^m A_r z^{m-r}$$

and $(\lambda_0, \lambda_1, \dots, \lambda_m)$ is the unique solution of the system of $m+1$ linear equations

$$\sum_{r=0}^m \lambda_r \omega_r^n = W_n \quad (n = 0, 1, \dots, m)$$

B-POWER FRACTIONS AS SERIES OF GENERAL n -BONACCI NUMBERS

(see [2], p. 35). Finally, we introduce, for any integer α ,

$$M(m) = \prod_{r=0}^m (B - \omega_r^\alpha).$$

Theorem: For integers $\alpha \geq 1$, $\beta \geq 0$, and any complex B satisfying

$$\max_{0 \leq r \leq m} |\omega_r^\alpha / B| < 1,$$

we have the formula

$$M(m) \cdot \sum_{k=0}^{\infty} B^{-k} W_{\alpha k + \beta} = B \cdot \sum_{r=0}^m \lambda_r \omega_r^\beta \cdot \prod_{\substack{0 \leq k \leq m \\ k \neq r}} (B - \omega_k^\alpha).$$

Remark: In the above formula, $M(m)$ and the right-hand side are in fact integers if $B, A_0, A_1, \dots, A_m, W_0, W_1, \dots, W_m$ are all integers.

Now we can comment on earlier results in more detail. In 1981, Hudson and Winans [1] handled the case of the ordinary Fibonacci sequence with $\beta = 0$, $B = 10^n$. According to [3] and [4], their result can be written as

$$\sum_{k=1}^{\infty} 10^{-n(k+1)} F_{\alpha k} = \frac{F_\alpha}{10^{2n} - 10^n L_\alpha - (-1)^\alpha},$$

where L_α denote the Lucas numbers. Also in 1981, Long [4] treated the case of the general Fibonacci sequence, i.e., $m = 1$ and arbitrary A_0, A_1, W_0, W_1 , and B , with the restriction, however, to $\alpha = 1$, $\beta = 0$. In 1985, Köhler [2] gave the generalization for arbitrary $m, A_0, A_1, \dots, A_m, W_0, W_1, \dots, W_m, B$, again with the restriction to $\alpha = 1$, $\beta = 0$. His result is

$$\sum_{k=1}^{\infty} B^{-k} W_{k-1} = p(B)/q(B),$$

where p is a polynomial of degree m with explicitly given coefficients. Also in 1985, Lee [3] discussed the cases $m = 1$ and $m = 2$ of general Fibonacci and Tribonacci sequences with arbitrary α and β . The results of [3] will be deduced from our Theorem in Examples 1 and 2 below. For this purpose, we introduce the notation

$$S_n = \sum_{r=0}^m \omega_r^n, \quad L(m) = M(m) \cdot \sum_{k=0}^{\infty} B^{-k} W_{\alpha k + \beta}.$$

Proof of Theorem: We have

$$\begin{aligned} \sum_{k=0}^{\infty} B^{-k} W_{\alpha k + \beta} &= \sum_{k=0}^{\infty} B^{-k} \sum_{r=0}^m \lambda_r \omega_r^{\alpha k + \beta} \\ &= \sum_{r=0}^m \lambda_r \omega_r^\beta \left(\sum_{k=0}^{\infty} (B^{-1} \omega_r^\alpha)^k \right) = B \sum_{r=0}^m \lambda_r \omega_r^\beta \cdot \frac{1}{B - \omega_r^\alpha}. \end{aligned}$$

Convergence is guaranteed by the condition on B . In the same way, we obtain

B-POWER FRACTIONS AS SERIES OF GENERAL n -BONACCI NUMBERS

$$\sum_{k=1}^{\infty} B^{-k} W_{\alpha k + \beta} = \sum_{r=0}^m \lambda_r \omega_r^\beta \cdot \frac{\omega_r^\alpha}{B - \omega_r^\alpha}.$$

Multiplying with $M(m)$ yields the Theorem.

Remark: Partial sums of the series in our Theorem can be expressed by these series, according to the formula

$$B^n \cdot \sum_{k=0}^{\infty} B^{-k} W_{\alpha k + \beta} = B^n \cdot \sum_{k=0}^n B^{-k} W_{\alpha k + \beta} + \sum_{k=1}^{\infty} B^{-k} W_{\alpha k + (\alpha n + \beta)}.$$

2. EXAMPLES

Example 1: The general Fibonacci sequence. Take $m = 1$. Then we have

$$W_{n+2} = A_0 W_{n+1} + A_1 W_n,$$

$$M(1) = (B - \omega_0^\alpha)(B - \omega_1^\alpha) = B^2 - BS_\alpha + (-A_1)^\alpha,$$

$$\begin{aligned} \sum_{k=1}^{\infty} B^{-k} W_{\alpha k + \beta} &= \{\lambda_0 \omega_0^{\alpha + \beta} (B - \omega_1^\alpha) + \lambda_1 \omega_1^{\alpha + \beta} (B - \omega_0^\alpha)\} / M(1) \\ &= (BW_{\alpha + \beta} - (-A_1)^\alpha W_\beta) / M(1), \end{aligned}$$

$$\sum_{k=0}^{\infty} B^{-k} W_{\alpha k + \beta} = B(BW_\beta - (-A_1)^\alpha W_{\beta - \alpha}) / M(1).$$

As to the partial sums, we get

$$\begin{aligned} \sum_{k=0}^n B^{n-k} W_{\alpha k + \beta} &= B^n L(1) / M(1) - \sum_{k=1}^{\infty} B^{-k} W_{\alpha k + \alpha n + \beta} \\ &= \frac{B^n L(1) - BW_{\alpha(n+1) + \beta} + (-A_1)^\alpha W_{\alpha n + \beta}}{B^2 - BS_\alpha + (-A_1)^\alpha}. \end{aligned}$$

These formulas are equal to Theorems 1-3 of [3].

Example 2: The general Tribonacci sequence. Take $m = 2$. Then we obtain

$$W_{n+3} = A_0 W_{n+2} + A_1 W_{n+1} + A_2 W_n,$$

$$M(2) = B^3 - B^2 S_\alpha + BA_2^\alpha S_{-\alpha} - A_2^\alpha,$$

$$\sum_{k=1}^{\infty} B^{-k} W_{\alpha k + \beta} = (B^2 W_{\alpha + \beta} + B(W_{2\alpha + \beta} - S_\alpha W_{\alpha + \beta}) + A_2^\alpha W_\beta) / M(2),$$

$$\sum_{k=0}^{\infty} B^{-k} W_{\alpha k + \beta} = B(B^2 W_\beta + B(W_{\alpha + \beta} - S_\alpha W_\beta) + A_2^\alpha W_{\beta - \alpha}) / M(2),$$

$$\sum_{k=0}^n B^{n-k} W_{\alpha k + \beta} = B^n L(2) / M(2) - \sum_{k=1}^{\infty} B^{-k} W_{\alpha k + \alpha n + \beta}$$

(continued)

B-POWER FRACTIONS AS SERIES OF GENERAL n -BONACCI NUMBERS

$$= \frac{B^n L(2) - B^2 W_{\alpha(n+1)+\beta} + B(S_\alpha W_{\alpha(n+1)+\beta} - W_{\alpha(n+2)+\beta}) - A_2^\alpha W_{\alpha n+\beta}}{B^3 - B^2 S_\alpha + B A_2^\alpha S_{-\alpha} - A_2^\alpha}$$

These formulas are equal to (9) and Theorems 7 and 8 in [3], and a misprint in Theorem 7 in [3] is corrected.

Example 3: The general Tetranacci sequence. Take $m = 3$. Then we have

$$W_{n+4} = A_0 W_{n+3} + A_1 W_{n+2} + A_2 W_{n+1} + A_3 W_n,$$

$$M(3) = B^4 - B^3 S_\alpha + B^2 (S_\alpha^2 - S_{2\alpha})/2 - B(-A_3)^\alpha S_{-\alpha} + (-A_3)^\alpha,$$

$$\begin{aligned} \sum_{k=1}^{\infty} B^{-k} W_{\alpha k+\beta} &= \{B^3 W_{\alpha+\beta} + B^2 (W_{2\alpha+\beta} - S_\alpha W_{\alpha+\beta}) \\ &\quad + B(-A_3)^\alpha (S_{-\alpha} W_\beta - W_{\beta-\alpha}) - (-A_3)^\alpha W_\beta\} / M(3), \end{aligned}$$

$$\begin{aligned} \sum_{k=0}^{\infty} B^{-k} W_{\alpha k+\beta} &= B\{B^3 W_\beta + B^2 (W_{\alpha+\beta} - S_\alpha W_\beta) + B(2W_{2\alpha+\beta} - 2S_\alpha W_{\alpha+\beta} \\ &\quad + (S_\alpha^2 - S_{2\alpha}) W_\beta) / 2 - (-A_3)^\alpha W_{\beta-\alpha}\} / M(3), \end{aligned}$$

$$\begin{aligned} \sum_{k=0}^n B^{-k} W_{\alpha k+\beta} &= \{B^n L(3) - B^3 W_{\alpha(n+1)+\beta} - B^2 (W_{\alpha(n+2)+\beta} - S_\alpha W_{\alpha(n+1)+\beta}) \\ &\quad - B(-A_3)^\alpha (S_{-\alpha} W_{\alpha n+\beta} - W_{\alpha(n+1)+\beta}) + (-A_3)^\alpha W_{\alpha n+\beta}\} / M(3). \end{aligned}$$

Formulas for $m \geq 4$ can be obtained in a similar manner.

ACKNOWLEDGMENTS

We are deeply thankful to Dr. Hong-Jinh Chang and to the referee for the earnest discussions and valuable suggestions made.

REFERENCES

1. R. H. Hudson & C. F. Winans. "A Complete Characterization of the Decimal Fractions That Can Be Represented as $\sum 10^{-k(i+1)} F_{\alpha i}$, Where $F_{\alpha i}$ Is the α th Fibonacci Number." *The Fibonacci Quarterly* 19, no. 5 (1981):414-421.
2. G. Köhler. "Generating Functions of Fibonacci-Like Sequences and Decimal Expansions of Some Fractions." *The Fibonacci Quarterly* 23, no. 1 (1985): 29-35.
3. J. S. Lee. "A Complete Characterization of B-Power Fractions That Can Be Represented as Series of General Fibonacci Numbers or of General Tribonacci Numbers." *Tamkang J. of Management Sciences* 6, no. 1 (1985):41-52.
4. C. T. Long. "The Decimal Expansion of 1/89 and Related Results." *The Fibonacci Quarterly* 19, no. 1 (1981):53-55.
5. F. Stancliff. "A Curious Property of a_{ii} ." *Scripta Mathematica* 19 (1953): 126.
6. C. F. Winans. "The Fibonacci Series in the Decimal Equivalents of Fractions." *A Collection of Manuscripts Related to the Fibonacci Sequence*. Santa Clara, Calif.: The Fibonacci Association, 1980, pp. 78-81.

◆◆◆◆