

SOME OPERATIONAL FORMULAS FOR THE  $q$ -LAGUERRE POLYNOMIALS

NADHLA A. AL-SALAM

University of Alberta, Edmonton, Alberta, Canada (T6C, 2C1)

(Submitted August 1982)

1. INTRODUCTION

L. B. Rédei [7] proved an operational identity for the Laguerre polynomials that was later generalized by Viskov [9]. Viskov's main results were as follows: if  $D = d/dx$ , then for  $n = 0, 1, 2, \dots$ , we have

$$L_n^{(\alpha)}(x) = \frac{(-1)^n}{n!} e^{x\{(\alpha + 1 + xD)D\}^n} e^{-x} = \frac{(-1)^n}{n!} e^{xB^n} e^{-x} \quad (1.1)$$

and

$$L_n^{(\alpha)}(x) = \frac{1}{n!} \{(1 + \alpha - x + xD)(1 - D)\}^n \cdot 1, \quad (1.2)$$

where  $L_n^{(\alpha)}(x)$  is the  $n^{\text{th}}$  Laguerre polynomial.

A third formula of a similar nature was given earlier by Carlitz [2]:

$$L_n^{(\alpha)}(x) = \frac{1}{n!} \prod_{k=1}^n (xD - x + \alpha + k) \cdot 1. \quad (1.3)$$

Recently, there has been renewed interest in  $q$ -identities and operators as well as in the  $q$ -Laguerre polynomial (see, e.g., [3], [5], [6]). Therefore, we felt it would also be interesting to discuss  $q$ -generalizations of the identities (1.1)-(1.3). In the following, we shall assume always that  $|q| < 1$ .

We first introduce the following notation:

$$[a]_0 = (a; q)_0 = 1;$$

$$[a]_n = (a; q)_n = (1 - a)(1 - aq) \dots (1 - aq^{n-1}) \quad (n = 1, 2, 3, \dots).$$

Also, we shall use  $[a]_\infty = (a; q)_\infty$  to mean the convergent product

$$\prod_{k=0}^{\infty} (1 - aq^k).$$

It is well known that  $[a]_\infty$  is a  $q$ -analog of the exponential function. Thus, we have

$$\lim_{q \rightarrow 1^-} (-1 - q)x; q)_\infty^{-1} = e^{-x}.$$

For this reason, the more suggestive notation

$$(x; q)_\infty^{-1} = e_q(x)$$

is used for a  $q$ -analog of the exponential function.

The  $q$ -derivatives of a function  $f(x)$  is given by

$$D_q f(x) = \frac{f(x) - f(xq)}{x},$$

so whenever  $f$  has a derivative at  $x$ , we have

$$\lim_{q \rightarrow 1} \frac{1}{(1 - q)} D_q f(x) = f'(x).$$

SOME OPERATIONAL FORMULAS FOR THE  $q$ -LAGUERRE POLYNOMIALS

We shall also use the substitution operator  $\eta: \eta f(x) = f(qx)$ . It is related to the  $q$ -derivative by means of  $x D_q = I - \eta$ , where  $I$  is the identity operator. Note that  $x$  and  $D_q$  do not commute.

We recall that the  $q$ -Laguerre polynomials [6] are defined by

$$L_n^{(\alpha)}(x|q) = \frac{[q^{\alpha+1}]_n}{[q]_n} \sum_{k=0}^n \frac{[q^{-n}]_k q^{\frac{1}{2}k(k+1)+k(n+\alpha)}}{[q]_k [q^{\alpha+1}]_k} x^k \tag{1.4}$$

so that

$$\lim_{q \rightarrow 1} L_n^{(\alpha)}((1-q)x|q) = L_n^{(\alpha)}(x), \quad n = 0, 1, 2, \dots$$

These polynomials, which are orthogonal and belong to an indetermined Stieltjes moment problem ([3] and [6]), were known to W. Hahn [5]. They have, among other properties, a Rodrigues formula:

$$L_n^{(\alpha)}(x|q) = \frac{x^{-\alpha} [-x]_{\infty}}{[q]_n} D_q^n \left\{ \frac{x^{\alpha+n}}{[-x]_{\infty}} \right\}. \tag{1.5}$$

Cigler [4] gave the representation

$$L_n^{(\alpha)}(x|q) = \frac{(-1)^n}{[q]_n} (\eta - D_q)^{n+\alpha} x^n = (-1)^n x^{-\alpha} \frac{1}{[q]_n} (\eta - D_q)^n x^{n+\alpha}. \tag{1.6}$$

Representations (1.5) and (1.6) are both of the same nature—the  $n^{\text{th}}$  iterate of the operator ( $D_q$  or  $\eta - D_q$ , respectively) acts on a function that depends on  $n$  also. In some applications, this is a drawback. This is why (1.1) and (1.2) are interesting.

2. A  $q$ -ANALOG OF THE REDEI-VISKOV OPERATOR

Put

$$B_q = \{(1 - q^{\alpha+1})I + q^{\alpha+1}x D_q\} D_q. \tag{2.1}$$

Thus, formally, we have

$$\lim_{q \rightarrow 1} \frac{B_q}{(1-q)^2} f(x) = (\alpha + 1 + xD) D f(x) = B f(x),$$

which is the operator that appears in the right-hand side of (1.1).

It is easy to see that

$$B_q x^n = (1 - q^n)(1 - q^{n+\alpha}) x^{n-1}, \tag{2.2}$$

from which we can verify another representation for the  $B_q$  operator, namely,

$$B_q = (I - q^{\alpha+1}\eta) D_q \tag{2.3}$$

and

$$B_q = x^{-\alpha} D_q x^{\alpha+1} D_q = D_q x^{1-\alpha} D_q x^{\alpha}. \tag{2.4}$$

The latter representation shows that the operator  $B_q$  is also a  $q$ -analog of the Bessel operator (see [10]):

$$B = x^{-\alpha} \frac{d}{dx} x^{\alpha+1} \frac{d}{dx}.$$

From the relation  $D_q(1 - q^{\alpha+1}\eta) = (1 - q^{\alpha+2}\eta) D_q$ , we get, by induction,

SOME OPERATIONAL FORMULAS FOR THE  $q$ -LAGUERRE POLYNOMIALS

$$B_q^n = \left\{ \prod_{k=1}^n (1 - q^{\alpha+k}\eta) \right\} D_q^n \quad (n = 0, 1, 2, \dots). \quad (2.5)$$

It is easy to see that  $D_q\{e_q(-x)f(x)\} = e_q(-x)(D_q - \eta)f(x)$ . Thus, we have, for any formal power series  $F(x)$ ,

$$F(D_q)\{e_q(-x)f(x)\} = e_q(-x)F(D_q - \eta)f(x). \quad (2.6)$$

Using mathematical induction and noting that

$$(D_q - \eta)(I - q^{\mu}(1+x)\eta) = (I - q^{\mu+1}(1+x)\eta)(D_q - \eta),$$

we get

$$\begin{aligned} B_q^n\{e_q(-x)f(x)\} &= e_q(-x)\{(I - q^{\alpha+1}(1+x)\eta)(D_q - \eta)\}^n \cdot f(x) \\ &= e_q(-x) \left\{ \prod_{k=1}^n (1 - q^{\alpha+k}(1+x)\eta) \right\} (D_q - \eta)^n \cdot f(x) \\ &= \frac{1}{x^n} e_q(-x) \prod_{k=1}^n (1 - q^{\alpha+k-n}(1+x)\eta)(1 - q^{k-n}(1+x)\eta) \cdot f(x). \end{aligned}$$

Now, to obtain operational representations for the  $q$ -Laguerre polynomials, we first calculate

$$\begin{aligned} B_q^n e_q(-x) &= B_q^n \sum_{k=0}^{\infty} \frac{(-1)^k}{[q]_k} x^k \\ &= \sum_{k=n}^{\infty} \frac{(1 - q^k)(1 - q^{k-1}) \dots (1 - q^{k-n+1})(1 - q^{k+\alpha}) \dots (1 - q^{k-n+\alpha+1})(-x)^{k-n}}{[q]_k} \\ &= (-1)^n [q^{\alpha+1}]_n \sum_{k=0}^{\infty} \frac{[q^{\alpha+n+1}]_k}{[q]_k [q^{\alpha+1}]_k} (-x)^k. \end{aligned}$$

Andrews [1] gave a  $q$ -analog of Kummer's Theorem, i.e.,

$$\sum_{k=0}^{\infty} \frac{[\beta]_k [\alpha]_k (-1)^k q^{\frac{1}{2}k(k-1)}}{[q]_k [\gamma]_k [x\alpha]_k} \left(\frac{x\gamma}{\beta}\right)^k = \frac{[x]_{\infty}}{[x\alpha]_{\infty}} \sum_{k=0}^{\infty} \frac{[\gamma/\beta]_k [\alpha]_k}{[q]_k [\gamma]_k} x^k \quad (2.8)$$

Putting  $\alpha = 0$  in this formula, replacing  $x$  by  $-x$ , and then taking  $\gamma = q^{\alpha+1}$ ,  $\beta = q^{-n}$ , we get that

$$B_q^n e_q(-x) = \frac{(-1)^n [q^{\alpha+1}]_n}{[-x]_{\infty}} \sum_{k=0}^n \frac{[q^{-n}]_k q^{\frac{1}{2}k(k-1) + k(\alpha+n+1)}}{[q]_k} x^k = \frac{(-1)^n [q]_n}{[-x]_{\infty}} L_n^{(\alpha)}(x|q).$$

Together with (2.7), this formula gives the following three representations:

$$L_n^{(\alpha)}(x|q) = \frac{(-1)^n}{[q]_n} \{e_q(-x)\}^{-1} B_q^n \{e_q(-x)\}; \quad (2.9)$$

$$= \frac{1}{[q]_n} \prod_{k=1}^n (I - q^{\alpha+k}(1+x)\eta) \cdot 1; \quad (2.10)$$

$$= \frac{(-1)^n}{[q]_n} \{(I - q^{\alpha+1}(1+x)\eta)(D_q - \eta)\}^n \cdot 1. \quad (2.11)$$

SOME OPERATIONAL FORMULAS FOR THE  $q$ -LAGUERRE POLYNOMIALS

If we let  $q \rightarrow 1$ , then (2.9), after suitable normalization by  $(1 - q)$ , reduces to (1.1); (2.11) reduces to (1.2); and (2.10) reduces to Carlitz's formula (1.3).

Using (2.2) and (1.4), we get, for  $m = 0, 1, 2, \dots$ ,

$$E_q^m L_n^{(\alpha)}(x|q) = (-1)^m \frac{[q^{\alpha+1}]_n}{[q^{\alpha+1}]_{n-m}} q^{m(m+\alpha)} L_{n-m}^{(\alpha)}(q^{2m}x|q). \quad (2.12)$$

Notice that the operation on  $L_n^{(\alpha)}(x|q)$  by  $B_q$  reduces the degree by one without changing the value of the parameter  $\alpha$ .

There is another  $q$ -analog of the exponential function  $e^{-x}$ , namely,

$$E_q(x) = \sum_{k=0}^{\infty} \frac{(-1)^k q^{\frac{1}{2}k(k-1)}}{[q]_k} x^k = \prod_{j=0}^{\infty} (1 - xq^j).$$

If we repeat the above calculation, we can show that

$$B_q^n E_q(x) = (-1)^n [q^{\alpha+1}]_n q^{\frac{1}{2}n(n-1)} \sum_{j=0}^n \frac{(-1)^j q^{\frac{1}{2}j(j-1) + n j} [q^{\alpha+n+1}]_j}{[q]_j [q^{\alpha+1}]_j} x^j.$$

Once again we can transform the right-hand side of this formula by using (2.8) (with  $\alpha = 0, \beta = q^{\alpha+n+1}, \gamma = q^{\alpha+1}$ , and  $x \rightarrow xq^{2n}$ ), to obtain

$$B_q^n \{E_q(x)\} = (-1)^n q^{\frac{1}{2}n(n-1) + \alpha n} [q]_n E_q(xq^{2n}) L_n^{(\alpha)}(-xq^{2n-1}|q^{-1}). \quad (2.13)$$

Comparison formulas to this are:

$$e_q(-\eta^{-2} B_q)\{x^n\} = (-1)^n q^{-n(n-1)} [q]_n L_n^{(\alpha)}(q^{\alpha+1}x|q) \quad (2.14)$$

and

$$E_q(-B_q)\{x^n\} = (-1)^n [q]_n q^{\frac{1}{2}n(n-1)} L_n^{(\alpha)}(xq^{2n-1}|q^{-1}). \quad (2.15)$$

Both formulas (2.14) and (2.15) reduce in the case  $q \rightarrow 1$  to the new formula for the ordinary Laguerre polynomials:

$$e^{-B} x^n = (-1)^n n! L_n^{(\alpha)}(x) \quad (n = 0, 1, 2, \dots). \quad (2.16)$$

On the other hand, formula (2.13) reduces to (1.1).

If we calculate the right-hand side of (2.9) directly, we get

$$\begin{aligned} B_q^n e_q(-x) &= (-1)^n \prod_{k=1}^n (1 - q^{\alpha+1} \eta) e_q(-x) \\ &= \sum_{j=0}^n (-1)^{n+j} \begin{bmatrix} n \\ j \end{bmatrix} q^{\frac{1}{2}j(j-1) + (\alpha+1)j} \eta^j e_q(-x) \\ &= e_q(-x) \sum_{j=0}^n (-1)^{n+j} \begin{bmatrix} n \\ j \end{bmatrix} q^{\frac{1}{2}j(j+1) + \alpha j} [-x]_j. \end{aligned} \quad (2.17)$$

The second equality is due to the Euler identity

$$\prod_{k=1}^n (1 - q^{k-1}x) = \sum_{j=0}^n (-1)^j \begin{bmatrix} n \\ j \end{bmatrix} q^{\frac{1}{2}j(j-1)} x^j,$$

SOME OPERATIONAL FORMULAS FOR THE  $q$ -LAGUERRE POLYNOMIALS

where  $\begin{bmatrix} n \\ j \end{bmatrix}$  stands for the  $q$ -binomial coefficient, i.e., for 1 if  $j = 0$  and for  $(1 - q)(1 - q^{n-1}) \dots (1 - q^{n-j+1}) / (1 - q)(1 - q^2) \dots (1 - q^j)$

if  $j \geq 1$ . Combining (2.9) and (2.17), we obtain another expansion for the  $q$ -Laguerre polynomial:

$$L_n^{(\alpha)}(x|q) = \frac{1}{[q]_n} \sum_{j=0}^n (-1)^j \begin{bmatrix} n \\ j \end{bmatrix} [-x]_j q^{\frac{1}{2}j(j+1) + \alpha j}. \quad (2.18)$$

REFERENCES

1. G. Andrews. "On the  $q$ -Analog of Kummer's Theorem and Applications." *Duke Math. J.* 40 (1972):525-28.
2. L. Carlitz. "A Note on the Laguerre Polynomials." *Michigan Math. J.* 7 (1960):219-23.
3. T. S. Chiharra. "Orthogonal Polynomials with Brenke Type Generating Functions." *Duke Math. J.* 35 (1968):505-18.
4. J. Cigler. "Operator methoden für  $q$ -Identitäten II:  $q$ -Laguerre Polynome." *Monatsh. Math.* 91 (1981):105-17.
5. W. Hahn. "Über orthogonalpolynome, die  $q$ -differenzen-gleichungen genügen." *Math. Math.* 2 (1949):4-34.
6. D. Moak. "The  $q$ -Analogue of the Laguerre Polynomials." *J. Math. Anal. Appl.* 81 (1981):20-47.
7. L. B. Rédei. "An Identity for the Laguerre Polynomials." *Acta Sci. Math.* 37 (1975):115-16.
8. G. Szégo: *Orthogonal Polynomials*. New York: American Mathematical Society, Rev. ed., 1959.
9. O. V. Viskov. "L. B. Redei Identity for the Laguerre Polynomials." *Acta Sci. Math.* 39 (1977):27-28. (Translated from the Russian.)
10. N. Meller. "On an Operational Calculus for the Operator  $B_\alpha$ ." *Vych. Math. Izd-vo Akad. Nauk SSSR* #6 (1960):161-68. (Translated from the Russian.)

◆◆◆◆