

# The Fibonacci Quarterly 1985 (23,4): 290-293

## ON $m$ -TH ORDER LINEAR RECURRENCES

CLAUDE LEVESQUE

Université Laval, Québec, P. Québec, Canada, G1K 7P4

(Submitted April 1983)

### 1. INTRODUCTION

Fix numbers  $u_0, u_1, \dots, u_{-1}$ , and for every  $n \geq 0$ , define  $u_{m+n}$  by means of the  $m$  preceding terms with the rule

$$u_{m+n} - k_1 u_{m+n-1} - \dots - k_m u_n = 0, \text{ with } k_m \neq 0. \quad (1.1)$$

In this note, we wish to present two formulas for these numbers  $u_n$  satisfying the above  $m$ -th order linear recurrence (Sections 2 and 3).

These results are probably known to some readers; however, since from time to time we happen to see in the literature special cases of these formulas, it may be worthwhile to present them once and for all.

Note that for  $m = 2$ ,  $k_1 = k_2 = 1$ ,  $u_0 = u_1 = 1$ , one is dealing with the Fibonacci numbers, which have been extensively studied by many authors (see, for instance, [13], [5], and [3]), and which were used by Matijasevič [9] in his notorious proof that Hilbert's tenth problem is recursively unsolvable.

### 2. GENERATING FUNCTION AND BINET'S FORMULA

Using the  $m$ -th order linear recurrence

$$u_{m+n} = k_1 u_{m+n-1} + k_2 u_{m+n-2} + \dots + k_m u_n, \quad k_m \neq 0, \quad (2.1)$$

(with the  $k_i$ 's in  $\mathbb{Z}$  for instance, or in a given field), we easily obtain

$$\left( \sum_{n=0}^{\infty} u_n X^n \right) (1 - k_1 X - \dots - k_m X^m) = \sum_{i=0}^{m-1} v_i X^i,$$

where the  $v_i$ 's, functions of the initial conditions on  $u_0, u_1, \dots, u_{m-1}$ , are defined by

$$v_i = - \sum_{j=0}^i u_{i-j} k_j, \quad (2.2)$$

(with  $k_0 = -1$  throughout this article). Associated with that recursive sequence is the following polynomial,

$$f(X) = X^m - k_1 X^{m-1} - \dots - k_{m-1} X - k_m = (X - \alpha_1)(X - \alpha_2) \dots (X - \alpha_m),$$

whose roots we assume distinct (and nonzero, since  $k_m \neq 0$ ).

Then we have

$$\sum_{n=0}^{\infty} u_n X^n = \frac{v_0 + v_1 X + \dots + v_{m-1} X^{m-1}}{1 - k_1 X - k_2 X^2 - \dots - k_m X^m}$$

(continued)

ON  $m$ -TH ORDER LINEAR RECURRENCES

$$\begin{aligned}
 &= \frac{v_0 + v_1X + \dots + v_{m-1}X^{m-1}}{(1 - \alpha_1X)(1 - \alpha_2X) \cdot \dots \cdot (1 - \alpha_mX)} \\
 &= (v_0 + v_1X + \dots + v_{m-1}X^{m-1})(1 + p_1X + \dots + p_jX^j + \dots),
 \end{aligned}$$

where  $p_j$  stands for the sum of all symmetric functions of weight  $j$  in  $\alpha_1, \alpha_2, \dots, \alpha_m$ ; in other words [2],  $p_j$  is the "sum of the homogeneous products of  $j$  dimensions" of the  $m$  symbols  $\alpha_1, \dots, \alpha_m$ .

Let us recall from Volume 1 of [2, p. 178] that

$$p_j = \sum_{i=1}^m \frac{\alpha_i^{m-1+j}}{f'(\alpha_i)} \text{ with } p_0 = 1,$$

and that  $p_{-1} = p_{-2} = \dots = p_{-m+1} = 0$  (which follows from Example 4 of p. 172). We therefore obtain for the  $m$ -th number  $u_n$  what can be called

BINET'S FORMULA: 
$$u_n = \sum_{j=0}^{m-1} v_j p_{n-j}.$$

EXAMPLES: (1) Let  $v_0 = v_1 = \dots = v_{m-2} = 0, v_{m-1} = 1$ ; then, as in Formula 9 of [7],

$$u_n = p_{n-m+1} = \frac{\alpha_1^n}{f'(\alpha_1)} + \frac{\alpha_2^n}{f'(\alpha_2)} + \dots + \frac{\alpha_m^n}{f'(\alpha_m)}.$$

(2) For  $m = 2, m = 3$ , we recover Binet's formulas of [3] and [11].

(3) For  $n \in \mathbb{N} = \{0, 1, 2, \dots\}$ , define  $s_n$  by

$$s_n = \alpha_1^n + \alpha_2^n + \dots + \alpha_m^n.$$

As is well known (see [2]), Newton's formulas state that:

$$s_n = \begin{cases} m & \text{if } n = 0 \\ k_1 & \text{if } n = 1 \\ k_1 s_{n-1} + k_2 s_{n-2} + \dots + k_{n-1} s_1 + n k_n & \text{if } 2 \leq n \leq m-1 \\ k_1 s_{n-1} + k_2 s_{n-2} + \dots + k_m s_{n-m} & \text{if } n \geq m. \end{cases}$$

In particular, if  $u_n = s_n$ , then  $\{u_n\}$  satisfies (2.1).

Thus, using the fact that  $k_0 = -1$ , we find that (2.2) gives:

$$\begin{cases} v_0 = m = -mk_0 \\ v_1 = s_1 - mk_1 = -(m-1)k_1 \\ v_2 = s_2 - s_1 k_1 - mk_2 = -(m-2)k_2 \\ \vdots \\ v_{m-1} = s_{m-1} - s_{m-2} k_1 - s_{m-3} k_2 - \dots - s_1 k_{m-2} - mk_{m-1} = -1k_{m-1}. \end{cases}$$

In short, for  $j = 0, 1, \dots, m-1, v_j = -(m-j)k_j$ , and Binet's formula becomes

$$s_n = -\sum_{j=0}^{m-1} (m-j)k_j p_{n-j}.$$

3. ANOTHER FORMULA

We can also use the multinomial theorem to obtain a formula for  $u_n$  that is a function of (i.e., the elementary symmetric functions of)  $k_1, k_2, \dots, k_m$ . Here we no longer require that the roots of  $f$  be distinct. Within a certain radius of convergence, we find that

$$\begin{aligned} \sum_{n=0}^{\infty} u_n X^n &= \frac{v_0 + v_1 X + \dots + v_{m-1} X^{m-1}}{1 - (k_1 X + k_2 X^2 + \dots + k_m X^m)} \\ &= (v_0 + v_1 X + \dots + v_{m-1} X^{m-1}) \left[ \sum_{j=0}^{\infty} (k_1 X + \dots + k_{m-1} X^{m-1} + k_m X^m)^j \right] \\ &= (v_0 + v_1 X + \dots + v_{m-1} X^{m-1}) \left( \sum_{i=0}^{\infty} A(i) X^i \right), \end{aligned}$$

where

$$A(i) = \sum \frac{(t_1 + t_2 + \dots + t_m)!}{t_1! t_2! \dots t_m!} k_1^{t_1} k_2^{t_2} \dots k_m^{t_m},$$

the last sum being taken over all  $m$ -tuples  $(t_1, t_2, \dots, t_m)$  of  $\mathbb{N}^m$  such that

$$t_1 + 2t_2 + \dots + mt_m = i.$$

Defining  $A(i)$  to be 0 for  $i < 0$ , we therefore conclude:

$$\boxed{u_n = \sum_{j=0}^{m-1} v_j A(n-j)} \quad (3.1)$$

**EXAMPLES:** (1) If  $v_0 = 1, v_1 = v_2 = \dots = v_{m-1} = 0$ , then

$$u_n = A(n).$$

(2) Let  $s_n = \alpha_1^n + \alpha_2^n + \dots + \alpha_m^n$ . Replacing  $v_j$  by  $-(m-j)k_j$ , and making in the  $j$ -th summation ( $j = 1, \dots, m-1$ ) of (3.1) the change of variable  $t_j \rightarrow t_j - 1$ , we obtain after a few calculations what is called in [2] Waring's formula for  $s_n$ :

$$s_n = \sum_{t_1 + 2t_2 + \dots + mt_m = n} \frac{n(t_1 + t_2 + \dots + t_m - 1)!}{t_1! t_2! \dots t_m!} k_1^{t_1} k_2^{t_2} \dots k_m^{t_m}.$$

REFERENCES

1. A. C. Aitken. *Determinants and Matrices*. 9th ed. New York: Interscience, 1956.
2. W. S. Burnside. *The Theory of Equations*. 7th ed. 2 vols. New York: Dover Publications.
3. G. Frei. "Binary Lucas and Fibonacci Polynomials, I." *Math. Nachr.* 96 (1980):83-112.
4. S. Goldberg. *Introduction to Difference Equations*. 3rd ed. New York: John Wiley & Sons, 1966.
5. V. E. Hoggatt, Jr. *Fibonacci and Lucas Numbers*. Santa Clara, Calif: The Fibonacci Association, 1973.

ON  $m$ -TH ORDER LINEAR RECURRENCES

6. Dov Jarden. *Recurring Sequences*. 3rd ed. Israel: Riveon Lematematika, 1973. Chapter 29.
7. D. Kalman. "Generalized Fibonacci Numbers by Matrix Methods." *The Fibonacci Quarterly* 20, no. 1 (1982):73-76.
8. D. Kalman. "The Generalized Vandermonde Matrix." *Math. Magazine* 57, no. 1 (1984):15-21.
9. Y. Matijasevič. "Enumerable Sets Are Diophantine." *Soviet Math. Doklady* 11 (1970):354-358.
10. K. S. Miller. *Linear Difference Equations*. New York: W. A. Benjamin, Inc., 1968.
11. W. R. Spickerman. "Binet's formula for the Tribonacci Sequence." *The Fibonacci Quarterly* 20, no. 2 (1982):118-120.
12. M. R. Spiegel. *Finite Differences and Difference Equations*. Schaum's Outline Series. New York: McGraw-Hill, 1971.
13. N. Vorob'ev. *The Fibonacci Numbers*. Boston: Heath, 1963.

◆◆◆◆