

SOME IDENTITIES INVOLVING GENERALIZED
SECOND-ORDER INTEGER SEQUENCES

Zhizheng Zhang

Department of Mathematics, Luoyang Teachers' College, Luoyang, Henan, 471022, P.R. China

(Submitted February 1996)

1. INTRODUCTION

In the notation of Horadam [2], write

$$W_n = W_n(a, b; p, q),$$

so that

$$W_n = pW_{n-1} - qW_{n-2}, \quad W_0 = a, \quad W_1 = b, \quad n \geq 2. \quad (1.1)$$

If α and β , assumed distinct, are the roots of $\lambda^2 - p\lambda + q = 0$, we have the Binet form [2]:

$$W_n = A\alpha^n + B\beta^n, \quad (1.2)$$

where $A = \frac{b - a\beta}{\alpha - \beta}$ and $B = \frac{a\alpha - b}{\alpha - \beta}$.

The sequence $\{W_n\}$ has been studied in the recent papers of Melham and Shannon [4], [5]. The purpose of this article is to establish some new identities involving W_n by using the method of Carlitz and Ferns [1].

Throughout this paper, the symbol $\binom{n}{i, j}$ is defined by $\binom{n}{i, j} = \frac{n!}{i!j!(n-i-j)!}$.

2. THE MAIN RESULTS

Carlitz and Ferns [1] have given a large number of interesting Fibonacci and Lucas identities. By adapting their method to the sequence $\{W_n\}$, we have obtained the following results.

Theorem 2.1:

$$W_{2n+k} = \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} p^j q^{n-j} W_{j+k}. \quad (2.1)$$

Lemma: Let $u = \alpha$ or β , then

$$(i) \quad -pq + (p^2 - q)u = u^3, \quad (2.2)$$

$$(ii) \quad -q^3 + pq^2u + u^6 = (p^2 - 2q)u^4, \quad (2.3)$$

$$(iii) \quad -q^5 + pq^4u + u^{10} = (p^4 - 4p^2q + 2q^2)u^6, \quad (2.4)$$

$$(iv) \quad -q^9 + pq^8u + u^{18} = \Delta u^{10}, \quad (2.5)$$

where $\Delta = p^8 - 8p^6q + 20p^4q^2 - 16p^2q^3 + 2q^4$.

Theorem 2.2:

$$(p^2 - q)W_{k+1} - pqW_k = W_{k+3}, \quad (2.6)$$

$$-q^3W_k + pq^2W_{k+1} + W_{k+6} = (p^2 - 2q)W_{k+4}, \quad (2.7)$$

$$-q^5W_k + pq^4W_{k+1} + W_{k+10} = (p^4 - 4p^2q + 2q^2)W_{k+6}, \quad (2.8)$$

$$-q^9W_k + pq^8W_{k+1} + W_{k+18} = \Delta W_{k+10}. \quad (2.9)$$

Theorem 2.3:

$$W_{3n+k} = \sum_{i+j+s=n} \binom{n}{i, j} (-1)^{i+s} p^{2j+s} q^{i+s} W_{i+j+k}, \quad (2.10)$$

$$W_{n+k} = (-q)^{-n} \sum_{i+j+s=n} \binom{n}{i, j} (-1)^j p^{2j+s} q^s W_{3i+j+k}. \quad (2.11)$$

Theorem 2.4:

$$W_{n+k} = (pq^2)^{-n} \sum_{i+j+s=n} \binom{n}{i, j} (-1)^j q^{3s} (p^2 - 2q)^i W_{4i+6j+k}, \quad (2.12)$$

$$W_{4n+k} = (p^2 - 2q)^{-n} \sum_{i+j+s=n} \binom{n}{i, j} (-1)^s p^j q^{2j+3s} W_{6i+j+k}, \quad (2.13)$$

$$W_{6n+k} = \sum_{i+j+s=n} \binom{n}{i, j} (-1)^j p^j q^{3s+2j} (p^2 - 2q)^i W_{4i+j+k}. \quad (2.14)$$

Theorem 2.5:

$$W_{n+k} = (pq^4)^{-n} \sum_{i+j+s=n} \binom{n}{i, j} (-1)^j q^{5s} (p^4 - 4p^2q + 2q^2)^i W_{6i+10j+k}, \quad (2.15)$$

$$W_{6n+k} = (p^4 - 4p^2q + 2q^2)^{-n} \sum_{i+j+s=n} \binom{n}{i, j} (-1)^s p^j q^{4j+5s} W_{10i+j+k}, \quad (2.16)$$

$$W_{10n+k} = \sum_{i+j+s=n} \binom{n}{i, j} (-1)^j p^j q^{5s+4j} (p^4 - 4p^2q + 2q^2)^i W_{6i+j+k}. \quad (2.17)$$

Theorem 2.6:

$$W_{n+k} = (pq^8)^{-n} \sum_{i+j+s=n} \binom{n}{i, j} (-1)^j q^{9s} \Delta^i W_{10i+18j+k}, \quad (2.18)$$

$$W_{10n+k} = \Delta^{-n} \sum_{i+j+s=n} \binom{n}{i, j} (-1)^s p^j q^{8j+9s} W_{18i+j+k}, \quad (2.19)$$

$$W_{18n+k} = \sum_{i+j+s=n} \binom{n}{i, j} (-1)^j p^j q^{9s+8j} \Delta^i W_{10i+j+k}. \quad (2.20)$$

3. THE PROOFS OF THE MAIN RESULTS

Since α and β are roots of $\lambda^2 - p\lambda + q = 0$, then

$$\alpha^2 = p\alpha - q, \quad (3.1)$$

$$\beta^2 = p\beta - q. \quad (3.2)$$

Now, by the binomial theorem, we have

$$\alpha^{2n} = \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} p^j q^{n-j} \alpha^j, \quad (3.3)$$

$$\beta^{2n} = \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} p^j q^{n-j} \beta^j. \quad (3.4)$$

Theorem 2.1 follows if we multiply both sides of (3.3) and (3.4) by α^k and β^k , respectively, and use the Binet form (1.2).

The Lemma can be proved by using (3.1) and (3.2). We prove only (2.3) since the proofs of (2.2), (2.4), and (2.5) are similar.

Proof of (2.3): Using (3.1) and (3.2), we have

$$\begin{aligned} -q^3 + pq^2u + u^6 &= q^2(pu - q) + u^4(pu - q) \\ &= q^2u^2 + pu^5 - qu^4 = q^2u^2 + pu^3(pu - q) - qu^4 \\ &= (p^2 - q)u^4 + q^2u^2 - pqu^3 = (p^2 - q)u^4 - qu^2(pu - q) = (p^2 - 2q)u^4. \end{aligned}$$

This completes the proof of (2.3).

Theorem 2.2 can be proved by using the results of the Lemma and proceeding in the same manner as the proof of Theorem 2.1.

The proofs of Theorems 2.3-2.6 are similar. Therefore, we prove only Theorem 2.4.

Proof of Theorem 2.4: By using (2.3) and the multinomial theorem, we have

$$\begin{aligned} (pq^2)^n u^n &= \sum_{i+j+s=n} \binom{n}{i, j} (-1)^j q^{3s} (p^2 - 2q)^i u^{4i+6j}, \\ (p^2 - 2q)^n u^{4n} &= \sum_{i+j+s=n} \binom{n}{i, j} (-1)^s p^j q^{2j+3s} u^{6i+j}, \\ u^{6n} &= \sum_{i+j+s=n} \binom{n}{i, j} (-1)^j p^j q^{3s+2j} (p^2 - 2q)^i u^{4i+j}. \end{aligned}$$

If we multiply both sides in the preceding identities by u^k and use the Binet form (1.2), we obtain (2.12), (2.13), and (2.14), respectively. This completes the proof of Theorem 2.4.

4. SOME CONGRUENCE PROPERTIES

From (2.12), (2.15), and (2.18), by using the decomposition

$$\sum_{i+j+s=n} = \sum_{i=0}^{n} + \sum_{i \neq 0}^{n},$$

we obtain

Theorem 4.1:

$$p^n q^{2n} W_{n+k} - \sum_{j=0}^n \binom{n}{j} (-1)^j q^{3n-3j} W_{6j+k} \equiv 0 \pmod{(p^2 - 2q)}, \tag{4.1}$$

$$p^n q^{4n} W_{n+k} - \sum_{j=0}^n \binom{n}{j} (-1)^j q^{5n-5j} W_{10j+k} \equiv 0 \pmod{(p^4 - 4p^2q + 2q^2)}, \tag{4.2}$$

$$p^n q^{8n} W_{n+k} - \sum_{j=0}^n \binom{n}{j} (-1)^j q^{9n-9j} W_{18j+k} \equiv 0 \pmod{\Delta}. \tag{4.3}$$

From (2.14), (2.17), and (2.20), by also using the above decomposition and Theorem 2.1, we get the following result:

Theorem 4.2:

$$W_{6n+k} - (-1)^n q^{2n} W_{2n+k} \equiv 0 \pmod{(p^2 - 2q)}, \quad (4.4)$$

$$W_{10n+k} - (-1)^n q^{4n} W_{2n+k} \equiv 0 \pmod{(p^4 - 4p^2q + 2q^2)}, \quad (4.5)$$

$$W_{18n+k} - (-1)^n q^{8n} W_{2n+k} \equiv 0 \pmod{\Delta}. \quad (4.6)$$

5. A REMARK

Some of the results in this paper are not as "practical" as others. For example, if we put $n = 10$ and $k = 0$ in (2.13), then we seek to find W_{40} . However, on the right-hand side, we need to know $W_6, W_{12}, W_{18}, \dots, W_{60}$ (and many other terms) in order to find W_{40} . In contrast, (2.14) is more practical since, in order to find W_{60} , we need to know the value of terms whose subscripts are much less than 60.

ACKNOWLEDGMENT

The author wishes to thank the anonymous referee for his/her patience and suggestions that have significantly improved the presentation of this paper.

REFERENCES

1. L. Carlitz & H. H. Ferns. "Some Fibonacci and Lucas Identities." *The Fibonacci Quarterly* **8.1** (1970):61-73.
2. A. F. Horadam. "Basic Properties of a Certain Generalized Sequence of Numbers." *The Fibonacci Quarterly* **3.2** (1965):161-76.
3. L. C. Hsu & Jiang Maosen. "A Kind of Invertible Graphical Process for Finding Reciprocal Formulas with Applications." *Acta Scien. Nat. Univ. Jilinensis* **4** (1980):43-55.
4. R. S. Melham & A. G. Shannon. "A Generalization of the Catalan Identity and Some Congruences." *The Fibonacci Quarterly* **33.1** (1995):82-84.
5. R. S. Melham & A. G. Shannon. "Some Congruence Properties of Generalized Second-Order Integer Sequences." *The Fibonacci Quarterly* **32.5** (1994):424-28.

AMS Classification Numbers: 11B37, 11B39



NEW EDITOR

On August 31, 1998, after eighteen years of continuous service, the current editor of *The Fibonacci Quarterly* will retire. He will be replaced by:

Professor Curtis Cooper
 Department of Mathematics and Computer Science
 Central Missouri State University
 Warrensburg, MO 64093-5045
 e-mail: ccooper@cmsuvmvmb.cmsu.edu

During the interim, Professor Cooper will serve as the coeditor.