



Nikita Gogin | Mika Hirvensalo

# On the Generating Function of Discrete Chebyshev Polynomials

TURKU CENTRE *for* COMPUTER SCIENCE

TUCS Technical Report  
No 819, April 2007





# On the Generating Function of Discrete Chebyshev Polynomials

**Nikita Gogin**

The author is with the University of Turku.  
`ngiri@lsi.ru`

**Mika Hirvensalo**

TUCS–Turku Centre for Computer Science, and  
Department of Mathematics, University of Turku  
FIN-20014 Turku, Finland  
`mikhirve@utu.fi`

Supported by the Academy of Finland under grant 208797

TUCS Technical Report

No 819, April 2007

### **Abstract**

We give a closed form for the generating function of the discrete Chebyshev polynomials. The closed form consists of the MacWilliams transform of Jacobi polynomials together with a binomial multiplicative factor. It turns out that the desired closed form is a solution to a special case of Heun differential equation, and that the closed form implies combinatorial identities that appear quite challenging to prove directly.

**Keywords:** Orthogonal polynomials, Discrete Chebyshev polynomials, Krawtchouk polynomials, MacWilliams transform, Generating function, Heun equation

**TUCS Laboratory**

Discrete Mathematics for Information Technology

# 1 Introduction

The discrete Chebyshev polynomials belong to the rich family of *orthogonal polynomials* (see [8] for a general treatise on the orthogonal polynomials and [3] for a previous work of the authors). The inner product associated to the discrete Chebyshev polynomials is defined with a discrete weight function, and hence the vector space  $\mathcal{P}_N$  of polynomials having degree at most  $N$  forms a natural reference for the orthogonal polynomials discussed in this article.

The sum and the scalar product in  $\mathcal{P}_N$  are defined pointwise, and the inner product is defined as

$$\langle p, q \rangle_w = \sum_{l=0}^N w_l p(l) q(l). \quad (1)$$

The *Krawtchouk polynomials* (see [6])  $K_0^{(N)}, K_1^{(N)}, \dots, K_N^{(N)}$  (of order  $N$ ) are orthogonal with respect to weight function  $w_l = \binom{N}{l}$  and the discrete Chebyshev polynomials  $D_0^{(N)}, D_1^{(N)}, \dots, D_N^{(N)}$  of order  $N$  with respect to weight function  $w_l = 1$  for each  $l$ . In addition to orthogonality, we have  $\deg(K_k^{(N)}) = \deg(D_k^{(N)}) = k$  for each  $k \in \{0, 1, \dots, N\}$ .

As (orthogonal) polynomials with ascending degree, the discrete Chebyshev polynomials form a basis of  $\mathcal{P}_N$ , and hence any polynomial  $p$  of degree at most  $N$  can be uniquely represented as

$$p = d_0 D_0^{(N)} + d_1 D_1^{(N)} + \dots + d_N D_N^{(N)}, \quad (2)$$

where  $d_l \in \mathbb{C}$ . Coefficients  $d_l$  in (2) are called the *discrete Chebyshev coefficients* of  $p$ . Since the discrete Chebyshev polynomials are orthogonal with respect to constant weight function, they have the following property important in the approximation theory: With respect to norm  $\|p - q\|^2 = \sum_{l=0}^N (p(l) - q(l))^2$ , the best approximation of  $p$  in  $\mathcal{P}_M$  can be found by simply taking  $M + 1$  first summands of (2) (see [4], for instance).

## 2 Preliminaries

### 2.1 The Discrete Chebyshev Polynomials

There are various ways to construct polynomials orthogonal with respect to inner product (1) with weight function  $w_l = 1$  so that  $\deg(D_k^{(N)}) = k$ . We choose a construction analogous to that of *Legendre polynomials* [8]. We first define the difference operator  $\Delta$  by  $\Delta f(x) = f(x + 1) - f(x)$ , the binomial coefficient by  $\binom{x}{k} = \frac{1}{k!} x(x-1) \dots (x-k+1)$ , and finally

$$D_k^{(N)}(x) = (-1)^k \Delta^k \left( \binom{x}{k} \binom{x-N-1}{k} \right). \quad (3)$$

It is straightforward to see that polynomials  $D_k$  (here and hereafter, we omit the superscript  $N$  if there is no danger of confusion) defined above form a basis of  $\mathcal{P}_N$

orthogonal with respect to inner product (1) with weight  $w_l = 1$ . Moreover, clearly  $\deg(D_k) = k$ , since one application of  $\Delta$  decreases the degree of a polynomial by one [2].

In this article, we consider (3) as the definition of discrete Chebyshev polynomials, but it is also easy to see that the following explicit expressions hold (see [2]):

$$\begin{aligned} D_k^{(N)}(x) &= \sum_{l=0}^k (-1)^l \binom{k}{l} \binom{N-x}{k-l} \binom{x}{l} \\ &= \sum_{l=0}^k (-1)^l \binom{k+l}{k} \binom{N-l}{k-l} \binom{x}{l}. \end{aligned} \quad (4)$$

Also, it is rather easy to verify that the discrete Chebyshev polynomials satisfy the following recurrence relation:

$$k^2 D_k = (2k-1) D_1 D_{k-1} - (N+k)(N-k+2) D_{k-2}, \quad (5)$$

$D_0 = 1$ ,  $D_1 = N - 2x$  (see [2]). The recurrence (5) also extends the definition of  $D_k$  to cases  $k > N$ .

The method of using *generating functions* is among the cornerstones of various areas of mathematics, and does not need any further introduction. We merely focus on the very simple form of the generating function of Krawtchouk polynomials (see [6]):

$$(1+t)^{N-x} (1-t)^x = \sum_{k=0}^{\infty} K_k^{(N)}(x) t^k. \quad (6)$$

In fact, when studying binomial distributions, it is quite natural to *define* the Krawtchouk polynomials via (6).

On the other hand, the quest for the generating function of the discrete Chebyshev polynomials seems to be a more complicated task. In what follows, we give a closed form for the generating function

$$\sum_{k=0}^{\infty} D_k^{(N)}(x) t^k. \quad (7)$$

It should be noticed, however, that some useful closed-form expressions carrying information about the discrete Chebyshev polynomials have been found before. For instance in [5] an expression

$$(1+t)^k (1+s)^{N-x} (1-st)^x \quad (8)$$

having the property that the coefficient of  $s^k t^k$  equals to  $D_k^{(N)}(x)$  is given.

## 2.2 Jacobi Differential Equation

For a natural number  $n$ , the *Jacobi polynomial*  $P_n^{(\alpha, \beta)}(x)$  is, up to the constant factor, the unique entire rational solution to the *Jacobi differential equation*

$$(1-x^2)y'' + (\beta - \alpha - (\alpha + \beta + 2)x)y' + n(n + \alpha + \beta + 1)y = 0 \quad (9)$$

(see [1]).

In this article, we are interested in Jacobi polynomials with parameters  $\alpha = 0$ ,  $\beta = -(N + 1)$ , where  $N > 0$  is a fixed integer. We also substitute  $x$  for  $n$  and  $t$  for  $x$  in equation (9), and denote  $J_x^{(N+1)}(t) = P_x^{(0, -N-1)}(t)$ . We usually omit superscript  $N + 1$  and denote  $J_x(t) = J_x^{(N+1)}(t)$ . Then  $J_x(t)$  satisfies differential equation

$$(1 - t^2)J_x''(t) - (N + 1 - (N - 1)t)J_x'(t) + x(x - N)J_x(t) = 0. \quad (10)$$

Recall that in this context,  $x$  is a fixed nonnegative integer. Polynomial  $J(t)$  can be expressed as

$$J_x(t) = \frac{1}{2^x} \sum_{k=0}^x \binom{x}{k} \binom{x - N - 1}{k} (t - 1)^k (t + 1)^{x-k} \quad (11)$$

(coefficient  $\frac{1}{2^x}$  is chosen for a special purpose) [1]. Since equation (10) is clearly invariant under substitution  $x \leftarrow N - x$ , we have symmetry

$$J_{N-x}(t) = J_x(t) \quad (12)$$

(see [1]).

### 2.3 MacWilliams Transform

The *MacWilliams transform* of order  $x$  for a polynomial  $P$  is defined as

$$\widehat{P}_x(t) = (1 + t)^x P\left(\frac{1 - t}{1 + t}\right). \quad (13)$$

As definition (13) shows, MacWilliams transform is a special case of Möbius transformation together with factor  $(1 + t)^x$ . If the subscript  $x$  is clear by context, we may omit it. It is also straightforward to see that if  $x$  is an integer so that  $\deg(P) \leq x$ , then  $\widehat{P}$  is again a polynomial. In this article, we will however face situations with non-integer values of  $x$ , and it is worth noticing already here that (13) shows that if  $t > -1$ , then  $\widehat{P}_x(t)$  is a uniquely defined differentiable function of real variable  $x$ .

In what follows,  $\widehat{J}_x(t)$  stands for the MacWilliams transform of  $J_x$  of order  $x$ . It is then straightforward to uncover a representation for  $\widehat{J}_x(t)$ :

$$\widehat{J}_x(t) = (\widehat{J_x})_x(t) = \sum_{k=0}^x (-1)^k \binom{x}{k} \binom{x - N - 1}{k} t^k. \quad (14)$$

The symmetry (12) implies straightforwardly

$$\begin{aligned} \widehat{J}_{N-x}(t) &= (\widehat{J_{N-x}})_{N-x}(t) = (1 + t)^{N-x} J_{N-x}\left(\frac{1 - t}{1 + t}\right) \\ &= (1 + t)^{N-2x} (1 + t)^x J_x\left(\frac{1 - t}{1 + t}\right) = (1 + t)^{N-2x} \widehat{J}_x(t). \end{aligned}$$

Equality

$$\widehat{J}_{N-x}(t) = (1 + t)^{N-2x} \widehat{J}_x(t) \quad (15)$$

thus obtained will be important in understanding the alternative representation of the generating function introduced in Section 5.

### 3 Heun Equation

A differential equation for the MacWilliams transform of  $J_x(t)$  can be found easily. For short, we denote  $J(t) = J_x(t)$  and  $\widehat{J}(t) = \widehat{J}_x(t)$  in the following lemmata.

**Lemma 1.**  $\widehat{J}(t)$  satisfies differential equation

$$t(1+t)\widehat{J}''(t) + (Nt+1-2t(x-1))\widehat{J}'(t) + x(x-N-1)\widehat{J}(t) = 0. \quad (16)$$

*Proof.* By computing the derivatives of  $\widehat{J}(t) = (1+t)^x J(\frac{1-t}{1+t})$  we can represent  $\widehat{J}(t)$ ,  $\widehat{J}'(t)$ , and  $\widehat{J}''(t)$  in terms of  $J(\frac{1-t}{1+t})$ ,  $J'(\frac{1-t}{1+t})$ , and  $J''(\frac{1-t}{1+t})$ . A direct calculation allows us also to reverse the representations to get

$$J\left(\frac{1-t}{1+t}\right) = (1+t)^{-x} \widehat{J}(t), \quad (17)$$

$$J'\left(\frac{1-t}{1+t}\right) = \frac{1}{2}x(1+t)^{-x+1}\widehat{J}(t) - \frac{1}{2}(1+t)^{-x+2}\widehat{J}'(t), \text{ and} \quad (18)$$

$$\begin{aligned} J''\left(\frac{1-t}{1+t}\right) &= \frac{1}{4}x(x-1)(1+t)^{-x+2}\widehat{J}(t) \\ &\quad - \frac{1}{2}(x-1)(1+t)^{-x+3}\widehat{J}'(t) + \frac{1}{4}(1+t)^{-x+4}\widehat{J}''(t). \end{aligned} \quad (19)$$

Replacing  $t$  with  $\frac{1-t}{1+t}$  in (10) and substituting (17)-(19) into (10) gives us the claim.

Another way to prove the lemma is to use (14) and verify by direct calculations that differential equation (16) is satisfied.  $\square$

**Lemma 2.** Let  $T(t)$  be defined as  $T(t) = (1+t)^{N-2x}\widehat{J}(-t^2)$ . Then  $T(t)$  satisfies differential equation

$$\begin{aligned} (t^3 - t)T''(t) &+ (2t(N-2x) + 3t^2 - 1)T'(t) \\ &+ (N - 2x - tN(N+2))T(t) = 0 \end{aligned} \quad (20)$$

*Proof.* As in the previous Lemma, we can express  $T(t)$ ,  $T'(t)$ , and  $T''(t)$  in terms of  $\widehat{J}(-t^2)$ ,  $\widehat{J}'(-t^2)$ , and  $\widehat{J}''(-t^2)$ , and then to reverse the representations to get

$$\widehat{J}(-t^2) = (1+t)^{2x-N}T(t) \quad (21)$$

$$\begin{aligned} \widehat{J}'(-t^2) &= \frac{1}{2t}(N-2x)(1+t)^{2x-N-1}T(t) \\ &\quad - \frac{1}{2t}(1+t)^{2x-N}T'(t) \end{aligned} \quad (22)$$

$$\begin{aligned} \widehat{J}''(-t^2) &= \frac{1}{4t^3}(N-2x)(1+t)^{2x-N-2}(t(N-2x+2)+1)T(t) \\ &\quad - \frac{1}{4t^3}(2t(N-2x)+t+1)(1+t)^{2x-N-1}T'(t) \\ &\quad + \frac{1}{4t^2}(1+t)^{2x-N}T''(t) \end{aligned} \quad (23)$$

by direct calculation. By substituting  $-t^2$  for  $t$  in (16) and by using (21)-(23), we get differential equation (20) after some direct calculations.  $\square$



Differential equation (20) is a special case of *Heun differential equation*

$$t(t-1)(t-q)y''(t) + (c(t-1)(t-q) + d \cdot t(t-q) + (a+b+1-c-d)t(t-1))y'(t) + (abt - \lambda)y(t) = 0$$

(see [7]). Differential equation (20) can be obtained by taking  $q = -1$ ,  $a = -N$ ,  $b = N + 2$ ,  $c = 1$ ,  $d = N - 2x + 1$ , and  $\lambda = 2x - N$ .

## 4 The Generating Function

By equality (14) function  $T(t) = (1+t)^{N-2x} \widehat{J}_x(-t^2)$  can be represented as

$$T(t) = (1+t)^{N-2x} \sum_{k=0}^x \binom{x}{k} \binom{x-N-1}{k} t^{2k}. \quad (24)$$

If  $t \in (-1, 1)$ , we should keep in mind that  $\widehat{J}_x(-t^2) = (1+t^2)^x J(\frac{1+t^2}{1-t^2})$  can be straightforwardly defined for any real values of  $x$ . Hence for  $t \in (-1, 1)$  also  $T(t) = (1+t)^{N-2x} \widehat{J}_x(-t^2)$  can be defined for an arbitrary real  $x$ , even though (24) is meaningful only for integer values of  $x$ . Another way of generalizing (24) even to complex values of  $x$  is to expand (24) straightforwardly to see that if we write

$$T(t) = \sum_{k=0}^{\infty} \tau_k(x) t^k, \quad (25)$$

then

$$\tau_k(x) = \sum_{0 \leq l \leq k/2} \binom{N-2x}{k-2l} \binom{x}{l} \binom{x-N-1}{l} \quad (26)$$

is a polynomial of degree  $k$ . For any fixed  $x$ ,  $T(t)$  is an analytic function of  $t$  in the disc  $|t| < 1$  (we can use the principal branch of the logarithm to defined to power), and hence it has a unique Maclaurin expansion (25) convergent when  $|t| < 1$ .

That (25) converges for  $|t| < 1$  can be also verified by using the ratio test, but to estimate  $|\tau_{k+1}(x)/\tau_k(x)|$  as  $k$  tends to infinity is not very straightforward. On the other hand, the recurrence of the next lemma reveals that  $\lim_{k \rightarrow \infty} |\tau_{k+1}(x)/\tau_k(x)| = 1$ .

**Remark 1.** Polynomials  $\tau_k(x)$  for small values of  $k$  are easy to find by using (26). For instance,  $\tau_0(x) = 1$ ,  $\tau_1(x) = N - 2x$ , and  $\tau_2(x) = 3x^2 - 3Nx + \frac{1}{2}N(N-1)$ .

**Lemma 3.** For  $k \geq 2$ , polynomials  $\tau_k(x)$  satisfy the recurrence relation

$$k^2 \tau_k(x) = (2k-1)(N-2x)\tau_{k-1}(x) - (N+k)(N-k+2)\tau_{k-2}(x). \quad (27)$$

*Proof.* This is a general property for a generic solution to Heun equation, see [7]. Recurrence (27) can be also obtained by differentiating and substituting (25) to Equation (20).  $\square$

**Remark 2.** From (27) it follows that

$$\frac{\tau_k(x)}{\tau_{k-1}(x)} = \frac{(2k-1)(N-2x)}{k^2} - \frac{(N+k)(N-k+2)}{k^2} \frac{\tau_{k-2}(x)}{\tau_{k-1}(x)},$$

which shows that  $\lim_{k \rightarrow \infty} |\tau_{k+1}(x)/\tau_k(x)| = \infty$  is impossible. Since clearly  $\tau_k(x)$  is a rational expression in  $k$ , the limit exists and is finite. Now

$$\frac{\tau_k(x)}{\tau_{k-1}(x)} \cdot \frac{\tau_{k-1}(x)}{\tau_{k-2}(x)} = \frac{(2k-1)(N-2x)}{k^2} \cdot \frac{\tau_{k-1}(x)}{\tau_{k-2}(x)} - \frac{(N+k)(N-k+2)}{k^2}$$

shows that  $\lim_{k \rightarrow \infty} |\tau_{k+1}(x)/\tau_k(x)| = 1$ .

We are now ready to state the main result.

**Theorem 1.** *Function*

$$T_{N,x}(t) = (1+t)^{N-2x} \widehat{J}_x(-t^2) \quad (28)$$

is the generating function of discrete Chebyshev polynomials, i.e.  $\tau_k(x) = D_k(x)$  for each  $k \geq 0$ .

*Proof.* By (5), the Discrete Chebyshev polynomials satisfy the same recurrence relation (27) as polynomials  $\tau_k(x)$  do. Since the initial conditions  $\tau_0(x) = D_0(x)$  and  $\tau_1(x) = D_1(x)$  hold by Remark 1, we have equality  $\tau_k(x) = D_k(x)$  for each  $k$ .  $\square$

**Remark 3.** It may be useful to compare (28) and (6).

## 5 Further Remarks

**Example 1.** Expression (28) shows that if  $x$  is an integer at most  $N/2$ , then  $T_{N,x}(t)$  is a polynomial in  $t$  of degree  $N - 2x + 2x = N$ . Thus we can find expressions

$$T_{N,x}(t) = \sum_{n=0}^N D_n^{(N)}(x) t^n$$

by simply evaluating  $D_n(N)(x)$  for  $n \in \{0, 1, \dots, N\}$  by using (5) or (26). For example,  $N = 6$  gives

$$\begin{aligned} T_{6,0}(t) &= 1 + 6t + 15t^2 + 20t^3 + 15t^4 + 6t^5 + t^6 = (1+t)^6 \\ T_{6,1}(t) &= 1 + 4t + 0 \cdot t^2 - 20t^3 - 35t^4 - 24t^5 - 6t^6 = (1+t)^4(1-6t^2) \\ T_{6,2}(t) &= 1 + 2t - 9t^2 - 20t^3 + 5t^4 + 30t^5 + 15t^6 \\ &= (1+t)^2(1-10t^2+15t^4) \\ T_{6,3}(t) &= 1 - 12t^2 + 30t^4 - 20t^6, \end{aligned}$$

which is in full accordance with (28) and (14). For  $x \in \{4, 5, 6\}$  the power  $6 - 2x$  of  $1 + t$  in (28) is no longer positive, so it is not clear that  $T_{6,x}(t)$  would be a polynomial anymore. But if  $T_{6,x}$  is not a polynomial for  $x \in \{4, 5, 6\}$ , there would be a rather mysterious asymmetry between  $x \leq 3$  and  $x > 3$ . Fortunately it is easy to show that  $T_{N,x}(t)$  is indeed a polynomial for each  $x \in \{0, 1, \dots, N\}$  and the asymmetry actually vanishes via trivial equality  $1 - t^2 = (1+t)(1-t)$ .

**Theorem 2.** *The generating function  $T_{N,x}(t)$  can be also represented as*

$$T_{N,x}(t) = (1-t)^{2x-N} \widehat{J}_{N-x}(-t^2). \quad (29)$$

*Proof.* Equality (15) implies

$$\widehat{J}_{N-x}(-t^2) = (1-t^2)^{N-2x} \widehat{J}_x(-t^2) = (1-t)^{N-2x} (1+t)^{N-2x} \widehat{J}_x(-t^2),$$

and the claim follows immediately.  $\square$

**Example 2** (Example 1 continued). Since by Theorem 2  $T_{6,x}(t)$  is a polynomial in  $t$  of degree 6, we can evaluate its values for  $x \in \{4, 5, 6\}$  as

$$\begin{aligned} T_{6,4}(t) &= 1 - 2t - 9t^2 + 20t^3 + 5t^4 - 30t^5 + 15t^6 \\ &= (1-t)^2(1-10t^2+15t^4) \\ T_{6,5}(t) &= 1 - 4t + 0 \cdot t^2 + 20t^3 - 35t^4 + 24t^5 - 6t^6 = (1-t)^4(1-6t^2) \\ T_{6,6}(t) &= 1 - 6t + 15t^2 - 20t^3 + 15t^4 - 6t^5 + t^6 = (1-t)^6. \end{aligned}$$

This is again in full accordance with (29) and (14).

To combine Theorems 1 and 2 into a single presentation is straightforward:

**Theorem 3** (The explicit polynomial form for  $x \in \{0, 1, \dots, N\}$ ). *The generating function  $T_{N,x}(t)$  can be presented as*

$$T_{N,x}(t) = (1+t \cdot \text{sign}(N-2x))^{|N-2x|} \widehat{J}_{\min\{x, N-x\}}^{(N)}(-t^2).$$

**Remark 4.** Theorem 1 implies that (5) and (26) are equal, i.e.

$$\sum_{0 \leq l \leq k/2} \binom{N-2x}{k-2l} \binom{x}{l} \binom{x-N-1}{l} = \sum_{l=0}^k (-1)^l \binom{k}{l} \binom{N-x}{k-l} \binom{x}{l}. \quad (30)$$

A direct combinatorial proof of (30) appears challenging. Theorem 2 implies an identity similar to (30).

## References

- [1] H. Bateman and A. Erdelyi: *Higher Transcendental Functions Vol 2.* McGraw-Hill (1953).
- [2] M. Hirvensalo: *Studies on Boolean Functions Related to Quantum Computing.* Ph.D thesis, University of Turku (2003).
- [3] N. Gogin and M. Hirvensalo: *Recurrent Construction of MacWilliams and Chebyshev Matrices.* TUCS Technical Report 812 (2007).
- [4] A.N. Kolmogorov and S.V. Fomin: *Introductory Real Analysis.* Dover (1975).

- [5] T. Laihonen: *Estimates on the Covering Radius When the Dual Distance is Known*. Ph.D thesis, University of Turku (1998).
- [6] F.J. MacWilliams and N.J.A. Sloane: *The Theory of Error-Correcting Codes*. North-Holland (1977).
- [7] S. Yu. Slavyanov and W. Lay: *Special Functions, A Unified Theory Based on Singularities*, Oxford University Press, 2000.
- [8] G. Szegő: *Orthogonal Polynomials*. Providence, Rhode Island: American Mathematical Society, Colloquium Publications, Volume XXIII (1975).



The logo features a dark blue background with several thin, white, abstract lines that form a network-like structure, resembling a stylized map or a complex diagram. The text is positioned on the left side of this blue area.

TURKU  
CENTRE *for*  
COMPUTER  
SCIENCE

Lemminkäisenkatu 14 A, 20520 Turku, Finland | [www.tucs.fi](http://www.tucs.fi)



**University of Turku**

- Department of Information Technology
- Department of Mathematics



**Åbo Akademi University**

- Department of Computer Science
- Institute for Advanced Management Systems Research



**Turku School of Economics and Business Administration**

- Institute of Information Systems Sciences

ISBN 978-952-12-1896-5  
ISSN 1239-1891