

# Expressiveness of the Interval Logics of Allen’s Relations on the Class of all Linear Orders: Complete Classification\*

Dario Della Monica<sup>†</sup>  
University of Udine, Italy

Valentin Goranko  
Technical University of Denmark

Angelo Montanari  
University of Udine, Italy

Guido Sciavicco

University of Murcia, Spain, and UIST Ohrid, Macedonia

## Abstract

We compare the expressiveness of the fragments of Halpern and Shoham’s interval logic (HS), i.e., of all interval logics with modal operators associated with Allen’s relations between intervals in linear orders. We establish a complete set of inter-definability equations between these modal operators, and thus obtain a complete classification of the family of  $2^{12}$  fragments of HS with respect to their expressiveness. Using that result and a computer program, we have found that there are 1347 expressively different such interval logics over the class of all linear orders.

## 1 Introduction

Interval reasoning naturally arises in various fields of artificial intelligence, such as theories of action and change, natural language analysis and processing, and constraint satisfaction problems. Interval temporal logics formalize reasoning about interval structures over ordered domains, where time intervals, rather than time instants, are the primitive ontological entities. The variety of binary relations between intervals in linear orders was first studied systematically by Allen [Allen, 1983], who explored their use in systems for time management and planning. The modal logic featuring modal operators corresponding to Allen’s interval relations was introduced by Halpern and Shoham in [Halpern and Shoham, 1991]; we hereafter call that logic HS. Temporal logics with interval-based semantics have also been proposed as a suitable formalism for the specification and verification of hardware [Moszkowski, 1983] and of real-time systems [Zhou and Hansen, 2004].

In [Halpern and Shoham, 1991], it was shown that the satisfiability problem for HS is undecidable in all natural classes

of linear orders. For a long time, these sweeping undecidability results have discouraged attempts for practical applications of interval logics. A renewed interest in the area has recently been stimulated by the discovery of several interesting decidable fragments of HS [Bresolin *et al.*, 2007a; 2007b; 2008; 2009; 2010; Montanari *et al.*, 2010a; 2010b]. In that context, and for the purpose of identifying expressive interval logics for various intended applications, the comparative analysis of the expressiveness of the variety of interval logics is a major research problem in the area. In particular, the important problem arises to analyze the mutual definabilities among the modal operators of the logic HS and to classify the fragments of HS with respect to their expressiveness.

In the present paper we address and solve that problem, by identifying a complete set of inter-definability formulae among the modal operators of HS and thus providing a complete classification of all fragments of HS with respect to their expressiveness for the *strict* semantics (excl. point intervals) over the class of all linear orders. Using that result we have found that there are exactly 1347 expressively different such fragments out of  $2^{12} = 4096$  sets of modal operators in HS.

The choice of strict semantics, excluding point intervals, instead of including them (non-strict semantics), conforms to the definition of interval adopted by Allen in [Allen, 1983]. It has at least two strong motivations. First, a number of representation paradoxes arise when the non-strict semantics is adopted, due to the presence of point intervals, as pointed out in [Allen, 1983]. Second, when point intervals are included, there seems to be no intuitive semantics for interval relations that makes them both pairwise disjoint and jointly exhaustive.

The structure of the paper: after the preliminary Section 2, in Section 3 we state the main result of the paper, and we prove that the proposed set of inter-definability equations is correct. The much more difficult proof of completeness is given in Section 4. Section 5 provides an assessment of the work done and it outlines future research directions.

## 2 Preliminaries

Let  $\mathbb{D} = \langle D, < \rangle$  be a linearly ordered set. An *interval* over  $\mathbb{D}$  is an ordered pair  $[a, b]$ , where  $a, b \in D$  and  $a \leq b$ . Intervals of the type  $[a, a]$  are called *point intervals*, while the others are called *strict intervals*. There are 12 different non-trivial relations (excluding the equality) between two strict intervals

\*We would like to acknowledge the Spanish project TIN2009-14372-C03-01 (G. Sciavicco), the HYLOCORE project, funded by the Danish Natural Science Research Council (V. Goranko), the Italian PRIN project *Innovative and multi-disciplinary approaches for constraint and preference reasoning* (D. Della Monica, A. Montanari), and the Spanish/South-African Project HS2008-0006 (V. Goranko, G. Sciavicco).

<sup>†</sup>Currently affiliated with University of Salerno, Italy.

$\langle A \rangle$	$[a, b]R_A[c, d] \Leftrightarrow b = c$	
$\langle L \rangle$	$[a, b]R_L[c, d] \Leftrightarrow b < c$	
$\langle B \rangle$	$[a, b]R_B[c, d] \Leftrightarrow a = c, d < b$	
$\langle E \rangle$	$[a, b]R_E[c, d] \Leftrightarrow b = d, a < c$	
$\langle D \rangle$	$[a, b]R_D[c, d] \Leftrightarrow a < c, d < b$	
$\langle O \rangle$	$[a, b]R_O[c, d] \Leftrightarrow a < c < b < d$	

Table 1: Allen’s interval relations and the corresponding HS modalities.

in a linear order, often called *Allen’s relations* [Allen, 1983]: the six relations depicted in Table 1 and the inverse relations.

We treat interval structures as Kripke structures and Allen’s relations as accessibility relations in them, thus associating a modal operator  $\langle X \rangle$  with each Allen’s relation  $R_X$ . For each operator  $\langle X \rangle$ , its *transpose*, denoted by  $\langle \bar{X} \rangle$ , corresponds to the inverse relation  $R_{\bar{X}}$  of  $R_X$  (that is,  $R_{\bar{X}} = (R_X)^{-1}$ ).

Halpern and Shoham’s logic HS is a multi-modal logic with formulae built over a set  $\mathcal{AP}$  of propositional letters, the propositional connectives  $\vee$  and  $\neg$ , and a set of modal operators associated with all Allen’s relations. With every subset  $\{R_{X_1}, \dots, R_{X_k}\}$  of these relations, we associate the fragment  $X_1X_2 \dots X_k$  of HS, the formulae of which are defined by the grammar:

$$\varphi ::= p \mid \neg\varphi \mid \varphi \vee \psi \mid \langle X_1 \rangle \varphi \mid \dots \mid \langle X_k \rangle \varphi.$$

The other propositional connectives,  $\wedge$  and  $\rightarrow$ , and the dual operators  $[X]$  are defined as usual, e.g.,  $[X]\varphi \equiv \neg\langle X \rangle\neg\varphi$ .

For a fragment  $\mathcal{F} = X_1X_2 \dots X_k$  and a modal operator  $\langle X \rangle$ , we write  $\langle X \rangle \in \mathcal{F}$  if  $X \in \{X_1, \dots, X_k\}$ . Given two fragments  $\mathcal{F}_1$  and  $\mathcal{F}_2$ , we write  $\mathcal{F}_1 \subseteq \mathcal{F}_2$  if  $\langle X \rangle \in \mathcal{F}_1$  implies  $\langle X \rangle \in \mathcal{F}_2$ , for every modal operator  $\langle X \rangle$ .

The semantics of HS is given in terms of *interval models*  $M = \langle \mathbb{I}(\mathbb{D}), V \rangle$ , where  $\mathbb{I}(\mathbb{D})$  is the set of all (strict) intervals over  $\mathbb{D}$ . The *valuation function*  $V : \mathcal{AP} \mapsto 2^{\mathbb{I}(\mathbb{D})}$  assigns to every  $p \in \mathcal{AP}$  the set of intervals  $V(p)$  on which  $p$  holds. The *truth* of a formula on a given interval  $[a, b]$  in an interval model  $M$  is defined by structural induction on formulae:

- $M, [a, b] \Vdash p$  iff  $[a, b] \in V(p)$ , for all  $p \in \mathcal{AP}$ ;
- $M, [a, b] \Vdash \neg\psi$  iff it is not the case that  $M, [a, b] \Vdash \psi$ ;
- $M, [a, b] \Vdash \varphi \vee \psi$  iff  $M, [a, b] \Vdash \varphi$  or  $M, [a, b] \Vdash \psi$ ;
- $M, [a, b] \Vdash \langle X \rangle \psi$  iff there exists an interval  $[c, d]$  such that  $[a, b]R_X[c, d]$  and  $M, [c, d] \Vdash \psi$ , where  $R_X$  is any of Allen’s relations.

A formula  $\phi$  of HS is *valid*, denoted  $\models \phi$ , if it is true on every interval in every interval model. Two formulae  $\phi$  and  $\psi$  are *equivalent*, denoted  $\phi \equiv \psi$ , if  $\models \phi \leftrightarrow \psi$ .

**Definition 2.1.** A modal operator  $\langle X \rangle$  of HS is *definable* in an HS-fragment  $\mathcal{F}$ , denoted  $\langle X \rangle \triangleleft \mathcal{F}$ , if  $\langle X \rangle p \equiv \psi$  for some formula  $\psi = \psi(p) \in \mathcal{F}$ , for any fixed propositional variable  $p$ . In such a case, the equivalence  $\langle X \rangle p \equiv \psi$  is called an *inter-definability equation for  $\langle X \rangle$  in  $\mathcal{F}$* .

It is known from [Halpern and Shoham, 1991] that, in the strict semantics, all modal operators in HS are definable in

the fragment containing the modalities  $\langle A \rangle$ ,  $\langle B \rangle$ , and  $\langle E \rangle$ , and their transposes  $\langle \bar{A} \rangle$ ,  $\langle \bar{B} \rangle$ , and  $\langle \bar{E} \rangle$  (In the non-strict semantics, the four modalities  $\langle B \rangle$ ,  $\langle E \rangle$ ,  $\langle \bar{B} \rangle$ , and  $\langle \bar{E} \rangle$  suffice, as shown in [Venema, 1990]).

In this paper, we compare and classify the expressiveness of all fragments of HS on the class of all interval structures over linear orders. Formally, let  $\mathcal{F}_1$  and  $\mathcal{F}_2$  be any pair of such fragments. We say that:

- $\mathcal{F}_2$  is *at least as expressive as*  $\mathcal{F}_1$ , denoted  $\mathcal{F}_1 \preceq \mathcal{F}_2$ , if every operator  $\langle X \rangle \in \mathcal{F}_1$  is definable in  $\mathcal{F}_2$ .
- $\mathcal{F}_1$  is *strictly less expressive than*  $\mathcal{F}_2$ , denoted  $\mathcal{F}_1 \prec \mathcal{F}_2$ , if  $\mathcal{F}_1 \preceq \mathcal{F}_2$  but not  $\mathcal{F}_2 \preceq \mathcal{F}_1$ .
- $\mathcal{F}_1$  and  $\mathcal{F}_2$  are *equally expressive* (or, *expressively equivalent*), denoted  $\mathcal{F}_1 \equiv \mathcal{F}_2$ , if  $\mathcal{F}_1 \preceq \mathcal{F}_2$  and  $\mathcal{F}_2 \preceq \mathcal{F}_1$ .
- $\mathcal{F}_1$  and  $\mathcal{F}_2$  are *expressively incomparable*, denoted  $\mathcal{F}_1 \not\equiv \mathcal{F}_2$ , if neither  $\mathcal{F}_1 \preceq \mathcal{F}_2$  nor  $\mathcal{F}_2 \preceq \mathcal{F}_1$ .

In order to show non-definability of a given modal operator in a given fragment, we use a standard technique in modal logic, based on the notion of *bisimulation* and the invariance of modal formulae with respect to bisimulations (see, e.g., [Blackburn *et al.*, 2002]). Let  $\mathcal{F}$  be an HS-fragment. An  $\mathcal{F}$ -bisimulation between two interval models  $M = \langle \mathbb{I}(\mathbb{D}), V \rangle$  and  $M' = \langle \mathbb{I}(\mathbb{D}'), V' \rangle$  over  $\mathcal{AP}$  is a relation  $Z \subseteq \mathbb{I}(\mathbb{D}) \times \mathbb{I}(\mathbb{D}')$  satisfying the following properties:

- *local condition*:  $Z$ -related intervals satisfy the same propositional letters over  $\mathcal{AP}$ ;
- *forward condition*: if  $([a, b], [a', b']) \in Z$  and  $([a, b], [c, d]) \in R_X$  for some  $\langle X \rangle \in \mathcal{F}$ , then there exists  $[c', d']$  such that  $([a', b'], [c', d']) \in R_X$  and  $([c, d], [c', d']) \in Z$ ;
- *backward condition*: likewise, but from  $M'$  to  $M$ .

The important property of bisimulations used here is that any  $\mathcal{F}$ -bisimulation preserves the truth of *all* formulae in  $\mathcal{F}$ . Thus, in order to prove that an operator  $\langle X \rangle$  is not definable in  $\mathcal{F}$ , it suffices to construct a pair of interval models  $M$  and  $M'$  and a  $\mathcal{F}$ -bisimulation between them, relating a pair of intervals  $[a, b] \in M$  and  $[a', b'] \in M'$ , such that  $M, [a, b] \Vdash \langle X \rangle p$ , while  $M', [a', b'] \not\Vdash \langle X \rangle p$ .

### 3 Comparing the expressiveness of the fragments of HS

In order to classify all fragments of HS with respect to their expressiveness, it suffices to identify all definabilities of modal operators  $\langle X \rangle$  in fragments  $\mathcal{F}$ , where  $\langle X \rangle \notin \mathcal{F}$ .

A definability  $\langle X \rangle \triangleleft \mathcal{F}$  is *optimal* if  $\langle X \rangle \not\triangleleft \mathcal{F}'$  for any fragment  $\mathcal{F}'$  such that  $\mathcal{F}' \prec \mathcal{F}$ . A set of such definabilities is *optimal* if it consists of optimal definabilities.

The main result of the paper is the following theorem.

**Theorem 3.1.** *The set of inter-definability equations given in Table 2 is sound, complete, and optimal.*

Most of the equations in Table 2 are known from [Halpern and Shoham, 1991], except the definability  $\langle L \rangle \triangleleft \bar{B}\bar{E}$  and its symmetric,  $\langle \bar{L} \rangle \triangleleft B\bar{E}$ , which are new. We will first prove the soundness of the given set of inter-definability equations.

**Lemma 3.2.** *The set of inter-definability equations given in Table 2 is sound.*

$\langle L \rangle p \equiv \langle A \rangle \langle A \rangle p$	$\langle L \rangle \triangleleft A$
$\langle \bar{L} \rangle p \equiv \langle \bar{A} \rangle \langle \bar{A} \rangle p$	$\langle \bar{L} \rangle \triangleleft \bar{A}$
$\langle O \rangle p \equiv \langle E \rangle \langle \bar{B} \rangle p$	$\langle O \rangle \triangleleft \bar{B}E$
$\langle \bar{O} \rangle p \equiv \langle \bar{B} \rangle \langle \bar{E} \rangle p$	$\langle \bar{O} \rangle \triangleleft \bar{B}\bar{E}$
$\langle D \rangle p \equiv \langle E \rangle \langle B \rangle p$	$\langle D \rangle \triangleleft BE$
$\langle \bar{D} \rangle p \equiv \langle \bar{E} \rangle \langle \bar{B} \rangle p$	$\langle \bar{D} \rangle \triangleleft \bar{B}\bar{E}$
$\langle L \rangle p \equiv \langle \bar{B} \rangle [E] \langle \bar{B} \rangle \langle E \rangle p$	$\langle L \rangle \triangleleft \bar{B}\bar{E}$
$\langle \bar{L} \rangle p \equiv \langle \bar{E} \rangle [B] \langle \bar{E} \rangle \langle B \rangle p$	$\langle \bar{L} \rangle \triangleleft \bar{B}\bar{E}$

Table 2: The complete set of inter-definability equations

*Proof.* We only need to prove the soundness for the new inter-definability equations  $\langle L \rangle p \equiv \langle \bar{B} \rangle [E] \langle \bar{B} \rangle \langle E \rangle p$  and its symmetric for  $\langle \bar{L} \rangle$ . The proofs are analogous, so we only prove the former. First, we prove the left-to-right direction. Suppose that  $M, [a, b] \Vdash \langle L \rangle p$  for some model  $M$  and interval  $[a, b]$ . This means that there exists an interval  $[c, d]$  such that  $b < c$  and  $M, [c, d] \Vdash p$ . We exhibit an interval  $[a, y]$ , with  $y > b$  such that, for every  $x$  (strictly) in between  $a$  and  $y$ , the interval  $[x, y]$  is such that there exist two points  $y'$  and  $x'$  such that  $y' > y$ ,  $x < x' < y'$ , and  $[x', y']$  satisfies  $p$ . Let  $y$  be equal to  $c$ . The interval  $[a, c]$ , which is started by  $[a, b]$ , is such that for any of its ending intervals, that is, for any interval of the form  $[x, c]$ , with  $a < x$ , we have that  $x < c < d$  and  $M, [c, d] \Vdash p$ . As for the other direction, we must show that  $\langle \bar{B} \rangle [E] \langle \bar{B} \rangle \langle E \rangle p$  implies  $\langle L \rangle p$ . To this end, suppose that  $M, [a, b] \Vdash \langle \bar{B} \rangle [E] \langle \bar{B} \rangle \langle E \rangle p$  for a model  $M$  and an interval  $[a, b]$ . Then, there exists an interval  $[a, c]$ , for some  $c > b$ , such that  $[E] \langle \bar{B} \rangle \langle E \rangle p$  is true on  $[a, c]$ . As a consequence, the interval  $[b, c]$  must satisfy  $\langle \bar{B} \rangle \langle E \rangle p$ , that means that there are two points  $x$  and  $y$  such that  $y > c$ ,  $b < x < y$ , and  $[x, y]$  satisfies  $p$ . Since  $x > b$ , then  $M, [a, b] \Vdash \langle L \rangle p$ .  $\square$

Proving completeness is the hard task; optimality will be established together with it. The completeness proof is organized as follows. For each HS operator  $\langle X \rangle$ , we show that  $\langle X \rangle$  is not definable in any fragment of HS that does not contain as definable (according to Table 2) all operators of some of the fragments in which  $\langle X \rangle$  is definable (according to Table 2). More formally, for each HS operator  $\langle X \rangle$ , the proof consists of the following steps:

1. using Table 2, find all fragments  $\mathcal{F}_i$  such that  $\langle X \rangle \triangleleft \mathcal{F}_i$ ;
2. identify the list  $\mathcal{M}_1, \dots, \mathcal{M}_m$  of all  $\subseteq$ -maximal fragments of HS that contain neither the operator  $\langle X \rangle$  nor any of the fragments  $\mathcal{F}_i$  identified by the previous step;
3. for each fragment  $\mathcal{M}_i$ , with  $i \in \{1, \dots, m\}$ , provide a bisimulation for  $\mathcal{M}_i$  which is not a bisimulation for  $X$ .

Details of the completeness proof will be provided in a series of lemmas (of increasing complexity) in the next section.

## 4 The completeness proof

In this section, we will prove that, for each modal operator  $\langle X \rangle$  of HS, the set of inter-definability equations for  $\langle X \rangle$  in Table 2 is complete for that operator, that is,  $\langle X \rangle$  is not definable in any fragment of HS that does not contain (as definable) all operators of some of the fragments listed in Table 2 in which  $\langle X \rangle$  is definable. Due to space limitations, we will

not prove in detail all the cases. A detailed proof can be found in the extended technical report and it will appear in a future journal version of the present paper.

### 4.1 Completeness for $\langle L \rangle$ and $\langle \bar{L} \rangle$

**Lemma 4.1.** *The set of inter-definability equations for  $\langle L \rangle$  and  $\langle \bar{L} \rangle$  given in Table 2 is complete.*

*Proof.* According to Table 2,  $\langle L \rangle$  is definable in terms of  $A$  and  $\bar{B}E$ . Hence, the fragments  $\text{BEDOALEDO}$  and  $\text{BDOALBEDO}$  are the only  $\subseteq$ -maximal ones not featuring  $\langle L \rangle$  and containing neither  $A$  nor  $\bar{B}E$ . To prove the thesis, it suffices to exhibit a bisimulation for each one of these two fragments that does not preserve the relation induced by  $\langle L \rangle$ . Thanks to Lemma 3.2,  $\text{BEDOALEDO}$  and  $\text{BDOALBEDO}$  are expressively equivalent to  $\text{BEOAED}$  and  $\text{BDOABE}$ , respectively. Thus, to all our purposes, we can simply refer to the latter ones instead of the former ones.

As for the first fragment, let  $M_1 = \langle \mathbb{I}(\mathbb{N}), V_1 \rangle$  and  $M_2 = \langle \mathbb{I}(\mathbb{N}), V_2 \rangle$  be two models and let  $V_1$  and  $V_2$  be such that  $V_1(p) = \{[2, 3]\}$  and  $V_2(p) = \emptyset$ , where  $p$  is the only propositional letter of the language. Moreover, let  $Z$  be a relation between (intervals of)  $M_1$  and  $M_2$  defined as  $Z = \{([0, 1], [0, 1])\}$ . It can be easily shown that  $Z$  is a  $\text{BEOAED}$ -bisimulation. The local property is trivially satisfied, since all  $Z$ -related intervals satisfy  $\neg p$ . As for the forward and backward conditions, it suffices to notice that, starting from the interval  $[0, 1]$ , it is not possible to reach any other interval using any of the modal operators of the fragment. At the same time,  $Z$  does not preserve the relation induced by the modality  $\langle L \rangle$ . Indeed,  $([0, 1], [0, 1]) \in Z$  and  $M_1, [0, 1] \Vdash \langle L \rangle p$ , but  $M_2, [0, 1] \Vdash \neg \langle L \rangle p$ . Therefore,  $\langle L \rangle$  is not definable in  $\text{BEDOALEDO}$ .

As for the second fragment, let  $M_1 = \langle \mathbb{I}(\mathbb{Z}^-), V_1 \rangle$  and  $M_2 = \langle \mathbb{I}(\mathbb{Z}^-), V_2 \rangle$  be two models based on the set  $\mathbb{Z}^- = \{\dots, -2, -1\}$ , and let  $V_1$  and  $V_2$  be such that  $V_1(p) = \{[-2, -1]\}$  and  $V_2(p) = \emptyset$ , where  $p$  is the only propositional letter of the language. Moreover, let  $Z$  be a relation between (intervals of)  $M_1$  and  $M_2$  defined as follows:  $([x, y], [w, z]) \in Z \stackrel{\text{def}}{\iff} [x, y] = [w, z]$  and  $[x, y] \neq [-2, -1]$ . We prove that  $Z$  is a  $\text{BDOABE}$ -bisimulation. First, the local property is trivially satisfied, since all  $Z$ -related intervals satisfy  $\neg p$ . Moreover, starting from any interval, the only interval that satisfies  $p$ , that is,  $[-2, -1]$ , cannot be reached using the set of modal operators featured by our fragment. At the same time,  $Z$  does not preserve the relation induced by  $\langle L \rangle$ , as  $([-4, -3], [-4, -3]) \in Z$  and  $M_1, [-4, -3] \Vdash \langle L \rangle p$ , but  $M_2, [-4, -3] \Vdash \neg \langle L \rangle p$ . Therefore,  $\langle L \rangle$  is not definable in  $\text{BDOALBEDO}$ .

A completely symmetric argument can be applied for the completeness proof of  $\langle \bar{L} \rangle$ .  $\square$

### 4.2 Completeness for $\langle E \rangle$ , $\langle \bar{E} \rangle$ , $\langle B \rangle$ , and $\langle \bar{B} \rangle$

**Lemma 4.2.** *The set of inter-definability equations for  $\langle E \rangle$ ,  $\langle \bar{E} \rangle$ ,  $\langle B \rangle$ , and  $\langle \bar{B} \rangle$  given in Table 2 is complete.*

*Proof.* According to Table 2, we will show that  $\langle E \rangle$  is not definable in terms of the only  $\subseteq$ -maximal fragment not fea-

turing it, namely,  $\overline{\text{ALBDOALBEDO}}$ . (The inverse modality  $\langle \overline{E} \rangle$  and the symmetric modalities  $\langle B \rangle$  and  $\langle \overline{B} \rangle$  can be dealt with using similar arguments.) Thanks to Lemma 3.2, it actually suffices to provide a bisimulation for  $\overline{\text{ABDOABE}}$ .

Let  $M_1 = \langle \mathbb{I}(\mathbb{R}), V_1 \rangle$  and  $M_2 = \langle \mathbb{I}(\mathbb{R}), V_2 \rangle$ , where  $p$  is the only propositional letter of the language, the valuation function  $V_1 : \mathcal{AP} \rightarrow 2^{\mathbb{I}(\mathbb{R})}$  is defined as:  $[x, y] \in V_1(p) \stackrel{\text{def}}{\iff} x \in \mathbb{Q} \text{ iff } y \in \mathbb{Q}$ , and the valuation function  $V_2 : \mathcal{AP} \rightarrow 2^{\mathbb{I}(\mathbb{R})}$  as:  $[w, z] \in V_2(p) \stackrel{\text{def}}{\iff} w \in \mathbb{Q} \text{ iff } z \in \mathbb{Q}$ , and  $([0, 3], [w, z]) \notin R_E$ . Moreover, let  $Z$  be a relation between (intervals of)  $M_1$  and  $M_2$  defined as follows:  $([x, y], [w, z]) \in Z \stackrel{\text{def}}{\iff} [x, y] \in V_1(p) \text{ iff } [w, z] \in V_2(p)$ .

We show that  $Z$  is an  $\overline{\text{ABDOABE}}$ -bisimulation between  $M_1$  and  $M_2$ . The satisfaction of the local condition immediately follows from the definition. The forward condition can be checked as follows. Let  $\overline{\mathbb{Q}} = \mathbb{R} \setminus \mathbb{Q}$  and let  $[x, y]$  and  $[w, z]$  be two  $Z$ -related intervals. For each modal operator  $\langle X \rangle$  of the language, let us assume that  $[x, y] R_X [x', y']$ . We have to exhibit an interval  $[w', z']$  such that  $[x', y']$  and  $[w', z']$  are  $Z$ -related, and  $[w, z]$  and  $[w', z']$  are  $R_X$ -related. We proceed case-by-case. Let  $\langle X \rangle = \langle A \rangle$  (and thus  $y = x'$ ). Suppose that  $[x', y'] \in V_1(p)$  (resp.,  $[x', y'] \notin V_1(p)$ ). We can always find a point  $z' > z$  such that  $[z, z'] \in V_2(p)$  (resp.,  $[z, z'] \notin V_2(p)$ ), independently from  $z$  belonging to  $\mathbb{Q}$  or  $\overline{\mathbb{Q}}$  (since both  $\mathbb{Q}$  and  $\overline{\mathbb{Q}}$  are right-unbounded). This implies that  $[x', y']$  and  $[z, z']$  are  $Z$ -related. Since  $[w, z]$  and  $[z, z']$  are obviously  $R_A$ -related, we have the thesis. If  $\langle X \rangle = \langle B \rangle$ , the argument is similar to the previous one, but, in this case, the density of  $\mathbb{Q}$  and  $\overline{\mathbb{Q}}$  is exploited. If  $\langle X \rangle = \langle D \rangle$ , it suffices to choose two points  $w'$  and  $z'$  such that  $w < w' < z' < z$ ,  $z' \neq 3$ ,  $w'$  belongs to  $\mathbb{Q}$  if and only if  $x'$  does, and  $z'$  belongs to  $\mathbb{Q}$  if and only if  $y'$  does. As in the previous case, the existence of such points is guaranteed by the density of  $\mathbb{Q}$  and  $\overline{\mathbb{Q}}$ . If  $\langle X \rangle = \langle O \rangle$ ,  $w'$  and  $z'$  are required to be such that  $w < w' < z < z'$ , and both density and right-unboundedness of  $\mathbb{Q}$  and  $\overline{\mathbb{Q}}$  must be exploited. The remaining cases as well as the backward condition can be verified in a very similar way. At the same time,  $Z$  does not preserve the relation induced by  $\langle E \rangle$ : we have that  $([0, 3], [0, 3]) \in Z$ ,  $M_1, [0, 3] \Vdash \langle E \rangle p$ , but  $M_2, [0, 3] \not\Vdash \langle E \rangle p$ . Therefore,  $\langle E \rangle$  cannot be defined in the fragment  $\overline{\text{ALBDOALBEDO}}$ .  $\square$

### 4.3 Completeness for $\langle A \rangle$ and $\langle \overline{A} \rangle$

In the proofs of Lemma 4.3 and Lemma 4.4, in order to get the bisimulation we want, we need to exploit a well-known property of the set of real numbers  $\mathbb{R}$ :  $\mathbb{R}$  (resp.,  $\mathbb{Q}$ ,  $\overline{\mathbb{Q}}$ ) can be partitioned into a countable number of pairwise disjoint subsets, each one of which is dense in  $\mathbb{R}$ . More formally, there are countably many nonempty sets  $\mathbb{R}_i$  (resp.,  $\mathbb{Q}_i$ ,  $\overline{\mathbb{Q}}_i$ ), with  $i \in \mathbb{N}$ , such that, for each  $i \in \mathbb{N}$ ,  $\mathbb{R}_i$  (resp.,  $\mathbb{Q}_i$ ,  $\overline{\mathbb{Q}}_i$ ) is dense in  $\mathbb{R}$ ,  $\mathbb{R} = \bigcup_{i \in \mathbb{N}} \mathbb{R}_i$  (resp.,  $\mathbb{Q} = \bigcup_{i \in \mathbb{N}} \mathbb{Q}_i$ ,  $\overline{\mathbb{Q}} = \bigcup_{i \in \mathbb{N}} \overline{\mathbb{Q}}_i$ ), and  $\mathbb{R}_i \cap \mathbb{R}_j = \emptyset$ , (resp.,  $\mathbb{Q}_i \cap \mathbb{Q}_j = \emptyset$ ,  $\overline{\mathbb{Q}}_i \cap \overline{\mathbb{Q}}_j = \emptyset$ ), for each  $i, j \in \mathbb{N}$  with  $i \neq j$ .

**Lemma 4.3.** *The set of inter-definability equations for  $\langle A \rangle$  and  $\langle \overline{A} \rangle$  given in Table 2 is complete.*

*Proof.* According to Table 2, it suffices to show that  $\langle A \rangle$  is not definable in the only  $\subseteq$ -maximal fragment not containing it, namely,  $\overline{\text{LBEDOALBEDO}}$ , which, by Lemma 3.2, turns out to be equivalent to  $\overline{\text{LBEABE}}$ .

Let  $M_1 = \langle \mathbb{I}(\mathbb{R}), V_1 \rangle$  and  $M_2 = \langle \mathbb{I}(\mathbb{R}), V_2 \rangle$  be two models built on the only propositional letter  $p$ . In order to define the valuation functions  $V_1$  and  $V_2$ , we take advantage of two partitions of the set  $\mathbb{R}$ , one for  $M_1$  and the other one for  $M_2$ , each of them consisting of exactly four sets that are dense in  $\mathbb{R}$ . Formally, for  $j = 1, 2$  and  $i = 1, \dots, 4$ , let  $\mathbb{R}_j^i$  be dense in  $\mathbb{R}$ . Moreover, for  $j = 1, 2$ , let  $\mathbb{R} = \bigcup_{i=1}^4 \mathbb{R}_j^i$  and  $\mathbb{R}_j^i \cap \mathbb{R}_j^{i'} = \emptyset$  for each  $i, i' \in \{1, 2, 3, 4\}$  with  $i \neq i'$ .

For  $j = 1, 2$ , we force points in  $\mathbb{R}_j^1$  (resp.,  $\mathbb{R}_j^2, \mathbb{R}_j^3, \mathbb{R}_j^4$ ) to behave in the same way with respect to the truth of  $p/\neg p$  over the intervals they initiate and terminate by imposing the following constraints:

$$\begin{aligned} &\forall x, y \text{ (if } x \in \mathbb{R}_j^1, \text{ then } M_j, [x, y] \Vdash \neg p); \\ &\forall x, y \text{ (if } x \in \mathbb{R}_j^2, \text{ then } M_j, [x, y] \Vdash \neg p); \\ &\forall x, y \text{ (if } x \in \mathbb{R}_j^3, \text{ then } (M_j, [x, y] \Vdash p \text{ iff } y \in \mathbb{R}_j^1 \cup \mathbb{R}_j^3)); \\ &\forall x, y \text{ (if } x \in \mathbb{R}_j^4, \text{ then } (M_j, [x, y] \Vdash p \text{ iff } y \in \mathbb{R}_j^2 \cup \mathbb{R}_j^4)). \end{aligned}$$

It can be easily shown that, from the given constraints, it immediately follows that:

$$\begin{aligned} &\forall x, y \text{ (if } y \in \mathbb{R}_j^1, \text{ then } (M_j, [x, y] \Vdash p \text{ iff } x \in \mathbb{R}_j^3)); \\ &\forall x, y \text{ (if } y \in \mathbb{R}_j^2, \text{ then } (M_j, [x, y] \Vdash p \text{ iff } x \in \mathbb{R}_j^4)); \\ &\forall x, y \text{ (if } y \in \mathbb{R}_j^3, \text{ then } (M_j, [x, y] \Vdash p \text{ iff } x \in \mathbb{R}_j^1)); \\ &\forall x, y \text{ (if } y \in \mathbb{R}_j^4, \text{ then } (M_j, [x, y] \Vdash p \text{ iff } x \in \mathbb{R}_j^2)). \end{aligned}$$

The above constraints univocally induces the following definition of the valuation functions  $V_j(p) : \mathcal{AP} \rightarrow 2^{\mathbb{I}(\mathbb{R})}$ :

$$[x, y] \in V_j(p) \stackrel{\text{def}}{\iff} \begin{cases} (x \in \mathbb{R}_j^3 \wedge y \in \mathbb{R}_j^1 \cup \mathbb{R}_j^3) \\ \vee (x \in \mathbb{R}_j^4 \wedge y \in \mathbb{R}_j^2 \cup \mathbb{R}_j^4). \end{cases}$$

Now, let  $Z$  be the relation between (intervals of)  $M_1$  and  $M_2$  defined as follows. Two intervals  $[x, y]$  and  $[w, z]$  are  $Z$ -related if and only if at least one of the following conditions holds:

1.  $x \in \mathbb{R}_1^1 \cup \mathbb{R}_2^1$  and  $w \in \mathbb{R}_1^2 \cup \mathbb{R}_2^2$ ;
2.  $x \in \mathbb{R}_1^3, w \in \mathbb{R}_2^3$ , and  $(y \in \mathbb{R}_1^1 \cup \mathbb{R}_1^3 \text{ iff } z \in \mathbb{R}_2^1 \cup \mathbb{R}_2^3)$ ;
3.  $x \in \mathbb{R}_1^3, w \in \mathbb{R}_2^4$ , and  $(y \in \mathbb{R}_1^1 \cup \mathbb{R}_1^3 \text{ iff } z \in \mathbb{R}_2^2 \cup \mathbb{R}_2^4)$ ;
4.  $x \in \mathbb{R}_1^4, w \in \mathbb{R}_2^3$ , and  $(y \in \mathbb{R}_1^2 \cup \mathbb{R}_1^4 \text{ iff } z \in \mathbb{R}_2^1 \cup \mathbb{R}_2^3)$ ;
5.  $x \in \mathbb{R}_1^4, w \in \mathbb{R}_2^4$ , and  $(y \in \mathbb{R}_1^2 \cup \mathbb{R}_1^4 \text{ iff } z \in \mathbb{R}_2^2 \cup \mathbb{R}_2^4)$ .

We show that the relation  $Z$  is an  $\overline{\text{LBEABE}}$ -bisimulation. It can be easily checked that every pair  $([x, y], [w, z])$  of  $Z$ -related intervals is such that either  $[x, y] \in V_1(p)$  and  $[w, z] \in V_2(p)$  or  $[x, y] \notin V_1(p)$  and  $[w, z] \notin V_2(p)$ . In order to verify the forward condition, let  $[x, y]$  and  $[w, z]$  be two  $Z$ -related intervals. For each modal operator  $\langle X \rangle$  of the language and each interval  $[x', y']$  such that  $[x, y] R_X [x', y']$ , we have to exhibit an interval  $[w', z']$  such that  $[x', y']$  and  $[w', z']$  are  $Z$ -related, and  $[w, z]$  and  $[w', z']$  are  $R_X$ -related. We proceed case-by-case. Let  $\langle X \rangle = \langle L \rangle$ . We must consider five sub-cases depending on the sets  $x'$  and  $y'$  belong to: (i) if  $x' \in \mathbb{R}_1^1 \cup \mathbb{R}_1^2$ , then for each  $w' \in \mathbb{R}_2^1$  such that  $w' > z$ , we have that, for every  $z' > w'$ ,  $([x', y'], [w', z']) \in Z$  and  $[w, z] R_L [w', z']$  (the existence of  $w'$  is guaranteed by right-unboundedness of  $\mathbb{R}_2^1$ ); (ii) if  $x' \in \mathbb{R}_1^3$  and  $y' \in \mathbb{R}_1^1 \cup \mathbb{R}_1^3$ , then

for each  $w', z'$  such that  $z < w' < z'$  and  $w', z' \in \mathbb{R}_2^3$ , we have that  $([x', y'], [w', z']) \in Z$  and  $[w, z]R_L[w', z']$  (right-unboundedness of  $\mathbb{R}_2^3$ ); (iii) if  $x' \in \mathbb{R}_1^3$  and  $y' \in \mathbb{R}_1^2 \cup \mathbb{R}_1^4$ , then for each  $w', z'$  such that  $z < w' < z'$ ,  $w' \in \mathbb{R}_2^3$ , and  $z' \in \mathbb{R}_2^4$ , we have that  $([x', y'], [w', z']) \in Z$  and  $[w, z]R_L[w', z']$  (right-unboundedness of  $\mathbb{R}_2^3$  and  $\mathbb{R}_2^4$ ); (iv) if  $x' \in \mathbb{R}_1^4$  and  $y' \in \mathbb{R}_1^1 \cup \mathbb{R}_1^3$ , then for each  $w', z'$  such that  $z < w' < z'$ ,  $w' \in \mathbb{R}_2^4$ , and  $z' \in \mathbb{R}_2^3$ , we have that  $([x', y'], [w', z']) \in Z$  and  $[w, z]R_L[w', z']$  (right-unboundedness of  $\mathbb{R}_2^3$  and  $\mathbb{R}_2^4$ ); (v) if  $x' \in \mathbb{R}_1^4$  and  $y' \in \mathbb{R}_1^2 \cup \mathbb{R}_1^4$ , then for each  $w', z'$  such that  $z < w' < z'$  and  $w', z' \in \mathbb{R}_2^4$ , we have that  $([x', y'], [w', z']) \in Z$  and  $[w, z]R_L[w', z']$  (right-unboundedness of  $\mathbb{R}_2^4$ ). Assume now  $\langle X \rangle = \langle B \rangle$ . If  $x \in \mathbb{R}_1^1 \cup \mathbb{R}_1^2$  and  $w \in \mathbb{R}_2^1 \cup \mathbb{R}_2^2$ , then for any  $w < z' < z$ , both  $([x, y'], [w, z']) \in Z$  and  $[w, z]R_B[w, z']$  hold. If  $x \in \mathbb{R}_1^i$  and  $w \in \mathbb{R}_2^j$ , for some  $i \in \{3, 4\}$ , and  $y' \in \mathbb{R}_1^k$ , for some  $k \in \{1, 2, 3, 4\}$ , then for any  $w < z' < z$  such that  $z' \in \mathbb{R}_2^k$ , it holds that  $([x, y'], [w, z']) \in Z$  and  $[w, z]R_B[w, z']$  (the existence of  $z'$  is guaranteed by density of  $\mathbb{R}_2^k$  in  $\mathbb{R}$ ). Finally, if  $x \in \mathbb{R}_1^i$  and  $w \in \mathbb{R}_2^{i'}$  for  $i, i' \in \{3, 4\}$  with  $i \neq i'$ , then if  $y' \in \mathbb{R}_1^1 \cup \mathbb{R}_1^3$  (resp.,  $y' \in \mathbb{R}_1^2 \cup \mathbb{R}_1^4$ ) for any  $w < z' < z$  such that  $z' \in \mathbb{R}_2^2 \cup \mathbb{R}_2^4$  (resp.,  $z' \in \mathbb{R}_2^1 \cup \mathbb{R}_2^3$ ), it holds that  $([x, y'], [w, z']) \in Z$  and  $[w, z]R_B[w, z']$  (density of  $\mathbb{R}_2^2$  and  $\mathbb{R}_2^4$ , resp.,  $\mathbb{R}_2^1$  and  $\mathbb{R}_2^3$ , in  $\mathbb{R}$ ). The remaining cases can be dealt with in a similar way. Let us consider now two intervals  $[x, y]$  and  $[w, z]$  such that  $x \in \mathbb{R}_1^1$ ,  $w \in \mathbb{R}_2^1$ ,  $y \in \mathbb{R}_3^3$ , and  $z \in \mathbb{R}_2^1$ . By definition of  $Z$ ,  $[x, y]$  and  $[w, z]$  are  $Z$ -related, and by definition of  $V_1$  and  $V_2$ , there exists  $y' > y$  such that  $M_1, [y, y'] \Vdash p$ , but there is no  $z' > z$  such that  $M_2, [z, z'] \Vdash p$ . This allows us to conclude that  $Z$  does not preserve the relation induced by  $\langle A \rangle$ , and thus  $\langle A \rangle$  is not definable in  $\text{LBEDO}\overline{\text{ALBEDO}}$ .

A completely symmetric argument can be applied for the completeness proof of  $\langle \overline{A} \rangle$ .  $\square$

#### 4.4 Completeness for $\langle D \rangle$ , $\langle \overline{D} \rangle$ , $\langle O \rangle$ , and $\langle \overline{O} \rangle$

To deal with the modalities  $\langle D \rangle$ ,  $\langle \overline{D} \rangle$ ,  $\langle O \rangle$ , and  $\langle \overline{O} \rangle$ , we proceed as follows. We first introduce the notion of  $f$ -model, that is, for any given function  $f : \mathbb{R} \rightarrow \mathbb{Q}$ , we define a model  $M_f$ , called  $f$ -model, whose valuation is based on  $f$ . Then, for any given pair of functions  $f_1$  and  $f_2$ , we define a suitable relation  $Z_{f_1}^{f_2}$  between the models  $M_{f_1}$  and  $M_{f_2}$  (from now on, we will simply write  $Z$  when there is no ambiguity about the involved models). Finally, we specify the requirements that  $f_1$  and  $f_2$  must satisfy to make  $Z$  the bisimulation we want (these requirements vary from one modality to the other).

**Lemma 4.4.** *The set of inter-definability equations for  $\langle D \rangle$ ,  $\langle \overline{D} \rangle$ ,  $\langle O \rangle$ , and  $\langle \overline{O} \rangle$  given in Table 2 is complete.*

*Proof.* We will detail the case of the modality  $\langle D \rangle$ . The other cases can be proved using similar arguments.

According to Table 2,  $\langle D \rangle$  is definable in terms of BE. The fragments  $\text{ALBOALBEDO}$  and  $\text{ALEOALBEDO}$  are thus the only  $\subseteq$ -maximal ones not featuring  $\langle D \rangle$  and not containing BE. We should provide a bisimulation, not preserving the relation induced by  $\langle D \rangle$ , for each of these fragments, but, thanks to the symmetry of the operators, it suffices to consider only one of them, say  $\text{ALBOALBEDO}$  (by Lemma 3.2,

we have that  $\text{ALBOALBEDO}$  is expressively equivalent to  $\text{ABOABE}$ ). Given a function  $f : \mathbb{R} \rightarrow \mathbb{Q}$ , we define the  $f$ -model  $M_f$ , over a language with one propositional letter  $p$  only, as the pair  $\langle \mathbb{I}(\mathbb{R}), V_f \rangle$ , where  $V_f : \mathbb{I}(\mathbb{R}) \rightarrow 2^{\mathcal{A}^P}$  is defined as follows:  $[x, y] \in V_f(p) \stackrel{\text{def}}{\iff} y \geq f(x)$ . For any given pair of functions  $f_1$  and  $f_2$  (from  $\mathbb{R}$  to  $\mathbb{Q}$ ), the relation  $Z$  is defined as follows:

$$([x, y], [w, z]) \in Z \stackrel{\text{def}}{\iff} x \equiv w, y \equiv z, \text{ and } [x, y] \equiv_l [w, z],$$

where  $u \equiv v \stackrel{\text{def}}{\iff} u \in \mathbb{Q}$  iff  $v \in \mathbb{Q}$  and  $[u, u'] \equiv_l [v, v'] \stackrel{\text{def}}{\iff} u' \sim f_1(u)$  and  $v' \sim f_2(v)$ , for  $\sim \in \{<, =, >\}$ . Finally, the following constraints are imposed on  $f$  (if we replace  $\langle D \rangle$  by one of the other modalities, the constraints must be suitably replaced as well): (i) for every  $x \in \mathbb{R}$ ,  $f(x) > x$ , (ii) for every  $x \in \mathbb{Q}$ , both  $f^{-1}(x) \cap \mathbb{Q}$  and  $f^{-1}(x) \cap \overline{\mathbb{Q}}$  are left-unbounded (notice that surjectivity of  $f$  immediately follows), and (iii) for every  $x, y \in \mathbb{R}$ , if  $x < y$ , then there exists  $u_1 \in \mathbb{Q}$  (resp.,  $u_2 \in \overline{\mathbb{Q}}$ ) such that  $x < u_1 < y$  (resp.,  $x < u_2 < y$ ) and  $y < f(u_1)$  (resp.,  $y < f(u_2)$ ).

Now, we show that if both  $f_1$  and  $f_2$  satisfy the above conditions, then  $Z$  is an  $\text{ABOABE}$ -bisimulation between  $M_{f_1}$  and  $M_{f_2}$ . Let  $[x, y]$  and  $[w, z]$  be two  $Z$ -related intervals. By definition,  $y \sim f_1(x)$  and  $z \sim f_2(w)$  for some  $\sim \in \{<, =, >\}$ . If  $\sim \in \{=, >\}$ , then both  $[x, y]$  and  $[w, z]$  satisfy  $p$ ; otherwise, both of them satisfy  $\neg p$ . The local condition is thus satisfied. As for the forward condition, let  $[x, y]$  and  $[x', y']$  be two intervals in  $M_{f_1}$  and  $[w, z]$  an interval in  $M_{f_2}$ . We have to prove that if  $[x, y]$  and  $[w, z]$  are  $Z$ -related, then, for each modal operator  $\langle X \rangle$  of  $\text{ABOABE}$  such that  $[x, y]R_X[x', y']$ , there exists an interval  $[w', z']$  such that  $[x', y']$  and  $[w', z']$  are  $Z$ -related and  $[w, z]R_X[w', z']$ . Once more, we proceed case-by-case. For the sake of brevity, we only detail the case of  $\langle A \rangle$ . The other modalities can also be dealt with by exploiting the requirements for the functions  $f_1$  and  $f_2$  in a suitable way. Let  $\langle X \rangle = \langle A \rangle$ . By definition of  $\langle A \rangle$ ,  $x' = y$  and we are forced to choose  $w' = z$ . By  $y \equiv z$ , it immediately follows  $x' \equiv w'$ . We must find a point  $z' > z$  such that  $y' \equiv z'$  and both  $y' \sim f_1(y)$  and  $z' \sim f_2(z)$  for some  $\sim \in \{<, =, >\}$ . Let us suppose that  $y' < f_1(y)$ . In such a case, we choose a point  $z'$  such that  $z < z' < f_2(z)$  and  $y' \equiv z'$ . The existence of such a point is guaranteed by condition (i) on  $f_2$  and by the density of  $\mathbb{Q}$  and  $\overline{\mathbb{Q}}$  in  $\mathbb{R}$ . Otherwise, if  $y' = f_1(y)$ , we choose  $z' = f_2(z)$ . By definition of  $f_1$  and  $f_2$  (the codomain of  $f_1$  and  $f_2$  is  $\mathbb{Q}$ ), both  $y'$  and  $z'$  belong to  $\mathbb{Q}$  and thus  $y' \equiv z'$ . Finally, if  $y' > f_1(y)$ , we choose  $z' > f_2(z)$  such that  $y' \equiv z'$ . The existence of such a point is guaranteed by right-unboundedness of  $\mathbb{Q}$  and  $\overline{\mathbb{Q}}$ . Satisfaction of the backward condition for all modalities can be checked in a similar way.

To complete the proof, we exhibit two functions that meet the requirements we have imposed to  $f_1$  and  $f_2$ , but do not preserve the relation induced by  $\langle D \rangle$ . Let  $\mathcal{P}(\mathbb{Q}) = \{\mathbb{Q}_q \mid q \in \mathbb{Q}\}$  and  $\mathcal{P}(\overline{\mathbb{Q}}) = \{\overline{\mathbb{Q}}_q \mid q \in \mathbb{Q}\}$  be infinite and countable partitions of  $\mathbb{Q}$  and  $\overline{\mathbb{Q}}$ , respectively, such that for every  $q \in \mathbb{Q}$ , both  $\mathbb{Q}_q$  and  $\overline{\mathbb{Q}}_q$  are dense in  $\mathbb{R}$ . For every  $q \in \mathbb{Q}$ , let  $\mathbb{R}_q = \mathbb{Q}_q \cup \overline{\mathbb{Q}}_q$ . We define a function  $g : \mathbb{R} \rightarrow \mathbb{Q}$  that maps every

real number  $x$  to the index  $q$  (a rational number) of the class  $\mathbb{R}_q$  it belongs to. Formally, for every  $x \in \mathbb{R}$ ,  $g(x) = q$ , where  $q \in \mathbb{Q}$  is the unique rational number such that  $x \in \mathbb{R}_q$ . The two functions  $f_1 : \mathbb{R} \rightarrow \mathbb{Q}$  and  $f_2 : \mathbb{R} \rightarrow \mathbb{Q}$  are defined as follows:

$$f_1(x) = \begin{cases} g(x) & \text{if } x < g(x), x \neq 1, \text{ and } x \neq 0 \\ 2 & \text{if } x = 1 \\ \lceil x + 3 \rceil & \text{otherwise} \end{cases}$$

$$f_2(x) = \begin{cases} g(x) & \text{if } x < g(x) \text{ and } x \notin [0, 3] \\ \lceil x + 3 \rceil & \text{otherwise} \end{cases}$$

It is not difficult to check that the above-defined functions meet the requirements for  $f_1$  and  $f_2$ , and thus  $Z$  is an  $\text{ABO}\overline{\text{ABE}}$ -bisimulation. On the other hand,  $Z$  does not preserve the relation induced by  $\langle D \rangle$ . Consider the interval  $[0, 3]$  in  $M_{f_1}$  and the interval  $[0, 3]$  in  $M_{f_2}$ . It is immediate to see that these two intervals are  $Z$ -related. However,  $M_{f_1}, [0, 3] \Vdash \langle D \rangle p$  (as  $M_{f_1}, [1, 2] \Vdash p$ ), but  $M_{f_2}, [0, 3] \Vdash \neg \langle D \rangle p$ . This allows us to conclude that  $\langle D \rangle$  is not definable in the fragment  $\text{ALBOALBEDO}$ .  $\square$

#### 4.5 Harvest

The proof of Theorem 3.1 follows now immediately.

We have used the equations in Table 2 as the basis of a simple program that identifies and counts all expressively different fragments of HS with respect to the strict semantics. Using that program, we have found that, under our assumptions (strict semantics, over the class of all linear orders) there are exactly 1347 genuine, that is, expressively different, fragments out of  $2^{12} = 4096$  different subsets of HS-operators.

## 5 Conclusions

In this paper, we have obtained a sound, complete, and optimal set of inter-definability equations among all modal operators in HS, thus providing a characterization of the relative expressive power of all interval logics definable as fragments of HS. Such a classification has a number of important applications. As an example, it allows one to properly identify the (small) set of HS fragments for which the decidability of the satisfiability problem is still an open problem.

It should be emphasized that the set of inter-definability equations listed in Table 2 and the resulting classification do not apply if the non-strict semantics is considered. For instance, if the non-strict semantics is assumed, then, as shown in [Venema, 1990],  $\langle A \rangle$  (resp.,  $\langle \overline{A} \rangle$ ) can be defined in  $\overline{\text{BE}}$  (resp.,  $\text{B}\overline{\text{E}}$ ). Also, if the semantics is restricted to specific classes of linear orders, the completeness of the set of equations in Table 2 is no longer guaranteed. For instance, in discrete linear orders,  $\langle A \rangle$  can be defined in  $\overline{\text{BE}}$ :  $\langle A \rangle p \equiv \varphi(p) \vee \langle E \rangle \varphi(p)$ , where  $\varphi(p)$  is a shorthand for  $[E] \perp \wedge \langle \overline{B} \rangle ([E][E] \perp \wedge \langle E \rangle (p \vee \langle \overline{B} \rangle p))$ ; likewise,  $\langle \overline{A} \rangle$  is definable in  $\text{B}\overline{\text{E}}$ . As another example, in dense linear orders,  $\langle L \rangle$  can be defined in  $\text{DO}$ :  $\langle L \rangle p \equiv \langle O \rangle (\langle O \rangle \top \wedge [O] (\langle O \rangle p \vee \langle D \rangle p \vee \langle D \rangle \langle O \rangle p))$ ; likewise,  $\langle \overline{L} \rangle$  is definable in  $\text{D}\overline{\text{O}}$ . (In view of these inter-definabilities, Lemma 4.1 cannot be proved by using bisimulation between models over the reals.)

The classification of the expressiveness of HS fragments with respect to the non-strict semantics, as well as over specific classes of linear orders, is currently under investigation and will be reported in a forthcoming publication.

## References

- [Allen, 1983] J. F. Allen. Maintaining knowledge about temporal intervals. *Communications of the ACM*, 26(11):832–843, 1983.
- [Blackburn *et al.*, 2002] P. Blackburn, M. de Rijke, and Y. Venema. *Modal Logic*. CUP, 2002.
- [Bresolin *et al.*, 2007a] D. Bresolin, A. Montanari, and P. Sala. An optimal tableau-based decision algorithm for Propositional Neighborhood Logic. In *Proc. of STACS 2007*, volume 4393 of *LNCS*, pages 549–560. Springer, 2007.
- [Bresolin *et al.*, 2007b] D. Bresolin, A. Montanari, and G. Sciavicco. An optimal decision procedure for Right Propositional Neighborhood Logic. *Journal of Automated Reasoning*, 38(1-3):173–199, 2007.
- [Bresolin *et al.*, 2008] D. Bresolin, A. Montanari, P. Sala, and G. Sciavicco. Optimal Tableaux for Right Propositional Neighborhood Logic over Linear Orders. In *Proc. of the 11th European Conference on Logics in AI*, volume 5293 of *LNAI*, pages 62–75. Springer, 2008.
- [Bresolin *et al.*, 2009] D. Bresolin, V. Goranko, A. Montanari, and G. Sciavicco. Propositional interval neighborhood logics: Expressiveness, decidability, and undecidable extensions. *Annals of Pure and Applied Logic*, 161(3):289–304, 2009.
- [Bresolin *et al.*, 2010] D. Bresolin, V. Goranko, A. Montanari, and P. Sala. Tableaux for logics of subinterval structures over dense orderings. *Journal of Logic and Computation*, 20:133–166, 2010.
- [Halpern and Shoham, 1991] J. Halpern and Y. Shoham. A propositional modal logic of time intervals. *Journal of the ACM*, 38(4):935–962, 1991.
- [Montanari *et al.*, 2010a] A. Montanari, G. Puppis, and P. Sala. Maximal decidable fragments of Halpern and Shoham’s modal logic of intervals. In *Proc. of ICALP’2010*, volume 6199 of *LNCS*, pages 345–356. Springer, 2010.
- [Montanari *et al.*, 2010b] A. Montanari, G. Puppis, P. Sala, and G. Sciavicco. Decidability of the interval temporal logic  $\text{AB}\overline{\text{B}}$  on natural numbers. In *Proc. of STACS 2010*, pages 597–608, 2010.
- [Moszkowski, 1983] B. Moszkowski. *Reasoning about digital circuits*. Tech. rep. stan-cs-83-970, Dept. of Computer Science, Stanford University, Stanford, CA, 1983.
- [Venema, 1990] Y. Venema. Expressiveness and completeness of an interval tense logic. *Notre Dame Journal of Formal Logic*, 31(4):529–547, 1990.
- [Zhou and Hansen, 2004] C. Zhou and M. R. Hansen. *Duration Calculus: A Formal Approach to Real-Time Systems*. Springer, 2004.