



Multiplicative properties of integer valued polynomials over split-quaternions

Antonio Cigliola, Dario Spirito & Francesca Tartarone

To cite this article: Antonio Cigliola, Dario Spirito & Francesca Tartarone (2021) Multiplicative properties of integer valued polynomials over split-quaternions, Communications in Algebra, 49:3, 1338-1351, DOI: [10.1080/00927872.2020.1834572](https://doi.org/10.1080/00927872.2020.1834572)

To link to this article: <https://doi.org/10.1080/00927872.2020.1834572>



Published online: 28 Oct 2020.



Submit your article to this journal [↗](#)



Article views: 52



View related articles [↗](#)



View Crossmark data [↗](#)



Multiplicative properties of integer valued polynomials over split-quaternions

Antonio Cigliola^a, Dario Spirito^b, and Francesca Tartarone^c

^aDipartimento di Scienze Statistiche, Sapienza Università di Roma, Istituto Comprensivo “G. Bagnera”, Roma, Italy; ^bDipartimento di Matematica “Tullio Levi Civita”, Università degli Studi di Padova, Padova, Italy; ^cDipartimento di Matematica e Fisica, Università degli Studi Roma Tre, Roma, Italy

ABSTRACT

We study localization properties and the prime spectrum of the integer-valued polynomial ring $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$, where $\mathbb{P}_{\mathbb{Z}}$ (respectively $\mathbb{P}_{\mathbb{Q}}$) is the algebra of split-quaternion over \mathbb{Z} (respectively over \mathbb{Q}).

ARTICLE HISTORY

Received 2 July 2020
Revised 18 September 2020
Communicated by Alfred Geroldinger

KEYWORDS

Krull dimension; prime ideal; quaternion algebra

2010 MATHEMATICS

SUBJECT

CLASSIFICATION

Primary: 11R52;
16H05; 13F20

Introduction

In [14] N. Werner studied the ring of integer-valued polynomials in a noncommutative setting, by considering quaternion algebras. Precisely, he considered the algebras $\mathbb{H}_{\mathbb{Z}}$ and $\mathbb{H}_{\mathbb{Q}}$ (respectively over \mathbb{Z} and over \mathbb{Q}) generated by the unit elements $1, \mathbf{i}, \mathbf{j}, \mathbf{k}$, linked by the relations $\mathbf{i}^2 = \mathbf{j}^2 = \mathbf{k}^2 = -1$, $\mathbf{i}\mathbf{j} = \mathbf{k} = -\mathbf{j}\mathbf{i}$, $\mathbf{j}\mathbf{k} = \mathbf{i} = -\mathbf{k}\mathbf{j}$ and $\mathbf{k}\mathbf{i} = \mathbf{j} = -\mathbf{i}\mathbf{k}$, and considered the set $\text{Int}_{\mathbb{H}_{\mathbb{Q}}}(\mathbb{H}_{\mathbb{Z}})$ of all polynomials $f \in \mathbb{H}_{\mathbb{Q}}[x]$ such that $f(\mathbb{H}_{\mathbb{Z}}) \subseteq \mathbb{H}_{\mathbb{Z}}$. After proving that $\text{Int}_{\mathbb{H}_{\mathbb{Q}}}(\mathbb{H}_{\mathbb{Z}})$ is indeed a noncommutative ring (which strictly contains $\mathbb{H}_{\mathbb{Q}}[x]$), he investigated the ideal structure of this ring, describing some prime ideals above the zero and the maximal ideals of $\mathbb{H}_{\mathbb{Z}}$.

Moving from these ideas, in [3] A. Cigliola, K.A. Loper and N. Werner focused on similar problems in a different setting: instead of $\mathbb{H}_{\mathbb{Z}}$ they considered the set of integer split-quaternions $\mathbb{P}_{\mathbb{Z}}$, i.e. the \mathbb{Z} -algebra generated by the unit elements $1, \mathbf{i}, \mathbf{j}, \mathbf{k}$ with the relations $-\mathbf{i}^2 = \mathbf{j}^2 = \mathbf{k}^2 = 1$ and $\mathbf{i}\mathbf{j}\mathbf{k} = 1$ (see Definition 1.1).

In this paper, we continue the study of the ring $\mathbb{P}_{\mathbb{Z}}$ (Section 1) and of the ring $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$ of integer-valued polynomials over $\mathbb{P}_{\mathbb{Z}}$ (Section 2). We study some denominator sets of $\mathbb{P}_{\mathbb{Z}}$ and $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$ that are not subsets of \mathbb{Z} (in particular, they are not central) and their ring of fractions. Thus, we partially answer to one of the open questions posed in [3, §5] which asks whether it is possible to find and to localize $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$ with respect to noncentral sets. We then study the ring $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ of the polynomials in $\mathbb{Q}[x]$ that are integer valued over $\mathbb{P}_{\mathbb{Z}}$. There is a strict connection between the prime spectrum of this ring and the prime spectrum of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$. This allows to

calculate the Krull dimension of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}(p)})$, for an odd prime integer p , starting from the dimension of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ and thus to get a partial but interesting information about the Krull dimension of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$. Finally, in [Section 4](#), we study in more detail the ideal $p\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ and show that it is not prime. In this last Section we will be able to construct explicitly some polynomials of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$.

Throughout the paper, all the rings we consider are unitary but not necessarily commutative.

1. Localizations of $\mathbb{P}_{\mathbb{Z}}$

We recall some definitions and basic properties.

Definition 1.1. Let R be a commutative ring. We denote by \mathbb{P}_R the R -algebra generated by the four unit elements $1, \mathbf{i}, \mathbf{j}$ and \mathbf{k} with the relations

$$- \mathbf{i}^2 = \mathbf{j}^2 = \mathbf{k}^2 = \mathbf{i} \mathbf{j} \mathbf{k} = 1.$$

Formally $\mathbb{P}_R := \{q = a + b \mathbf{i} + c \mathbf{j} + d \mathbf{k} \mid a, b, c, d \in R\}$.

We call \mathbb{P}_R the *ring of split-quaternions* over R .

Let $q = a + b \mathbf{i} + c \mathbf{j} + d \mathbf{k} \in \mathbb{P}_R$, then:

- (a) a, b, c , and d are the *coefficients* of q , and a is the *real part* of q ;
- (b) the *bar conjugate* of q is $\bar{q} := a - b \mathbf{i} - c \mathbf{j} - d \mathbf{k}$;
- (c) the *norm* of q is $N(q) := q\bar{q} = a^2 + b^2 - c^2 - d^2$;
- (d) the *trace* of q is $T(q) = q + \bar{q} = 2a$;
- (e) the *minimal polynomial* of q is ([3, Definition 2.4])

$$m_q(x) := \begin{cases} x - q & \text{if } q \in R \\ x^2 - T(q)x + N(q) & \text{if } q \in \mathbb{P}_R \setminus R. \end{cases}$$

where $m_q(x)$ is minimal in the way that $m_q(q) = 0$ and that $m_q(x)$ is the monic polynomial of least degree having q as a root.

In this section, we study some localizations of $\mathbb{P}_{\mathbb{Z}}$. We start with the description of its prime and maximal ideals. Recall that an ideal P of a (not necessarily commutative) ring R is *prime* if, given $a, b \in R$ such that $aPb \subseteq P$, then $a \in P$ or $b \in P$.

Theorem 1.2. [3, Theorem 2.11]. *The prime ideals of $\mathbb{P}_{\mathbb{Z}}$ are:*

- (i) (0) ;
- (ii) $p\mathbb{P}_{\mathbb{Z}}$ where p is an odd prime of \mathbb{Z} ;
- (iii) $\mathcal{M} = (1 + \mathbf{i}; 1 + \mathbf{j})$.

Moreover, the primes $p\mathbb{P}_{\mathbb{Z}}$ and \mathcal{M} are maximal, and \mathcal{M} is the only prime ideal containing 2.

Lemma 1.3. *Let $q \in \mathbb{P}_{\mathbb{Z}}$ such that $2 \mid N(q)$. Then $q \in \mathcal{M}$. In particular \mathcal{M} contains all the zero-divisors of $\mathbb{P}_{\mathbb{Z}}$.*

Proof. Let $q = a + b \mathbf{i} + c \mathbf{j} + d \mathbf{k}$ be such that $N(q) = a^2 + b^2 - c^2 - d^2 = 2m$, for some $m \in \mathbb{Z}$. By hypothesis, q must have zero, two or four even coefficients. In the case that all coefficients are even, then trivially $q \in (2) \subseteq \mathcal{M}$. If q has exactly two even coefficients, then q is congruent modulo $2\mathbb{P}_{\mathbb{Z}}$ to the sum of two of $1, \mathbf{i}, \mathbf{j}$ and \mathbf{k} , and all of them are elements of \mathcal{M} . Finally, if all coefficients of q are odd, then $q \equiv 1 + \mathbf{i} + \mathbf{j} + \mathbf{k} \pmod{2\mathbb{P}_{\mathbb{Z}}}$, and so $q \in \mathcal{M}$ since $1 + \mathbf{i} + \mathbf{j} + \mathbf{k} = (1 + \mathbf{i})(1 + \mathbf{j}) \in \mathcal{M}$. \square

Definition 1.4. Let R be a ring and S a multiplicative subset in R . We say that S is a *right denominator set* if:

- (i) for any $a \in R$ and $s \in S$, $aS \cap sR \neq \emptyset$ (this condition is known as *right Ore condition* and S is called a *right Ore set*);
- (ii) for $a \in R$, if $s'a = 0$ for some $s' \in S$, then $as = 0$ for some $s \in S$ (we say that S is *right reversible*).

Remark 1.5.

- (a) We can define left denominator sets in a completely symmetrical way.
- (b) Condition (ii) (reversibility) is automatically satisfied when S does not contain zero-divisors.
- (c) It is easily seen that the multiplicative subsets contained in the center of R are always denominator subsets.

By Lam [9, Theorem 10.6], if R is a ring and S a multiplicative subset of R , then R has a right ring of fractions with respect to S (namely, the ring $RS^{-1} := \{as^{-1} \mid a \in R, s \in S\}$) if and only if S is a right denominator set. Similarly we can construct the ring $S^{-1}R := \{s^{-1}a \mid a \in R, s \in S\}$ if and only if S is a left denominator set. If S is both a right and left denominator set, then $RS^{-1} \simeq S^{-1}R$ by Lam [9, Corollary 10.14].

Lemma 1.6. Let R be a commutative ring and S a multiplicative subset of \mathbb{P}_R , closed under norm (i.e., if $s \in S$ then $N(s) \in S$). Then S verifies both the right and the left Ore condition.

Proof. Fix $a \in \mathbb{P}_R$ and $s \in S$. Since $N(s) \in R$ is a central element, we have that $aN(s) = N(s)a$. It follows that $aN(s) = s(\bar{s}a)$, so S is a right Ore set since $aS \cap s\mathbb{P}_R \neq \emptyset$. Analogously, $(a\bar{s})s = N(s)a$ so S is a left Ore set since $Sa \cap \mathbb{P}_{RS} \neq \emptyset$. □

By the previous lemma, if $S = \mathcal{R}(R)$ is the set of all (right and left) regular elements of R , then S is a denominator set and RS^{-1} is the *total ring of fractions of R* , which we denote by $\mathcal{Q}(R)$.

For commutative rings, the most important way of constructing localizations of a ring R is through the sets $R \setminus P$, where P is a prime ideal; however, if R is not commutative, the complement of a prime ideal may not be multiplicatively closed. For example, if $p = 2k + 1$ is an odd prime number, then $p\mathbb{P}_{\mathbb{Z}}$ is prime, but $\mathbb{P}_{\mathbb{Z}} \setminus p\mathbb{P}_{\mathbb{Z}}$ is not multiplicatively closed since $((k + 1) + k \mathbf{j})((k + 1) - k \mathbf{j}) = p \in p\mathbb{P}_{\mathbb{Z}}$.

Following the notation of Goldie [6], we give the following definition:

Definition 1.7. Let be given a ring R and let Q be a proper prime ideal of R . We set:

$$\mathcal{C}(Q) := \{x \in R \mid xr \notin Q, \forall r \notin Q\},$$

and

$$\mathcal{C}'(Q) := \{x \in R \mid rx \notin Q, \forall r \notin Q\}.$$

Proposition 1.8. Let R be a ring and let $Q \subsetneq R$ be a prime ideal of R . Then $\mathcal{C}(Q)$ is a multiplicatively closed subset of R containing 1 but not 0, and $\mathcal{C}(Q) \subseteq R \setminus Q$. The same properties hold for $\mathcal{C}'(Q)$.

Proof. For each $r \notin Q$, we have that $1 \cdot r = r \notin Q$ and that $0 \cdot r \in Q$. Then, by definition, $1 \in \mathcal{C}(Q)$ and $0 \notin \mathcal{C}(Q)$. Take now $a, b \in \mathcal{C}(Q)$ and $r \notin Q$. Since $b \in \mathcal{C}(Q)$, then $br \notin Q$. Again, since $a \in \mathcal{C}(Q)$, we have $a(br) \notin Q$. Thus for all $r \notin Q$ we have $(ab)r = a(br) \notin Q$.

Finally, if $x \in \mathcal{C}(Q)$ then, since $1 \notin Q$, we have $x \cdot 1 = x \notin Q$. Hence, $\mathcal{C}(Q) \subseteq R \setminus Q$. □

Proposition 1.9. Let R be a ring and let $Q \subsetneq R$ be a prime ideal of R . Then $\mathcal{C}(Q)$ is the set of left regular elements of R modulo Q and $\mathcal{C}'(Q)$ is the set of right regular elements of R modulo Q .

Proof. Take $x \in R$. Then x is a left zero-divisor modulo Q if and only if there is $r \in R/Q, r \neq 0$, such that $xr = 0$. This is equivalent to saying that there is an $r \notin Q$ such that $xr \in Q$. In other words, $x \notin \mathcal{C}(Q)$. Similarly for $\mathcal{C}'(Q)$. \square

In particular, we have that $\mathcal{C}(0) = \mathcal{R}_l(R)$ is the set of the left regular elements of R , while $\mathcal{C}'(0) = \mathcal{R}_r(R)$ is the set of the right regular elements of R .

We now focus on some properties of the sets $\mathcal{C}(Q)$ associated to the prime ideals of $\mathbb{P}_{\mathbb{Z}}$.

Proposition 1.10. Let Q be a prime ideal of $\mathbb{P}_{\mathbb{Z}}$. Then:

- (i) $\mathcal{C}(Q)$ is closed under bar conjugation;
- (ii) $\mathcal{C}(Q)$ is closed under norm;
- (iii) $\mathcal{C}(Q) = \{x \in \mathbb{P}_{\mathbb{Z}} \mid N(x) \notin Q\}$;
- (iv) $\mathcal{C}(Q)$ does not contain any zero-divisor.

Proof. By Goodearl and Warfield [7, Proposition 1.6] $\mathbb{P}_{\mathbb{Z}}$ is a Noetherian ring. Thus, from [6, Section 3], $\mathcal{C}(Q) = \mathcal{C}'(Q)$.

Consider first $Q = (0)$. Then $\mathcal{C}(0)$ equals $\mathcal{R}(\mathbb{P}_{\mathbb{Z}})$, the set of all (two-sided) regular elements, and so

$$\mathcal{C}(Q) = \mathcal{R}(\mathbb{P}_{\mathbb{Z}}) = \{x \in \mathbb{P}_{\mathbb{Z}} \mid N(x) \neq 0\}.$$

This proves the claim in the case $Q = (0)$.

Let now be $Q = p\mathbb{P}_{\mathbb{Z}}$, for an odd prime integer p . We notice that:

- $\mathcal{C}(Q)(\text{mod } Q) = \mathcal{C}(\bar{0})$ in $\mathbb{P}_{\mathbb{Z}}/Q = \mathbb{P}_{\mathbb{Z}_p}$ (apply Proposition 1.9);
- $N(x)(\text{mod } p) = N(\bar{x})$, for $x \in \mathbb{P}_{\mathbb{Z}}$ and $\bar{x} = x(\text{mod } p\mathbb{P}_{\mathbb{Z}})$.

Using these equalities, points (i)-(ii)-(iii) reduce to the case $Q = (0)$, which has been already proved. For $p = 2$, the same reasoning applies reducing modulo \mathcal{M} .

For the point (iv), if $p = 2$ the claim follows from Lemma 1.3.

If p is an odd prime, then suppose that $xr' = 0$, for some $x \in \mathcal{C}(Q)$ and $0 \neq r' \in \mathbb{P}_{\mathbb{Z}}$. If we write $r' = p^m r$, for some $r \notin Q$, we get $xr = 0 \in Q$ (since p is not a zero divisor from Lemma 1.3) which is absurd. \square

In particular, we observe that $\mathcal{C}(p\mathbb{P}_{\mathbb{Z}}) = \{x \in \mathbb{P}_{\mathbb{Z}} \mid p \nmid N(x)\}$ and $\mathcal{C}(\mathcal{M}) = \{x \in \mathbb{P}_{\mathbb{Z}} \mid 2 \nmid N(x)\}$.

We will work with the following multiplicative subsets of $\mathbb{P}_{\mathbb{Z}}$:

- the multiplicative subsets of \mathbb{Z} ;
- the sets $\mathcal{C}(0)$, $\mathcal{C}(\mathcal{M})$ and $\mathcal{C}(p\mathbb{P}_{\mathbb{Z}})$, for any odd prime integer p .

For a general noncommutative ring, given a prime ideal Q , $\mathcal{C}(Q)$ may not be a denominator set: such an example is given, for instance, in [1, Example 2.3]. However we show that $\mathcal{C}(Q)$ is a denominator sets in $\mathbb{P}_{\mathbb{Z}}$ and also in $\text{Int}(\mathbb{P}_{\mathbb{Z}})$ (Proposition 2.4), for each prime ideal Q of $\mathbb{P}_{\mathbb{Z}}$.

Proposition 1.11. The sets $\mathbb{Z} \setminus (0), \mathbb{Z} \setminus p\mathbb{Z}$, for p prime, and $\mathcal{C}(Q)$, for Q prime ideal of $\mathbb{P}_{\mathbb{Z}}$, are (right and left) denominator sets of $\mathbb{P}_{\mathbb{Z}}$.

Proof. Let $S = \mathbb{Z} \setminus (0)$ or $S = \mathbb{Z} \setminus p\mathbb{Z}$, for a prime p . Then the statement easily follows from the fact that S is contained in the center of $\mathbb{P}_{\mathbb{Z}}$.

If $S = \mathcal{C}(Q)$, then S does not contain zero-divisors (Proposition 1.10), so $\mathcal{C}(Q)$ is right and left reversible. Finally, $\mathcal{C}(Q)$ is a right (left) Ore set by Lemma 1.6, since it is closed under bar conjugation (Proposition 1.10). Thus $\mathcal{C}(Q)$ is a right and left denominator set of $\mathbb{P}_{\mathbb{Z}}$. \square

Proposition 1.12. *Let $S = \mathcal{C}(0)$ or $S = \mathbb{Z} \setminus (0)$. Then*

$$\mathbb{P}_{\mathbb{Z}}S^{-1} = S^{-1}\mathbb{P}_{\mathbb{Z}} = \mathbb{P}_{\mathbb{Q}} = \mathcal{Q}(\mathbb{P}_{\mathbb{Z}}),$$

which is the total ring of fractions of $\mathbb{P}_{\mathbb{Z}}$.

Proof. By Proposition 1.11, S is a denominator set. So the ring $\mathbb{P}_{\mathbb{Z}}S^{-1}$ exists and its elements are the fractions rs^{-1} , where $r, s \in \mathbb{P}_{\mathbb{Z}}$ and $N(s) \neq 0$. Then $rs^{-1} = \frac{1}{N(s)}r\bar{s} \in \mathbb{P}_{\mathbb{Q}}$. Thus $\mathbb{P}_{\mathbb{Z}}S^{-1} \subseteq \mathbb{P}_{\mathbb{Q}}$. Conversely, given $q \in \mathbb{P}_{\mathbb{Q}}$, write q in the form $p \cdot a^{-1}$, where $p \in \mathbb{P}_{\mathbb{Z}}$ and a is a common denominator for the coefficients of q . Obviously, $a \in S$ and so $pa^{-1} \in \mathbb{P}_{\mathbb{Z}}S^{-1}$, i.e., $\mathbb{P}_{\mathbb{Z}}S^{-1} \supseteq \mathbb{P}_{\mathbb{Q}}$. Thus $\mathbb{P}_{\mathbb{Z}}S^{-1} = \mathbb{P}_{\mathbb{Q}}$. Similarly, $S^{-1}\mathbb{P}_{\mathbb{Z}} = \mathbb{P}_{\mathbb{Q}}$. Finally $\mathbb{P}_{\mathbb{Q}}$ is the total ring of fractions of $\mathbb{P}_{\mathbb{Z}}$ because we localize with respect to the set of regular elements of $\mathbb{P}_{\mathbb{Z}}$. \square

Similarly, if we localize $\mathbb{P}_{\mathbb{Z}}$ at $S = \mathbb{Z} \setminus p\mathbb{Z}$ or $S = \mathcal{C}(Q)$, where $Q = p\mathbb{P}_{\mathbb{Z}}$, for a prime number p , we get the algebra of split-quaternions with coefficients in $\mathbb{Z}_{(p)}$, the localization of \mathbb{Z} at the ideal $p\mathbb{Z}$ (as we see in the following Proposition). In the following, \mathbb{Z}_p will denote the field with p elements.

Proposition 1.13. *Let p be a prime number and let $S = \mathbb{Z} \setminus p\mathbb{Z}$ or $S = \mathcal{C}(Q)$, where Q is a prime ideal of $\mathbb{P}_{\mathbb{Z}}$ such that $Q \cap \mathbb{Z} = p\mathbb{Z}$.*

Then

$$\mathbb{P}_{\mathbb{Z}}S^{-1} = S^{-1}\mathbb{P}_{\mathbb{Z}} = \mathbb{P}_{\mathbb{Z}_{(p)}}.$$

Proof. We know that S is a denominator set of $\mathbb{P}_{\mathbb{Z}}$ by Proposition 1.11. So the ring $\mathbb{P}_{\mathbb{Z}}S^{-1}$ exists.

Let $S = \mathbb{Z} \setminus p\mathbb{Z}$. It is easy to see that $\mathbb{P}_{\mathbb{Z}}S^{-1} \subseteq \mathbb{P}_{\mathbb{Z}_{(p)}}$. For the reverse inclusion, notice that the minimum common denominator of any element of $\mathbb{Z}_{(p)}$ is an element of $\mathbb{Z} \setminus p\mathbb{Z}$. So $\mathbb{P}_{\mathbb{Z}}S^{-1} = \mathbb{P}_{\mathbb{Z}_{(p)}}$. Similarly it can be proved that $S^{-1}\mathbb{P}_{\mathbb{Z}} = \mathbb{P}_{\mathbb{Z}_{(p)}}$.

Let $S = \mathcal{C}(Q)$. Since the norm of the elements of S is not divisible by p (Proposition 1.10), a right fraction $ps^{-1} \in \mathbb{P}_{\mathbb{Z}}S^{-1}$, for some $p \in \mathbb{P}_{\mathbb{Z}}$ and $s \in S$, can be seen as a rational split-quaternion $q = \frac{1}{N(s)}p\bar{s} = a + b\mathbf{i} + c\mathbf{j} + d\mathbf{k}$, where $a, b, c, d \in \mathbb{Q}$ and their denominators are not divisible by p . Thus $\mathbb{P}_{\mathbb{Z}}S^{-1} \subseteq \mathbb{P}_{\mathbb{Z}_{(p)}}$. For the reverse inclusion let $q \in \mathbb{P}_{\mathbb{Z}_{(p)}}$. Taking a common denominator, write $q = \frac{1}{n}p$, for some $p \in \mathbb{P}_{\mathbb{Z}}$ and $n \in \mathbb{Z}$. Since the minimum common denominator of some elements of $\mathbb{Z}_{(p)}$ is an element of $\mathbb{Z} \setminus p\mathbb{Z}$, then n is not divisible by p . Thus neither $n^2 = N(n)$ is divisible by p . So $n \in S$ and $\mathbb{P}_{\mathbb{Z}}S^{-1} = \mathbb{P}_{\mathbb{Z}_{(p)}}$. In the same manner we can prove that $S^{-1}\mathbb{P}_{\mathbb{Z}} = \mathbb{P}_{\mathbb{Z}_{(p)}}$. \square

Imitating Proposition 1.12 we can give this general result.

Proposition 1.14. *Let R be a commutative ring and let $\mathcal{Q}(R)$ be its total ring of fractions. Then*

$$\mathcal{Q}(\mathbb{P}_R) = \mathbb{P}_{\mathcal{Q}(R)}.$$

Proof. Let S be the set of regular elements of R . Then, S is contained in the center of \mathbb{P}_R , and thus it is a denominator set of \mathbb{P}_R ; it is also easy to see that $S^{-1}\mathbb{P}_R = \mathbb{P}_{S^{-1}R} = \mathbb{P}_{\mathcal{Q}(R)}$ (see the proof of Propositions 1.12).

We claim that the elements of $\mathbb{P}_{\mathcal{Q}(R)}$ are either invertible or zero-divisors. Take $q \in \mathbb{P}_{\mathcal{Q}(R)}$. If $N(q)$ is regular, then it is invertible in $\mathcal{Q}(R)$, and thus $\frac{1}{N(q)}\bar{q} \in \mathbb{P}_{\mathcal{Q}(R)}$ is the inverse of q . Conversely, if $N(q)$ is not regular, then there is $z \in R, z \neq 0$, such that $zN(q) = 0$. If $zq \neq 0$, then also $z\bar{q} = \overline{zq} \neq 0$. So we have that:

$$0 = zN(q) = z\bar{q}q = (z\bar{q})q;$$

hence, q is a zero-divisor.

Thus, $\mathbb{P}_{\mathbb{Q}(R)}$ is a total ring of fractions, and so it is the total ring of fractions of \mathbb{P}_R . \square

2. Integer-valued polynomials

The ring of *integer-valued polynomials* over $\mathbb{P}_{\mathbb{Z}}$ is

$$\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}}) = \{f(x) \in \mathbb{P}_{\mathbb{Q}}[x] \mid f(\mathbb{P}_{\mathbb{Z}}) \subseteq \mathbb{P}_{\mathbb{Z}}\}.$$

This set is actually a ring ([15, Theorem 1.2]), and in [3] the authors describe explicitly some proper ideals of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$. A similar construction can be done if, instead of $\mathbb{P}_{\mathbb{Z}}$, we use $\mathbb{P}_{\mathbb{Z}(p)}$ or $\mathbb{P}_{\mathbb{Q}}$; in the former case, Werner [15, Theorem 1.2] guarantees that $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}(p)})$ is a ring, while in the latter $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Q}}) = \mathbb{P}_{\mathbb{Q}}[x]$ is the whole ring of polynomials (and, in particular, is a ring).

For simplicity of notation, in this Section we will write $\text{Int}(\mathbb{P}_{\mathbb{Z}})$ instead of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$.

A class of ideals of $\text{Int}(\mathbb{P}_{\mathbb{Z}})$ can be constructed in the following way: if $q = a + bi + cj + dk \in \mathbb{P}_{\mathbb{Z}}$ and I is a principal ideal of $\mathbb{P}_{\mathbb{Z}}$ generated by an element of \mathbb{Z} , then

$$\mathfrak{A}_{I,q} := \{f(x) \in \text{Int}(\mathbb{P}_{\mathbb{Z}}) \mid f(z) \in I \ \forall z \in C(q)\},$$

is an ideal of $\text{Int}(\mathbb{P}_{\mathbb{Z}})$, where $C(q) = \{a \pm bi \pm cj \pm dk\}$ ([3, Proposition 4.2]).

If P is a prime ideal of $\text{Int}(\mathbb{P}_{\mathbb{Z}})$, then $P \cap \mathbb{P}_{\mathbb{Z}}$ is a prime ideal of $\mathbb{P}_{\mathbb{Z}}$; since we have a classification of the prime ideals of $\mathbb{P}_{\mathbb{Z}}$ (Theorem 1.2), we can study the spectrum of $\text{Int}(\mathbb{P}_{\mathbb{Z}})$ according to the restriction to $\mathbb{P}_{\mathbb{Z}}$.

Proposition 2.1. *The following hold.*

- (1) [3, Corollary 4.10] *The prime ideals P of $\text{Int}(\mathbb{P}_{\mathbb{Z}})$ with $P \cap \mathbb{P}_{\mathbb{Z}} = (0)$ are exactly those of the form*

$$P = M(x) \cdot \mathbb{P}_{\mathbb{Q}}[x] \cap \text{Int}(\mathbb{P}_{\mathbb{Z}}) =: P_{M(x)},$$

where $M(x) \in \mathbb{Z}[x]$ is an irreducible polynomial.

In particular, if $m_q(x)$ is the minimal polynomial of an element $q \in \mathbb{P}_{\mathbb{Z}}$ then $P_{m_q(x)} = \mathfrak{A}_{0,q}$ is a prime ideal.

- (2) [3, Theorem 4.16] *Let $q := a + b\mathbf{i} + c\mathbf{j} + d\mathbf{k} \in \mathbb{P}_{\mathbb{Z}} \setminus \mathbb{Z}$ and let p be an odd prime. If $\gcd(b, c, d, p) = 1$, then $\mathfrak{A}_{p\mathbb{P}_{\mathbb{Z}},q}$ is prime if and only if $m_q(x)$ is irreducible mod p , in which case $\mathfrak{A}_{p\mathbb{P}_{\mathbb{Z}},q}$ is maximal.*
- (3) [3, Corollary 4.22] *Let $q = a + b\mathbf{i} + c\mathbf{j} + d\mathbf{k} \in \mathbb{P}_{\mathbb{Z}}$, and assume that either $b \equiv c \pmod{2}$ or $b \equiv d \pmod{2}$. Then,*

$$\mathfrak{M}_q := \{f \in \text{Int}(\mathbb{P}_{\mathbb{Z}}) \mid f(q) \in \mathcal{M}\}$$

is a maximal ideal of $\text{Int}(\mathbb{P}_{\mathbb{Z}})$.

Remark 2.2.

- (1) While the first case of the proposition completely classifies the prime ideals above (0) , the other two merely give some examples of the prime ideals above $p\mathbb{P}_{\mathbb{Z}}$ and \mathcal{M} , but not a complete list.
- (2) We refer to [3] for some results about equalities among these ideals.

Lemma 2.3. *The following hold.*

- (1) *If p is an odd prime number and $q \in \mathbb{P}_{\mathbb{Z}}$, then $\mathfrak{A}_{m_q(x)} \subset \mathfrak{A}_{p\mathbb{P}_{\mathbb{Z}},q}$.*

(2) If $q \in \mathbb{P}_{\mathbb{Z}}$ is as in Proposition 2.1(3), then $\mathfrak{P}_{m_q(x)} \subset \mathfrak{M}_q$.

Proof. Let $f(x) \in \mathfrak{P}_{m_q(x)}$: then, $f(x) = m_q(x)g(x)$ for some $g(x) \in \mathbb{P}_{\mathbb{Q}}[x]$. Since $m_q(x)$ has coefficients in the center of $\mathbb{P}_{\mathbb{Q}}$, we have $f(q) = m_q(q)g(q) = 0$. Hence, $f(x) \in \mathfrak{M}_q$; furthermore, $m_q(q') = 0$ for all $q' \in C(q)$ (since the elements of $C(q)$ have the same minimal polynomial of q [3, paragraph after Definition 4.1]) and thus $f(x) \in \mathfrak{P}_{p\mathbb{P}_{\mathbb{Z}}, q}$. Therefore $\mathfrak{P}_{m_q(x)}$ is contained in both $\mathfrak{P}_{p\mathbb{P}_{\mathbb{Z}}, q}$ and \mathfrak{M}_q .

By intersecting the ideals with \mathbb{Z} , it is easily seen that the inclusions are proper. □

When D is a Noetherian commutative domain, the integer-valued polynomials over D behave well with respect to the localization, that is, if S is a multiplicative subset of D then $S^{-1}\text{Int}(D) = \text{Int}(S^{-1}D)$ ([2, Theorem 1.2.3]). In [3, Theorem 3.4] an analogous result has been showed for $\text{Int}(\mathbb{P}_{\mathbb{Z}})$ when S is a multiplicatively closed subset $S \subseteq \mathbb{Z}$ (it is central). In the following we prove that $\text{Int}(\mathbb{P}_{\mathbb{Z}})$ behaves well with respect to localization also for denominator sets whose elements are not necessarily central, as $S = \mathcal{C}(Q)$, where Q is a prime ideal of $\mathbb{P}_{\mathbb{Z}}$.

Theorem 2.4. *Let Q be a prime ideal of $\mathbb{P}_{\mathbb{Z}}$ and let $S = \mathcal{C}(Q)$. Then S is also a denominator set of $\text{Int}(\mathbb{P}_{\mathbb{Z}})$ and $\text{Int}(\mathbb{P}_{\mathbb{Z}})S^{-1} = \text{Int}(\mathbb{P}_{\mathbb{Z}}S^{-1})$.*

Proof. To prove that S is a denominator set of $\text{Int}(\mathbb{P}_{\mathbb{Z}})$ it is sufficient to use the same argument of Lemma 1.6 and Proposition 1.11, observing that $N(s)$ is in the center of $\text{Int}(\mathbb{P}_{\mathbb{Z}})$ for each $s \in S$.

Let Q be a prime ideal of $\mathbb{P}_{\mathbb{Z}}$, and let $Q \cap \mathbb{Z} = p\mathbb{Z}$ (where p is either a prime number or 0). Set $T := \mathbb{Z} \setminus p\mathbb{Z}$. By Propositions 1.12 and 1.13, we have $\text{Int}(\mathbb{P}_{\mathbb{Z}}T^{-1}) = \text{Int}(\mathbb{P}_{\mathbb{Z}}\mathcal{C}(Q)^{-1}) = \text{Int}(\mathbb{P}_{\mathbb{Z}(p)})$.

To prove the statement it is enough to show that

$$(1) \quad \text{Int}(\mathbb{P}_{\mathbb{Z}})T^{-1} \subseteq \text{Int}(\mathbb{P}_{\mathbb{Z}})\mathcal{C}(Q)^{-1} \subseteq \text{Int}(\mathbb{P}_{\mathbb{Z}(p)}) \subseteq \text{Int}(\mathbb{P}_{\mathbb{Z}})T^{-1}.$$

The first inclusion follows from the fact that $T \subseteq \mathcal{C}(Q)$, while the last one from [3, Theorem 3.4] (it is actually an equality). Thus, we only need to prove that $\text{Int}(\mathbb{P}_{\mathbb{Z}})\mathcal{C}(Q)^{-1} \subseteq \text{Int}(\mathbb{P}_{\mathbb{Z}(p)})$. Again by [3, Theorem 3.4], we have $\text{Int}(\mathbb{P}_{\mathbb{Z}}) \subseteq \text{Int}(\mathbb{P}_{\mathbb{Z}(p)})$; furthermore, each element of $\mathcal{C}(Q)$ becomes invertible in $\mathbb{P}_{\mathbb{Z}(p)}$ and thus in $\text{Int}(\mathbb{P}_{\mathbb{Z}(p)})$. Hence, $\text{Int}(\mathbb{P}_{\mathbb{Z}})\mathcal{C}(Q)^{-1} \subseteq \text{Int}(\mathbb{P}_{\mathbb{Z}(p)})$ and all the containments must be equalities. □

Note that the exact same argument can be used if we localize on the left: if S is $S = \mathcal{C}^l(Q)$ then $S^{-1}\text{Int}(\mathbb{P}_{\mathbb{Z}}) = \text{Int}(S^{-1}\mathbb{P}_{\mathbb{Z}})$.

Corollary 2.5. *The following hold.*

- (1) If $S = \mathcal{R}(\mathbb{P}_{\mathbb{Z}})$ or $S = \mathbb{Z} \setminus (0)$ then $\text{Int}(\mathbb{P}_{\mathbb{Z}})S^{-1} = \text{Int}(\mathbb{P}_{\mathbb{Q}}) = \mathbb{P}_{\mathbb{Q}}[x]$.
- (2) If p is a prime number and $S = \mathbb{Z} \setminus p\mathbb{Z}$ or $S = \mathcal{C}(Q)$, with Q a prime ideal of $\mathbb{P}_{\mathbb{Z}}$ such that $Q \cap \mathbb{Z} = p\mathbb{Z}$, then $\text{Int}(\mathbb{P}_{\mathbb{Z}})S^{-1} = \text{Int}(\mathbb{P}_{\mathbb{Z}(p)})$.

Proof. For the first point, the equality $\text{Int}(\mathbb{P}_{\mathbb{Z}})S^{-1} = \text{Int}(\mathbb{P}_{\mathbb{Q}})$ follows from Theorem 2.4 and Proposition 1.12. The equality $\text{Int}(\mathbb{P}_{\mathbb{Q}}) = \mathbb{P}_{\mathbb{Q}}[x]$ follows directly from the definitions.

Similarly, the second point follows from Theorem 2.4 and from Proposition 1.13. □

These results allow us to represent $\mathbb{P}_{\mathbb{Z}}$ and $\text{Int}(\mathbb{P}_{\mathbb{Z}})$ as intersection of localizations.

Proposition 2.6. *Let \mathcal{P} be the set of prime numbers. Then, the following hold.*

$$\mathbb{P}_{\mathbb{Z}} = \bigcap_{p \in \mathcal{P}} \mathbb{P}_{\mathbb{Z}(p)}. \tag{1}$$

$$\text{Int}(\mathbb{P}_{\mathbb{Z}}) = \bigcap_{p \in \mathcal{P}} \text{Int}(\mathbb{P}_{\mathbb{Z}(p)}). \tag{2}$$

Proof. (1) The inclusion (\subseteq) is obvious since for every prime p , $\mathbb{P}_{\mathbb{Z}} \subseteq \mathbb{P}_{\mathbb{Z}(p)}$. For the reverse inclusion, take an element $\mathbf{q} = a + b \mathbf{i} + c \mathbf{j} + d \mathbf{k}$ of the intersection. Then $a, b, c, d \in \cap_p \mathbb{Z}(p) = \mathbb{Z}$ and $\mathbf{q} \in \mathbb{P}_{\mathbb{Z}}$.

(2) For all primes p , let Q_p be the maximal ideal of $\mathbb{P}_{\mathbb{Z}}$ above p . We have that $\text{Int}(\mathbb{P}_{\mathbb{Z}}) \subseteq (\text{Int}(\mathbb{P}_{\mathbb{Z}}))_{\mathcal{C}}(Q_p)^{-1} = \text{Int}(\mathbb{P}_{\mathbb{Z}(p)})$, and thus $\text{Int}(\mathbb{P}_{\mathbb{Z}})$ is inside the intersection. Conversely, if $f(x)$ belongs to the intersection and $\mathbf{q} \in \mathbb{P}_{\mathbb{Z}}$, then $f(\mathbf{q}) \in \mathbb{P}_{\mathbb{Z}(p)}$ for every prime number p , and thus $f(\mathbf{q}) \in \cap_p \mathbb{P}_{\mathbb{Z}(p)} = \mathbb{P}_{\mathbb{Z}}$ (by the previous point) and $f(x) \in \text{Int}(\mathbb{P}_{\mathbb{Z}})$. \square

3. Matrix representations

To study the spectrum of $\text{Int}(\mathbb{P}_{\mathbb{Z}})$, we introduce the related commutative ring

$$\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}}) := \{f(x) \in \mathbb{Q}[x] \mid \forall \mathbf{q} \in \mathbb{P}_{\mathbb{Z}}: f(\mathbf{q}) \in \mathbb{P}_{\mathbb{Z}}\},$$

and we define similarly $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$. These sets are easily seen to be rings by using polynomial evaluation. To avoid confusion in the notation, from now in we will go back to write $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$ and $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}(p)})$ for $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ and $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$, respectively. Note that, if we consider $\mathbb{Q}[x]$ as subring of $\mathbb{P}_{\mathbb{Q}}[x]$ in the obvious way, then $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}}) = \text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}}) \cap \mathbb{Q}[x]$.

The relation between $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ and $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$ passes through a matrix representation of the rings $\mathbb{P}_{\mathbb{Z}}$ and $\mathbb{P}_{\mathbb{Z}(p)}$. We denote by $\mathcal{M}_n(R)$ the ring of matrices of order n over R .

Proposition 3.1. [3, Proposition 2.2] *The following hold.*

- (1) *Let R be a commutative ring with identity such that 2 is a unit of R . Then, $\mathbb{P}_R \cong \mathcal{M}_2(R)$ as R -algebras.*
- (2) *Let $\mathcal{A} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a \equiv d, b \equiv c \pmod{2} \right\} \subseteq \mathcal{M}_2(\mathbb{Z})$. Then, $\mathbb{P}_{\mathbb{Z}} \cong \mathcal{A}$ as \mathbb{Z} -algebras.*

Let D be a domain with quotient field K . We define

$$\text{Int}_K(\mathcal{M}_n(D)) := \{f(x) \in K[x] \mid \forall A \in \mathcal{M}_n(D): f(A) \in \mathcal{M}_n(D)\}$$

and

$$\begin{aligned} \text{Int}_{\mathcal{M}_n(K)}(\mathcal{M}_n(D)) &:= \\ &= \{f(x) \in \mathcal{M}_n(K)[x] \mid \forall A \in \mathcal{M}_n(D): f(A) \in \mathcal{M}_n(D)\}. \end{aligned}$$

These rings roughly correspond, respectively, to $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ and $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$.

Proposition 3.2. *Let D be a domain with quotient field K . Then*

$$\text{Int}_{\mathcal{M}_n(K)}(\mathcal{M}_n(D)) \simeq \mathcal{M}_n(\text{Int}_K(\mathcal{M}_n(D))).$$

Moreover, the following hold.

- (i) *The ideals of $\text{Int}_{\mathcal{M}_n(K)}(\mathcal{M}_n(D))$ are in 1-1 correspondence with the sets of the form $\mathcal{M}_n(\mathcal{I})$, where \mathcal{I} is an ideal of $\text{Int}_K(\mathcal{M}_n(D))$.*
- (ii) *The prime ideals of $\text{Int}_{\mathcal{M}_n(K)}(\mathcal{M}_n(D))$ are in 1-1 correspondence with the sets of the form $\mathcal{M}_n(\mathcal{P})$, where \mathcal{P} is a prime ideal of $\text{Int}_K(\mathcal{M}_n(D))$.*
- (iii) *The maximal ideals of $\text{Int}_{\mathcal{M}_n(K)}(\mathcal{M}_n(D))$ are in 1-1 correspondence with the sets of the form $\mathcal{M}_n(\mathcal{M})$, where \mathcal{M} is a maximal ideal of $\text{Int}_K(\mathcal{M}_n(D))$.*

Proof. See Frisch [4, Theorem 7.2] and Frisch [4, Theorem 7.3]. The remaining part follows from Lam [8, Theorem 3.1]. \square

Putting together these two results, we have the following theorem.

Theorem 3.3. *Let p be an odd prime integer. Then, the prime ideals of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}(p)})$ are in 1-1 correspondence with the prime ideals of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$.*

Proof. By Proposition 3.1, $\mathbb{P}_{\mathbb{Q}} \simeq \mathcal{M}_2(\mathbb{Q})$, and the isomorphism brings $\mathbb{P}_{\mathbb{Z}(p)}$ into $\mathcal{M}_2(\mathbb{Z}(p))$. By Proposition 3.2,

$$\begin{aligned} \text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}(p)}) &\simeq \text{Int}_{\mathcal{M}_2(\mathbb{Q})}(\mathcal{M}_2(\mathbb{Z}(p))) \simeq \\ &\simeq \mathcal{M}_2(\text{Int}_{\mathbb{Q}}(\mathcal{M}_2(\mathbb{Z}(p)))) \simeq \mathcal{M}_2(\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})); \end{aligned}$$

thus the prime ideals of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}(p)})$ are in bijective correspondence with the prime ideals of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$, as claimed. \square

The main advantage of this theorem is that $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$ is a commutative ring properly contained in between the two well-studied rings $\mathbb{Z}[x]$ and $\mathbb{Q}[x]$.

Proposition 3.4. *The nonzero prime ideals P of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ such that $P \cap \mathbb{Z} = (0)$ are pairwise uncomparable.*

Proof. It is enough to notice that $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})S^{-1} = \mathbb{Q}[x]$, if $S = \mathbb{Z} \setminus (0)$. Then the prime ideals of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ contracting to (0) in \mathbb{Z} are in 1-1 correspondence with the prime ideals of $\mathbb{Q}[x]$, which are not comparable (except (0)). This correspondence preserves the order, thus the statement follows. \square

Theorem 3.5. *The rings $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ and $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$, for any prime p , have dimension 2.*

Proof. The ring $\mathbb{Z}[x]$ is a Noetherian ring of dimension 2, and thus by Gilmer [5, Theorem 30.9 and Corollary 30.10] every overring of $\mathbb{Z}[x]$ has dimension at most 2. Since $\mathbb{Z}[x] \subseteq \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}}) \subseteq \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$, it follows that the dimensions of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ and $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$ are at most 2.

Fix now a prime p . By Proposition 2.1, we can find an integer split-quaternion $q = a + b \mathbf{i} + c \mathbf{j} + d \mathbf{k} \in \mathbb{P}_{\mathbb{Z}}$ such that its minimal polynomial $m_q(x)$ is irreducible modulo p and such that $\mathfrak{P}_{p\mathbb{Z},q}$ (if $p \neq 2$) or \mathfrak{M}_q (if $p = 2$) is a maximal ideal of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$ over p . Then, its restriction $P_2 := \mathfrak{P}_{p\mathbb{Z},q} \cap \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ or $P_2 := \mathfrak{M}_q \cap \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ is a prime ideal of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$.

By construction, $m_q(x)$ is irreducible in $\mathbb{Q}[x]$, and thus $\mathfrak{P}_{m_q(x)}$ is a prime ideal of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$, which by Lemma 2.3 is contained inside $\mathfrak{P}_{p\mathbb{Z},q}$ and \mathfrak{M}_q ; hence, $P_1 := \mathfrak{P}_{m_q(x)} \cap \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ is a prime ideal of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ inside P_2 . Therefore, $(0) \subseteq P_1 \subseteq P_2$ is a chain of prime ideals of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$; furthermore, the inclusions are strict since $m_q(x) \in P_1 \setminus (0)$, while $p \in P_2 \setminus P_1$. Hence, $(0) \subsetneq P_1 \subsetneq P_2$ is a chain of length 2, and thus $\dim \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}}) \geq 2$; therefore, the dimension must be exactly 2.

For $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$, it is enough to note that the chain $(0) \subsetneq P_1 \subsetneq P_2$ lifts to a chain in $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$ (since this ring is a localization of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$) and thus also $\dim(\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})) \geq 2$ and $\dim(\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})) = 2$. \square

Note that this replicates the same pattern of $\text{Spec}(\text{Int}(\mathbb{Z}))$ shown in [2, Proposition V.2.7]. The correspondence with matrix rings allow also to say something about $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}(p)})$.

Corollary 3.6. *If p is an odd prime, then $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}(p)})$ has dimension 2. Furthermore, $\dim(\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})) \geq 2$.*

Proof. By Theorem 3.3, the dimension of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}(p)})$ is the same of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$, which is 2 by Theorem 3.5. The last claim follows since $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}(p)})$ is a localization of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$. \square

Theorem 3.3 does not work for $p = 2$, and thus the previous results do not allow to calculate the dimension of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$. However, we conjecture that $\dim(\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})) = \dim(\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}(2)})) = 2$.

An important difference between $\text{Int}(\mathbb{Z})$ and $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ is that the latter is not integrally closed (and thus it is not a Prüfer domain); see [Corollary 3.8](#) below. However, we can describe its integral closure by using algebraic integers.

Given a finite degree extension $\mathbb{Q}(\theta)$ of \mathbb{Q} , we indicate by \mathcal{A}_{θ} the ring of algebraic integers of $\mathbb{Q}(\theta)$. If $n \in \mathbb{N}$ is positive, the set of all algebraic integers of degree at most n over \mathbb{Q} is

$$\mathcal{A}_n := \bigcup_{[\mathbb{Q}(\theta):\mathbb{Q}] \leq n} \mathcal{A}_{\theta};$$

similarly, if p is a prime number, we denote by $\mathcal{A}_{n,p}$ is the set of algebraic numbers that are root of a monic irreducible polynomial of degree n over $\mathbb{Z}_{(p)}$.

In [10] the authors define the set of integer-valued polynomials over \mathcal{A}_n with rational coefficients to be the set

$$\text{Int}(\mathcal{A}_n) := \bigcap_{\theta \in \mathcal{A}_n} \text{Int}_{\mathbb{Q}}(\mathcal{A}_{\theta}).$$

The ring $\text{Int}(\mathcal{A}_n)$ can be seen as the set of all polynomials with rational coefficients that map \mathcal{A}_n into \mathcal{A}_n . They also show that $\text{Int}_{\mathbb{Q}}(\mathcal{A}_n)$ is a Prüfer domain for every n ([10, Theorem 3.9]).

Theorem 3.7. *Let p be an odd prime integer. Then $\text{Int}_{\mathbb{Q}}(\mathcal{A}_2)_{(p)} = \text{Int}_{\mathbb{Q}}(\mathcal{A}_{2,p})$ is the integral closure of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}_{(p)}})$ in $\mathbb{Q}[x]$.*

Proof. By Loper and Werner [10, Theorem 4.6], $\text{Int}_{\mathbb{Q}}(\mathcal{A}_2)$ is the integral closure of $\text{Int}_{\mathbb{Q}}(\mathcal{M}_2(\mathbb{Z}))$. Using [Proposition 3.1](#), and recalling that the localization at prime integers preserves the integral closure, we have that:

$$\begin{aligned} \text{Int}_{\mathbb{Q}}(\mathcal{A}_2)_{(p)} &= \overline{\text{Int}_{\mathbb{Q}}(\mathcal{M}_2(\mathbb{Z}))}_{(p)} = \overline{\text{Int}_{\mathbb{Q}}(\mathcal{M}_2(\mathbb{Z}))}_{(p)} = \\ &= \overline{\text{Int}_{\mathbb{Q}}(\mathcal{M}_2(\mathbb{Z}))}_{(p)} = \overline{\text{Int}_{\mathbb{Q}}(\mathcal{M}_2(\mathbb{Z}_{(p)}))} = \overline{\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}_{(p)}})}. \end{aligned}$$

Finally, using [11, Theorem 13] with $A = \mathbb{P}_{\mathbb{Z}_{(p)}}$, we have that $\overline{\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}_{(p)}})}$ is also the integral closure of $\text{Int}_{\mathbb{Q}}(\mathcal{A}_{2,p})$. \square

Corollary 3.8. *The ring $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ is not integrally closed.*

Proof. If $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ is integrally closed, then its localization at an odd prime p , $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}_{(p)}})$, is integrally closed too. Thus, from [Theorem 3.7](#), $\overline{\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}_{(p)}})} = \text{Int}_{\mathbb{Q}}(\mathcal{A}_2)_{(p)}$ and this is Prüfer. Since $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}_{(p)}}) \cong \text{Int}_{\mathbb{Q}}(\mathcal{M}_2(\mathbb{Z}_{(p)}))$, it follows that the ring

$$\text{Int}_{\mathbb{Q}}(B, \mathcal{M}_2(\mathbb{Z}_{(p)})) := \{f \in \mathbb{Q}[x] \mid f(B) \in \mathcal{M}_2(\mathbb{Z}_{(p)})\}$$

is an overring of $\text{Int}_{\mathbb{Q}}(\mathcal{M}_2(\mathbb{Z}_{(p)}))$, for every matrix $B \in \mathcal{M}_2(\mathbb{Z}_{(p)})$. Taking $B = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and arguing as in [10, §4], it can be shown that $\text{Int}_{\mathbb{Q}}(B, \mathcal{M}_2(\mathbb{Z}_{(p)}))$ is not integrally closed. The claim follows. \square

4. The ideal $p\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$

In this section we study in more detail the ideal $p\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ generated by a prime number p (not necessarily odd). Our first result can be seen as a refinement of the proof of [Theorem 3.5](#).

Proposition 4.1. *Let p be a prime number. Then, every prime ideal of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ containing p is maximal.*

Proof. We follow the proof of Cahen and Chabert [2, Lemma V.1.9].

Let u_1, \dots, u_k be a set of residues of $\mathbb{P}_{\mathbb{Z}}/p\mathbb{P}_{\mathbb{Z}}$ (with $k = p^4$), and let P be a prime ideal of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$ containing p . Take any $a(x) \in \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})$, and let $a_i(x) := a(x) - u_i$. Let $b(x) := a_1(x) \cdots a_k(x)$: by construction, for every $q \in \mathbb{P}_{\mathbb{Z}}$ there is an i such that $a(q) \equiv u_i \pmod{p\mathbb{P}_{\mathbb{Z}}}$.

Since the a_i have coefficients in the commutative ring \mathbb{Q} , we have $b(q) = a_1(q) \cdots a_k(q)$: hence, $b(q) \in p\mathbb{P}_{\mathbb{Z}}$ and so $b(x) \in p\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}}) \subseteq P$; since P is prime, there must be an i such that $a_i(x) \in P$. However, $a_i(x) \equiv u_i \pmod{p\mathbb{P}_{\mathbb{Z}}}$, and thus $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}})/P$ is isomorphic to $\mathbb{P}_{\mathbb{Z}}/p\mathbb{P}_{\mathbb{Z}} \simeq \mathbb{P}_{\mathbb{Z}_p}$. Hence, P is maximal, as claimed. □

Corollary 4.2. *Let p be an odd prime integer. Then, every prime ideal of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}(p)})$ containing p is maximal.*

Proof. It is enough to use Proposition 4.1 and the correspondence of Theorem 3.3. □

Remark 4.3.

- (1) The previous two results allow to give an alternative proof of Theorem 3.5. Indeed, if $(0) \subsetneq Q_1 \subsetneq Q_2 \subsetneq Q_3$ is a chain of prime ideals of length 3, then either $Q_1 \cap \mathbb{Z} = Q_2 \cap \mathbb{Z} = (0)$ or $Q_2 \cap \mathbb{Z} = Q_3 \cap \mathbb{Z} = p\mathbb{Z}$, for some prime number p . The latter case is made impossible by Proposition 4.1 (as Q_2 contains p but is not maximal); on the other hand the former case would imply that two nonzero prime ideals of $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$ over (0) are comparable, against Proposition 3.4.
- (2) The proof of Proposition 4.1 does *not* work in the ring $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$, since the evaluation of a product of polynomials cannot be done separately for each factor, and thus $b(q) \neq a_1(q) \cdots a_k(q)$ in general. Nevertheless, we conjecture (but we don't have a proof) that the same property holds also in $\text{Int}_{\mathbb{P}_{\mathbb{Q}}}(\mathbb{P}_{\mathbb{Z}})$.

A consequence of Proposition 4.1 is that the ideals $p\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$ are not prime. We now want to find an explicit description of the polynomials in $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$ and, as a corollary, to find two polynomials outside $p\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$ whose product is inside the ideal.

Proposition 4.4. *Let R, S be commutative rings and let $\pi: R \rightarrow S$ be a homomorphism. Then, the natural map*

$$\begin{aligned} \varphi: \mathbb{P}_R &\rightarrow \mathbb{P}_S \\ a + b \mathbf{i} + c \mathbf{j} + d \mathbf{k} &\mapsto \pi(a) + \pi(b) \mathbf{i} + \pi(c) \mathbf{j} + \pi(d) \mathbf{k} \end{aligned}$$

is a ring homomorphism. Furthermore, if π is surjective then φ is surjective and $\ker \varphi = (\ker \pi)\mathbb{P}_R = \mathbb{P}_{\ker \pi} = \{a + b \mathbf{i} + c \mathbf{j} + d \mathbf{k} \mid a, b, c, d \in \ker \pi\}$; in particular, $\mathbb{P}_R/\ker \varphi \simeq \mathbb{P}_S$.

Proof. Straightforward. □

An important particular case is when $R = \mathbb{Z}$ or $R = \mathbb{Z}_{(p)}$ and $S = \mathbb{Z}_p$: in this case, the kernel of π is generated by p , and thus we obtain the well-known isomorphisms $\frac{\mathbb{P}_{\mathbb{Z}}}{p\mathbb{P}_{\mathbb{Z}}} \simeq \frac{\mathbb{P}_{\mathbb{Z}(p)}}{p\mathbb{P}_{\mathbb{Z}(p)}} \simeq \mathbb{P}_{\mathbb{Z}_p}$.

In particular, the previous proposition shows that polynomial evaluation behaves well with respect to quotients. Given a surjection $\pi: R \rightarrow S$ and a polynomial $f(x) = \sum_{t=0}^n p_t x^t \in R[x]$, we denote by $\bar{f}(x) = \sum_{t=0}^n \pi(p_t) x^t \in S[x]$ the polynomial obtained by reducing the coefficients modulo $\ker \varphi$. Then, for every $q \in \mathbb{P}_{\mathbb{Z}}$, we have $\pi(f(q)) = \bar{f}(\pi(q))$.

Proposition 4.5. *Let p be a prime integer. Let $f(x) \in \mathbb{Z}[x]$ and $\bar{f}(x) \in \mathbb{Z}_p[x]$ be as above. Given an integer $n > 1$ such that $n = p^{\alpha}m$ with $p \nmid m$, then $\frac{1}{n}f(x) \in \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$ if and only if $f(q) \in p^{\alpha}\mathbb{P}_{\mathbb{Z}(p)}$, for all $q \in \mathbb{P}_{\mathbb{Z}(p)}$. In particular if $\alpha = 1$, $\frac{1}{n}f(x) \in \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$ if and only if $f(q) = 0$ in $\mathbb{P}_{\mathbb{Z}_p}$, for all $q \in \mathbb{P}_{\mathbb{Z}_p}$.*

Proof. We have that

$$\frac{1}{n}f(x) \in \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}_{(p)}}) \iff \frac{1}{n}f(\mathfrak{q}) \in \mathbb{P}_{\mathbb{Z}_{(p)}} \quad \forall \mathfrak{q} \in \mathbb{P}_{\mathbb{Z}_{(p)}} \iff f(\mathfrak{q}) \in n\mathbb{P}_{\mathbb{Z}_{(p)}} \quad \forall \mathfrak{q} \in \mathbb{P}_{\mathbb{Z}_{(p)}}.$$

Since $p \nmid m$, we have $n\mathbb{P}_{\mathbb{Z}_{(p)}} = p^{\alpha}\mathbb{P}_{\mathbb{Z}_{(p)}}$. □

Lemma 4.6. *Let R be a commutative domain. Take $\mathfrak{q} \in \mathbb{P}_R \setminus R$ and let $m_{\mathfrak{q}}(x) \in R[x]$ be its minimal polynomial over R . If a polynomial $f(x) \in R[x]$ is such that $f(\mathfrak{q}) = 0$, then $m_{\mathfrak{q}}(x) \mid f(x)$ in $R[x]$.*

Proof. Since $m_{\mathfrak{q}}(x)$ is monic we can divide $f(x)$ by $m_{\mathfrak{q}}(x)$ obtaining

$$f(x) = g(x)m_{\mathfrak{q}}(x) + r(x),$$

for some $g(x), r(x) \in R[x]$. In particular $r(x) = ax + b$ is linear as $m_{\mathfrak{q}}(x)$ is of degree two. Since $R[x]$ is contained in the center of $\mathbb{P}_R[x]$, we can evaluate the polynomial relation above in \mathfrak{q} , obtaining $0 = f(\mathfrak{q}) = g(\mathfrak{q}) \cdot 0 + a\mathfrak{q} + b$. Since R is a domain and $\mathfrak{q} \notin R$, necessarily $a = b = 0$. □

We observe that Lemma 4.6 does not hold if $f(x) \in \mathbb{P}_R[x] \setminus R[x]$. For example, consider $\mathfrak{i} \in \mathbb{P}_{\mathbb{Z}}$ and $f(x) = x^3 + \mathfrak{i}x + (\mathfrak{i} + 1)x + \mathfrak{i} + 1$. Then $f(\mathfrak{i}) = 0$ but $f(x) = (x^2 + 1)(x + \mathfrak{i}) + \mathfrak{i}x + 1$ and the remainder is nonzero.

Corollary 4.7. *With the hypothesis and notation of Proposition 4.5, let p be a prime integer and $n = pm$ with $p \nmid m$. Then $\frac{1}{n}f(x) \in \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}_{(p)}})$ if and only if $\bar{f}(x)$ is divided by all the minimal polynomials of the elements of $\mathbb{P}_{\mathbb{Z}_p}$.*

Proof. It is an immediate consequence of Proposition 4.5 and Lemma 4.6. □

Using the previous Corollary we can construct a nontrivial element of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}_{(p)}})$.

Example 4.8. *The polynomial*

$$\Phi_p(x) = \frac{1}{p}(x^p - x)(x^{p^2} - x)$$

belongs to $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}_{(p)}})$.

By Proposition 4.5, it is sufficient to show that $f(x) = (x^p - x)(x^{p^2} - x) \in \mathbb{Z}[x]$ vanishes over all elements of $\mathbb{P}_{\mathbb{Z}_p}$. Observe that every monic and irreducible polynomial of $\mathbb{Z}_p[x]$ of degree one or two is a factor of $f(x)$. In particular, if $g(x)$ is a linear polynomial then $g(x)^2$ divides $f(x)$, since $g(x)$ divides both $x^p - x$ and $x^{p^2} - x$. By Corollary 4.7, this also means that the minimal polynomial of every split-quaternion of $\mathbb{P}_{\mathbb{Z}_p}$ is a factor of $f(x)$.

In particular we can show that every monic and quadratic polynomial of $\mathbb{Z}_p[x]$ is the minimal polynomial for some element of $\mathbb{P}_{\mathbb{Z}_p}$. The proof is *mutatis mutandis* the same as the proof of [14, Lemma 3.5]. This means that the polynomial $\Phi_p(x)$ does not contain any redundant factor.

Proposition 4.9. *With the above notation we have the following proper inclusions:*

$$\mathbb{Z}_{(p)}[x] \subsetneq \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}_{(p)}}) \subsetneq \text{Int}(\mathbb{Z}_{(p)}).$$

Proof. The first inclusion follows from the fact that $\mathbb{Z} \subseteq \mathbb{P}_{\mathbb{Z}}$ and thus $\mathbb{Z}_{(p)}[x] \subseteq \mathbb{P}_{\mathbb{Z}_{(p)}}[x] \cap \mathbb{Q}[x] \subseteq \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}_{(p)}})$. It is proper since the polynomial $\Phi_p(x)$ given in Example 4.8 belongs to $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}_{(p)}})$ but not to $\mathbb{Z}_{(p)}[X]$.

The second inclusion is straightforward since $\mathbb{Z}_{(p)} = \mathbb{P}_{\mathbb{Z}_{(p)}} \cap \mathbb{Q}$.

To see that it is proper, consider the “binomial polynomial” $f(x) = \frac{x(x-1)(x-2)\dots(x-p+1)}{p} \in \text{Int}(\mathbb{Z}_{(p)})$. If $p = 2$ then $f(\mathfrak{i}) = \frac{-1-\mathfrak{i}}{2} \notin \mathbb{P}_{\mathbb{Z}_{(2)}}$; if p is odd then $pf(x)$ is not divided by $x^2 + 1$

(the minimal polynomial of \mathbf{i}), and thus $f(\mathbf{i}) \notin \mathbb{P}_{\mathbb{Z}(p)}$ by Corollary 4.7. It follows that $f(x) \notin \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$. □

The fact that the two containments of the previous proposition are strict also follows from [12, Theorem 2.12] (the first one) and [13, Theorem 2.11] (the second one).

Proposition 4.10. *The ideal $p\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$ is not a prime ideal of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$.*

Proof. Let us consider the polynomials:

$$\begin{aligned} f(x) &= (x^p - x)^2 \in \mathbb{Z}[x], \\ g(x) &= \frac{1}{p}(x^{p^2} - x)^2 \in \mathbb{Q}[x], \\ F(x) &= f(x)g(x) = \frac{1}{p}(x^p - x)^2(x^{p^2} - x)^2 \in \mathbb{Q}[x]. \end{aligned}$$

These three polynomials are elements of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$. Indeed, for $f(x)$ it follows from the inclusion $\mathbb{Z}[x] \subseteq \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$. For $F(x)$ and $g(x)$ observe that they are equal to $\Phi_p(x)$ (Example 4.8) multiplied by a polynomial with integer coefficients.

We claim that $F(x) \in p\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$ while $f(x)$ and $g(x)$ do not belong to this ideal.

Indeed, $\frac{1}{p}F(x) = (\Phi_p(x))^2 \in \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$ and thus $F(x) \in p\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$.

As regards $f(x)$, we have that $\tilde{f}(x)$ is not divisible by any quadratic irreducible polynomial over \mathbb{Z}_p , and thus by Corollary 4.7 $\frac{1}{p}f(x) \notin \text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$.

For $g(x)$, consider $\frac{1}{p}g(x) = \frac{1}{p^2}(x^{p^2} - x)^2$. If $p = 2$ then $\frac{1}{2}g(\mathbf{i}) = -\frac{\mathbf{i}}{2} \notin \mathbb{P}_{\mathbb{Z}(2)}$. If p is odd, then we set $\mathbf{q} := \mathbf{i} + (p - 1)\mathbf{k}$. We have that $\mathbf{q}^2 = p^2 - 2p$, and if we raise \mathbf{q} to an even power greater than 2, we obtain an integer divisible by p^2 . Since $\frac{1}{p}g(x)$ is a central polynomial, we can evaluate it in \mathbf{q} using its factorization. Thus, we have

$$\frac{1}{p}g(\mathbf{q}) = \frac{(q^{p^2} - q)^2}{p^2} = \frac{q^{2p^2} + q^2 - 2q^{p^2+1}}{p^2} = m + \frac{p - 2}{p} \notin \mathbb{P}_{\mathbb{Z}(p)}$$

for some $m \in \mathbb{Z}$.

Since $F(x) = f(x)g(x)$, we can conclude that $p\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$ is not a prime ideal of $\text{Int}_{\mathbb{Q}}(\mathbb{P}_{\mathbb{Z}(p)})$. □

Acknowledgements

We would like to thank the referee for his/her useful suggestions.

Funding

The second and third authors were partially supported by GNSAGA of Istituto Nazionale di Alta Matematica.

References

- [1] Allen, D. B. (1989). *Notes on Localization in Noncommutative Noetherian Rings*. Cuadernos de Algebra, 9. Granada: University of Granada.
- [2] Cahen, P. J., Chabert, J. L. (1997). *Integer-valued polynomials*. Providence, RI: Amer. Math. Soc. Surveys and Monographs.
- [3] Cigliola, A., Loper, K.A., Werner, N. (2015). Split quaternions and integer-valued polynomials. *Commun. Algebra* 43(1):182–196. DOI: [10.1080/00927872.2014.897561](https://doi.org/10.1080/00927872.2014.897561).

- [4] Frisch, S. (2013). Integer-valued polynomials on algebras. *J. Algebra* 373:414–425. DOI: [10.1016/j.jalgebra.2012.10.003](https://doi.org/10.1016/j.jalgebra.2012.10.003).
- [5] Gilmer, R. (1992). *Multiplicative ideal theory*. Queen's Papers in Pure and Applied Mathematics, 90, Corrected Reprint of the 1972 Edition. Kingston, ON: Queen's University.
- [6] Goldie, A. W. (1967). Localization in noncommutative Noetherian rings. *J. Algebra* 5(1):89–105. DOI: [10.1016/0021-8693\(67\)90028-2](https://doi.org/10.1016/0021-8693(67)90028-2).
- [7] Goodearl, K. R., Warfield, R. B. Jr., (2004). *An Introduction to Noncommutative Noetherian Rings*. Cambridge: Cambridge University Press.
- [8] Lam, T. Y. (2001). *A First Course in Noncommutative Rings*. Mathematics, 131. New York: Springer-Verlag.
- [9] Lam, T. Y. (1999). *Lectures on Modules and Rings*. Graduate Texts in Mathematics, 189. New York: Springer-Verlag.
- [10] Loper, K. A., Werner, N. (2012). Generalized rings of integer-valued polynomials. *J. Num. Theory* 132(11): 2481–2490. DOI: [10.1016/j.jnt.2012.05.009](https://doi.org/10.1016/j.jnt.2012.05.009).
- [11] Peruginelli, G., Werner, J. N. (2014). Integral closure of rings of integer-valued polynomials on algebras. In: Fontana, M., Frisch, S., Glaz, S., eds. *Commutative Algebra: Recent Advances in Commutative Rings, Integer-Valued Polynomials, and Polynomial Functions*. New York: Springer, p. 293–305.
- [12] Peruginelli, G., Werner, N. J. (2017). Non-triviality conditions for integer-valued polynomial rings on algebras. *Monatsh. Math.* 183(1):177–189. DOI: [10.1007/s00605-016-0951-8](https://doi.org/10.1007/s00605-016-0951-8).
- [13] Peruginelli, G., Werner, N. J. (2018). Decomposition of integer-valued polynomial algebras. *J. Pure Appl. Algebra* 222(9):2562–2579. DOI: [10.1016/j.jpaa.2017.10.007](https://doi.org/10.1016/j.jpaa.2017.10.007).
- [14] Werner, N. J. (2010). Integer-valued polynomials over quaternion rings. *J. Algebra* 324(7):1754–1769. DOI: [10.1016/j.jalgebra.2010.06.024](https://doi.org/10.1016/j.jalgebra.2010.06.024).
- [15] Werner, N. J. (2012). Integer-valued polynomials over matrix rings. *Commun. Algebra* 40(12):4717–4726. DOI: [10.1080/00927872.2011.606859](https://doi.org/10.1080/00927872.2011.606859).