

Silver dichotomy for countable cofinalities

Vincenzo Dimonte

September 11, 2018

Joint work with Xianghui Shi

Previously...

When λ is a strong limit cardinal of cofinality ω , descriptive set theory can be done in ${}^\lambda 2$

When λ is a strong limit cardinal of cofinality ω , descriptive set theory can be done in ${}^\lambda 2$, or equivalently in ${}^\omega \lambda$

When λ is a strong limit cardinal of cofinality ω , descriptive set theory can be done in ${}^\lambda 2$, or equivalently in ${}^\omega \lambda$, $\prod_{n \in \omega} \lambda_n$

When λ is a strong limit cardinal of cofinality ω , descriptive set theory can be done in ${}^\lambda 2$, or equivalently in ${}^\omega \lambda$, $\prod_{n \in \omega} \lambda_n$ or $V_{\lambda+1}$

When λ is a strong limit cardinal of cofinality ω , descriptive set theory can be done in ${}^\lambda 2$, or equivalently in ${}^\omega \lambda$, $\prod_{n \in \omega} \lambda_n$ or $V_{\lambda+1}$.

Many results in classical descriptive set theory hold also in this setting

When λ is a strong limit cardinal of cofinality ω , descriptive set theory can be done in ${}^\lambda 2$, or equivalently in ${}^\omega \lambda$, $\prod_{n \in \omega} \lambda_n$ or $V_{\lambda+1}$.

Many results in classical descriptive set theory hold also in this setting.

In general, the results that are dependent to some tree-structure generalize very well

When λ is a strong limit cardinal of cofinality ω , descriptive set theory can be done in ${}^\lambda 2$, or equivalently in ${}^\omega \lambda$, $\prod_{n \in \omega} \lambda_n$ or $V_{\lambda+1}$.

Many results in classical descriptive set theory hold also in this setting.

In general, the results that are dependent to some tree-structure generalize very well.

$I_0(\lambda)$ has an influence on this setting in the same way that AD has an influence on classical descriptive set theory

When λ is a strong limit cardinal of cofinality ω , descriptive set theory can be done in ${}^\lambda 2$, or equivalently in ${}^\omega \lambda$, $\prod_{n \in \omega} \lambda_n$ or $V_{\lambda+1}$.

Many results in classical descriptive set theory hold also in this setting.

In general, the results that are dependent to some tree-structure generalize very well.

$I_0(\lambda)$ has an influence on this setting in the same way that AD has an influence on classical descriptive set theory.

Theorem (Silver, *Counting the number of equivalence classes of borel and coanalytic equivalence relations*. 1980)

Let X be a Polish space and $E \subseteq X^2$ be a coanalytic equivalence relation on X . Then exactly one of the following holds:

- E has at most countably many classes;
- there is a continuous injection $\varphi : {}^\omega 2 \rightarrow X$ such that for distinct $x, y \in {}^\omega 2$ $\neg \varphi(x) E \varphi(y)$.

Is this true also for the generalized Baire space?

Theorem (Friedman, Kulikov 2014)

Suppose $V = L$ and κ inaccessible. Then the order $\langle \mathcal{P}(\kappa), \subset \rangle$ can be embedded into the set of Borel equivalence relations on 2^κ strictly below the identity, ordered with Borel reducibility.

Theorem (Silver, 1980)

Let E be a coanalytic equivalence relation on ${}^\omega 2$. Then exactly one of the following holds:

- E has at most countably many classes;
- there is a continuous injection $\varphi : 2^\omega \rightarrow {}^\omega 2$ such that for distinct $x, y \in 2^\omega$ $\neg \varphi(x) E \varphi(y)$.

Theorem?

Let E be a coanalytic equivalence relation on $\prod_{n \in \omega} \lambda_n$. Then exactly one of the following holds:

- E has at most countably many classes;
- there is a continuous injection $\varphi : \prod_{n \in \omega} \lambda_n \rightarrow \prod_{n \in \omega} \lambda_n$ such that for distinct $x, y \in \prod_{n \in \omega} \lambda_n$ $\neg \varphi(x) E \varphi(y)$.

Theorem?

Let E be a coanalytic equivalence relation on $\prod_{n \in \omega} \lambda_n$. Then exactly one of the following holds:

- E has at most λ many classes;
- there is a continuous injection $\varphi : \prod_{n \in \omega} \lambda_n \rightarrow \prod_{n \in \omega} \lambda_n$ such that for distinct $x, y \in \prod_{n \in \omega} \lambda_n$ $\neg \varphi(x) E \varphi(y)$.

Theorem! (D.-Shi)

Let λ_n be measurable cardinals. Let E be a coanalytic equivalence relation on $\prod_{n \in \omega} \lambda_n$. Then exactly one of the following holds:

- E has at most λ many classes;
- there is a continuous injection $\varphi : \prod_{n \in \omega} \lambda_n \rightarrow \prod_{n \in \omega} \lambda_n$ such that for distinct $x, y \in \prod_{n \in \omega} \lambda_n$ $\neg \varphi(x) E \varphi(y)$.

A theorem by Shelah appears!

“Definition”

Let E be an equivalence relation on some product space

“Definition”

Let E be an equivalence relation on some product space. We say that E has the “singleton property” if for all x, y , if they differ *only* in one coordinate, then $\neg xEy$

“Definition”

Let E be an equivalence relation on some product space. We say that E has the “singleton property” if for all x, y , if they differ *only* in one coordinate, then $\neg xEy$.

Theorem (Shelah, *Can the fundamental (homotopy) group of a space be the rationals?* 1988)

If E is a co-analytic equivalence relation on ${}^\omega 2$ with the singleton property, then there is a continuous injection $\varphi : {}^\omega 2 \rightarrow {}^\omega 2$ such that for distinct $x, y \in {}^\omega 2$ $\neg \varphi(x)E\varphi(y)$.

“Definition”

Let E be an equivalence relation on some product space. We say that E has the “singleton property” if for all x, y , if they differ *only* in one coordinate, then $\neg xEy$.

Theorem (Shelah *On nice equivalence relations on λ^2* 2004)

Let λ_n be measurable cardinals. If E is a co-analytic equivalence relation on $\prod_{n \in \omega} \lambda_n$ with the singleton property, then there is a continuous injection $\varphi : \prod_{n \in \omega} \lambda_n \rightarrow \prod_{n \in \omega} \lambda_n$ such that for distinct $x, y \in \prod_{n \in \omega} \lambda_n$ $\neg \varphi(x)E\varphi(y)$.

G_0 -dichotomy

Fix a dense subset S of ${}^{<\omega}2$ that intersects every level in exactly one element

Fix a dense subset S of ${}^{<\omega}2$ that intersects every level in exactly one element. Let G_0 be the directed graph that couples two elements if they start with an element of S and differ only in the next coordinate

Fix a dense subset S of ${}^{<\omega}2$ that intersects every level in exactly one element. Let G_0 be the directed graph that couples two elements if they start with an element of S and differ only in the next coordinate.

Theorem (G_0 -dichotomy, (maybe Kechris-Solecki-Todorćević *Borel chromatic numbers*, 1999))

Let G be an analytic directed graph on ${}^\omega 2$. Then exactly one of the following holds

Fix a dense subset S of ${}^{<\omega}2$ that intersects every level in exactly one element. Let G_0 be the directed graph that couples two elements if they start with an element of S and differ only in the next coordinate.

Theorem (G_0 -dichotomy, (maybe Kechris-Solecki-Todorćević *Borel chromatic numbers*, 1999))

Let G be an analytic directed graph on ${}^\omega 2$. Then exactly one of the following holds:

- there is a (Borel) \aleph_0 -colouring of G

Fix a dense subset S of ${}^{<\omega}2$ that intersects every level in exactly one element. Let G_0 be the directed graph that couples two elements if they start with an element of S and differ only in the next coordinate.

Theorem (G_0 -dichotomy, (maybe Kechris-Solecki-Todorćević *Borel chromatic numbers*, 1999))

Let G be an analytic directed graph on ${}^\omega 2$. Then exactly one of the following holds:

- there is a (Borel) \aleph_0 -colouring of G ;
- there is a continuous function from ${}^\omega 2$ to itself that is a homomorphism from G_0 to G

Fix a dense subset S of ${}^{<\omega}2$ that intersects every level in exactly one element. Let G_0 be the directed graph that couples two elements if they start with an element of S and differ only in the next coordinate.

Theorem (G_0 -dichotomy, (maybe Kechris-Solecki-Todorćević *Borel chromatic numbers*, 1999))

Let G be an analytic directed graph on ${}^\omega 2$. Then exactly one of the following holds:

- there is a (Borel) \aleph_0 -colouring of G ;
- there is a continuous function from ${}^\omega 2$ to itself that is a homomorphism from G_0 to G .

This actually generalizes nicely, with almost the same proof.

Fix a dense subset S of $\prod_{n \in \omega} \lambda_n$ that intersects every level in exactly one element. Let G_0 be the directed graph that couples two elements if they start with an element of S and differ only in the next coordinate.

Theorem?

Let G be an analytic directed graph on $\prod_{n \in \omega} \lambda_n$. Then exactly one of the following holds:

- there is a \aleph_0 -colouring of G ;
- there is a continuous function from $\prod_{n \in \omega} \lambda_n$ to itself that is a homomorphism from G_0 to G .

This actually generalizes nicely, with almost the same proof.

Fix a dense subset S of $\prod_{n \in \omega} \lambda_n$ that intersects every level n in exactly κ_{n-1} element. Let G_0 be the directed graph that couples two elements if they start with an element of S and differ only in the next coordinate.

Theorem?

Let G be an analytic directed graph on $\prod_{n \in \omega} \lambda_n$. Then exactly one of the following holds:

- there is a \aleph_0 -colouring of G ;
- there is a continuous function from $\prod_{n \in \omega} \lambda_n$ to itself that is a homomorphism from G_0 to G .

This actually generalizes nicely, with almost the same proof.

Fix a dense subset S of $\prod_{n \in \omega} \lambda_n$ that intersects every level n in exactly κ_{n-1} element. Let G_0 be the directed graph that couples two elements if they start with an element of S and differ only in the next coordinate.

Theorem?

Let G be an analytic directed graph on $\prod_{n \in \omega} \lambda_n$. Then exactly one of the following holds:

- there is a λ -colouring of G ;
- there is a continuous function from $\prod_{n \in \omega} \lambda_n$ to itself that is a homomorphism from G_0 to G .

This actually generalizes nicely, with almost the same proof.

Fix a dense subset S of $\prod_{n \in \omega} \lambda_n$ that intersects every level n in exactly κ_{n-1} element. Let G_0 be the directed graph that couples two elements if they start with an element of S and differ only in the next coordinate.

Theorem! (D.-Shi)

Let G be an analytic directed graph on $\prod_{n \in \omega} \lambda_n$. Then exactly one of the following holds:

- there is a λ -colouring of G (actually, something more complicated, but equivalent for graphs that are the complement of an equivalence relation);
- there is a continuous function from $\prod_{n \in \omega} \lambda_n$ to itself that is a homomorphism from G_0 to G .

This actually generalizes nicely, with almost the same proof.

Now, let E be a co-analytic equivalence relation on $\prod_{n \in \omega} \lambda_n$

Now, let E be a co-analytic equivalence relation on $\prod_{n \in \omega} \lambda_n$. Then its complement G is an analytic directed graph, therefore either E has $\leq \lambda$ equivalence classes

Now, let E be a co-analytic equivalence relation on $\prod_{n \in \omega} \lambda_n$. Then its complement G is an analytic directed graph, therefore either E has $\leq \lambda$ equivalence classes, or there is a continuous function $\varphi : \prod_{n \in \omega} \lambda_n \rightarrow \prod_{n \in \omega} \lambda_n$ such that $x, y \in G_0$ iff $\neg \varphi(x) E \varphi(y)$

Now, let E be a co-analytic equivalence relation on $\prod_{n \in \omega} \lambda_n$. Then its complement G is an analytic directed graph, therefore either E has $\leq \lambda$ equivalence classes, or there is a continuous function $\varphi : \prod_{n \in \omega} \lambda_n \rightarrow \prod_{n \in \omega} \lambda_n$ such that $x, y \in G_0$ iff $\neg \varphi(x) E \varphi(y)$. The problem is now that φ is possibly not injective

Now, let E be a co-analytic equivalence relation on $\prod_{n \in \omega} \lambda_n$. Then its complement G is an analytic directed graph, therefore either E has $\leq \lambda$ equivalence classes, or there is a continuous function $\varphi : \prod_{n \in \omega} \lambda_n \rightarrow \prod_{n \in \omega} \lambda_n$ such that $x, y \in G_0$ iff $\neg \varphi(x) E \varphi(y)$. The problem is now that φ is possibly not injective.

Classically, from the G_0 -dichotomy to Silver Dichotomy we use the meagre-comeagre structure of ${}^\omega 2$

Now, let E be a co-analytic equivalence relation on $\prod_{n \in \omega} \lambda_n$. Then its complement G is an analytic directed graph, therefore either E has $\leq \lambda$ equivalence classes, or there is a continuous function $\varphi : \prod_{n \in \omega} \lambda_n \rightarrow \prod_{n \in \omega} \lambda_n$ such that $x, y \in G_0$ iff $\neg \varphi(x) E \varphi(y)$. The problem is now that φ is possibly not injective.

Classically, from the G_0 -dichotomy to Silver Dichotomy we use the meagre-comeagre structure of ${}^\omega 2$. This creates many problems in $\prod_{n \in \omega} \lambda_n$

Now, let E be a co-analytic equivalence relation on $\prod_{n \in \omega} \lambda_n$. Then its complement G is an analytic directed graph, therefore either E has $\leq \lambda$ equivalence classes, or there is a continuous function $\varphi : \prod_{n \in \omega} \lambda_n \rightarrow \prod_{n \in \omega} \lambda_n$ such that $x, y \in G_0$ iff $\neg \varphi(x) E \varphi(y)$. The problem is now that φ is possibly not injective.

Classically, from the G_0 -dichotomy to Silver Dichotomy we use the meagre-comeagre structure of ${}^\omega 2$. This creates many problems in $\prod_{n \in \omega} \lambda_n$, but Shelah's theorem can save us: the complement of G_0 has the singleton property, and we can use a similar argument to finally prove the Silver Dichotomy.

A look into the future...

Can we get rid of the measurable cardinals?

Can we get rid of the measurable cardinals?

Are measurable cardinals the key to understand the Baire structure of ${}^\lambda 2$?

One of the main points of the Axiom of Determinacy is that it generalizes regularity properties for all subsets of reals

One of the main points of the Axiom of Determinacy is that it generalizes regularity properties for all subsets of reals. This is true also for Silver Dichotomy:

Theorem (AD, Hjorth, *A dichotomy for the definable universe*, 1995)

Let E be an equivalence relation on ${}^\omega 2$. Then exactly one of the following holds:

- the classes of E are well-ordered;
- there is a continuous injection $\varphi : {}^\omega 2 \rightarrow {}^\omega 2$ such that for distinct $x, y \in {}^\omega 2$ $\neg \varphi(x) E \varphi(y)$.

One of the main points of I0 is that it generalizes AD-like results to higher cardinal

One of the main points of I_0 is that it generalizes AD-like results to higher cardinal. Does it work also in this case?

Open problem $I_0(\lambda)$

Let E be an equivalence relation on ${}^\lambda 2$. Is it true that exactly one of the following holds?

- the classes of E are well-ordered;
- there is a continuous injection $\varphi : {}^\lambda 2 \rightarrow {}^\lambda 2$ such that for distinct $x, y \in {}^\lambda 2$ $\neg \varphi(x) E \varphi(y)$.

Forbidden slide 1 (not enough time)

Brief summary of proof of Shelah's result.

Consider the double diagonal Prikry forcing \mathbb{P} that adds *two* Prikry sequences in λ . This forcing has two important characteristics:

- if M is a model of cardinality λ , then there is a M -generic set for \mathbb{P} in V ;
- only the tails of the generic are meaningful, so changing just one coordinate maintain the genericity.

Forbidden slide 2 (not enough time)

The fact that E is co-analytic is also important: this means that the formula that defines E is absolute between models that contain V_λ .

So the proof goes like this: pick M small model that contains everything. If there is a condition of \mathbb{P} that forces that the two elements of the generic are E -related, then also those in V are E -related. Switching one coordinate we do the same, but this contradicts the singleton property or the fact that E is an equivalence relation.

Using generic absoluteness, we have a partial result:

Theorem

Suppose $I_0(\lambda)$, as witness by j , and let $(\lambda_n)_{n \in \omega}$ be the critical sequence of j

Using generic absoluteness, we have a partial result:

Theorem

Suppose $I_0(\lambda)$, as witness by j , and let $(\lambda_n)_{n \in \omega}$ be the critical sequence of j . Suppose that all subsets of $V_{\lambda+1}$ are $U(j)$ -representable

Using generic absoluteness, we have a partial result:

Theorem

Suppose $I_0(\lambda)$, as witness by j , and let $(\lambda_n)_{n \in \omega}$ be the critical sequence of j . Suppose that all subsets of $V_{\lambda+1}$ are $U(j)$ -representable. Then if $E \in L(V_{\lambda+1})$ is an equivalence relation with the singleton property, there is a continuous injection $\prod_{n \in \omega} \lambda_n \rightarrow \prod_{n \in \omega} \lambda_n$ such that for distinct $x, y \in \prod_{n \in \omega} \lambda_n$ $\neg \varphi(x) E \varphi(y)$.

Thanks for watching.